

Physiology
of
Fungi

VINCENT W COCHRANE

Professor of Biology

Wesleyan University

Physiology
of
Fungi



NEW YORK JOHN WILEY & SONS INC

LONDON CHAPMAN & HALL LIMITED

COPYRIGHT © 1958 BY JOHN WILEY & SONS INC

All rights reserved. This book or any part thereof
must not be reproduced in any form with-
out the written permission of the publisher

LIBRARY OF CONGRESS CATALOG CARD NUMBER 58-13456
PRINTED IN THE UNITED STATES OF AMERICA

Preface

Although study of the physiology of the fungi has not kept pace with that of for example bacterial metabolism it has in recent decades had much the same development. It seems fair to say that there have been two disparate types of study. Research on some problems—reproduction parasitism development—has been with significant but not very numerous exceptions descriptive. At the other extreme it has been possible to develop basic biochemical knowledge on the implicit assumption that the microbial cell is a small bag of enzymes which only require to be extracted and enumerated. Our need now is to extend the valuable results of the abstract analytical study to the type of problem that has remained in the descriptive natural history stage.

This however is for the future. We work with what we have and it is the purpose of this book to summarize and organize our knowledge of the physiology of the fungi indicating whenever possible some of the problems that lie immediately ahead. If there is one guiding principle it is that the fungi however unusual or unique they may be as functioning intact organisms can be studied at the cellular and molecular levels only by taking account of the findings of other disciplines particularly bacterial physiology. This is the point of view of comparative biochemistry. No doubt in a decade or two some more useful principle of organization will emerge but for the present an

COPYRIGHT © 1958 BY JOHN WILEY & SONS, INC.

All rights reserved. This book or any part thereof
must not be reproduced in any form without the
written permission of the publisher.

LIBRARY OF CONGRESS CATALOG CARD NUMBER 58-13456
PRINTED IN THE UNITED STATES OF AMERICA

In the naming of organisms I have followed the usage of the original author unless confusion is thereby introduced. A few exceptions to this rule have been made e.g. in the use of *Streptomyces* for the aerobic actinomycetes forming conidia in chains.

Two important and interesting topics have been almost entirely omitted—the genetics of fungi and the interactions of fungi with other organisms in nature. Both of these subjects deserve more space than could here be given to them and the second—the ecology of fungi—probably requires a type of conceptual approach different from that of this book.

Individual chapters have been read in manuscript by S. H. Hutner, Frederick Kavanagh, Albert Kelner, Leonard Machlis, S. E. A. McCallan, W. C. Snyder and C. E. Yarwood. It is a pleasure to acknowledge their advice and help. I am grateful too to the many authors and publishers who have allowed me to use published material and to the staffs of the Olin Library of Wesleyan University, the Mann Library of Cornell University and the New York Agricultural Experiment Station Library.

A considerable part of the work embodied here was done during a sabbatical leave spent as guest of the Department of Plant Pathology of the University of California at Berkeley to the members of which I am greatly indebted. The Board of Trustees of Wesleyan University generously provided support for this period.

Jean Conn Cochrane has given invaluable assistance in all phases of preparation of the manuscript which without her help would have been poorer in quality and later in appearance.

VINCENT W. COCHRANE

October 1958

ordering of knowledge around this central principle seems valid

To stress the fundamental biochemical similarities of all organisms is to assert in a concrete instance the general proposition of the unity of all living things the unity of the science of biology. The separation of the various fields of biology has caused incalculable damage to all possibly most to the botanical sciences. This is not to deny the need for specialization within biology but when this specialization reaches the point where a student can for example be considered well trained in botany without having had any contact with animal sciences or with bacteriology it seems time to reconsider. Some way must be found both at the teaching and at the research level to ensure that biologists of all sorts will be conversant with each other's problems and will be able to draw on each other's experience in the prosecution of their own specialties.

The literature on the physiology of fungi is very large and is growing rapidly. Much of it is of necessity repetitious and limited in scope to particular organisms or to practical applications. Some of it fails to meet criteria of accuracy or logic. I have attempted to exercise the author's prerogative of selection so as to exclude work which in my judgment falls into either category. However I have preferred to err on the side of citing too much literature rather than too little. All inclusiveness is of course out of the question the references are designed to give immediately most of the work of the last three decades and to open up the entire literature for the student or research worker. In this connection it is to be regretted that the old fashioned review of the literature has been largely squeezed out of modern journal articles. No doubt it was often overdone and wasted valuable space but the current practice of publishing the past month's work in the form of a letter to the editor has even less to recommend it.

Professor Frank Dickens once spoke in a seminar to the effect that for the purposes of biochemistry the yeasts are considered to be animals. With something of the sort in mind I have omitted much of the physiological work on yeasts feeling that it has been well treated elsewhere and that its importance to the filamentous fungi is of a background nature only. On the other hand I have included the aerobic actinomycetes on the ground that ecologically and physiologically they have close affinities to the true fungi. Both groups are filamentous both require oxygen for growth both form exogenous spores and both occur in similar habitats. The scope of the book is then the filamentous fungi and actinomycetes and that work on the yeasts which has a direct bearing on the physiology of the true fungi.

Contents

	<i>Page</i>
<i>Chapter 1 Cultivation and Growth</i>	<i>1</i>
1 Methods of Cultivation	4
2 Preservation	4
3 The Dynamics of Growth	5
4 Dimorphism	10
5 The Measurement of Growth	11
6 Temperature and Growth	13
7 pH and Growth	19
8 Oxygen and Growth	23
9 Carbon Dioxide and Growth	24
10 Water and Growth	24
11 Bioluminescence	26
Bibliography	27
 <i>2 The Composition of Fungus Cells</i>	 <i>35</i>
1 The Major Constituents of the Cell	36
2 Mineral Constituents of the Cell	37
3 The Cell Wall	39
4 Cellular Carbohydrates	42
5 Nitrogenous Constituents	43
6 Cell Lipids	45
Bibliography	49

	<i>Page</i>
<i>Chapter 7 Respiration</i>	<i>202</i>
1 Problems of Method	203
2 Factors in Respiration	206
3 Endogenous Respiration	210
4 Oxidative Assimilation	212
5 The Embden Meyerhof Pathway	212
6 The Phosphogluconate Oxidation Pathway	217
7 The Non phosphorylative Oxidation of Glucose	220
8 Pentose Fermentation and Oxidation	221
9 The Citric Acid Cycle	222
10 Dicarboxylic Acids in Respiration	226
11 Carbon Dioxide Fixation	228
12 Terminal Respiration	229
Bibliography	232
 <i>8 Nitrogen Nutrition and Metabolism</i>	 <i>241</i>
1 Inorganic Sources of Nitrogen	242
2 Proteins and Peptides	250
3 Utilization of Amino Acids and Amides	256
4 Biosynthesis of Amino Acids	265
5 Amines	273
6 Metabolism of Nucleic Acids and Their Derivatives	274
7 Urea	277
8 Ergot Alkaloids	279
9 Miscellaneous Nitrogen Compounds	280
Bibliography	283
 <i>9 Inorganic Nutrition and Metabolism</i>	 <i>300</i>
1 Phosphorus	301
2 Potassium	303
3 Sulfur	304
4 Magnesium	306
5 Iron	307
6 Zinc	308
7 Copper	309
8 Manganese	310
9 Molybdenum	310
10 Calcium	311
11 Cobalt	312
12 Other Inorganic Nutrients	313
13 Bioassay of Minerals	313
14 Problems in Mineral Nutrition	314
Bibliography	317

Chapter 3 Carbon Nutrition**Page**
55

- 1 Problems of Method 55
- 2 Monosaccharides and Their Derivatives 60
- 3 Oligosaccharides 66
- 4 Polysaccharides 75
- 5 Organic Acids 80
- 6 The Lipids 82
- 7 Other Carbon Sources 83
- 8 The Concentration Factor in Carbon Nutrition 85
- 9 Enzyme Induction Adaptive Growth and the Utilization of Mixed Carbon Sources 86
- 10 Efficiency of Growth 89
- 11 Conclusion 90
- Bibliography 92

4 Carbon Metabolism**99****PART I INTRODUCTION**

- Bibliography 103

5 Carbon Metabolism**104****PART II THE METABOLISM OF CARBOHYDRATES AND RELATED COMPOUNDS**

- 1 The Metabolism of Monosaccharides 104
- 2 The Metabolism of Oligosaccharides 105
- 3 The Metabolism of Other Glycosides 109
- 4 The Metabolism of Polysaccharides 110
- 5 The Metabolism of Amino Sugars 121
- 6 The Metabolism of Polyhydric Alcohols 122
- Bibliography 124

6 Carbon Metabolism**131****PART III ORGANIC ACIDS LIPIDS AND OTHER COMPOUNDS**

- 1 The Metabolism of Aliphatic Organic Acids 131
- 2 The Metabolism of Lipids 148
- 3 The Metabolism of Aromatic Compounds 161
- 4 Derivatives of Pyrone 178
- 5 Derivatives of Furan 180
- 6 Biological Methylation 181
- 7 Biological Chlorination 183
- 8 Penicillin 181
- Bibliography 187

6	Nutrition and Germination	400
7	pH and Germination	403
8	Water and Germination	404
9	Temperature and Germination	407
10	Oxygen and Germination	412
11	Carbon Dioxide and Germination	413
12	Biological Effects on Spore Germination	413
	Bibliography	415

Chapter 13 The Action of Physical Agents 423

1	High Temperature	423
2	Low Temperature	425
3	Visible Light	427
4	Ultraviolet Radiation	429
5	Ionizing Radiations	433
6	Electromagnetic Radiation of Long Wavelength	435
	Bibliography	435

14 The Action of Chemical Agents 440

1	The Response to Toxicants and Its Measurement	441
2	Tolerance and Resistance	448
3	Selective Accumulation	450
4	Surface Action	453
5	Hydrogen Ion Concentration	454
6	Structure and Activity	457
7	The Action of Heavy Metals	462
8	The Action of Sulfur	472
9	Quinones and Other Unsaturated Compounds	475
10	Chelation and Fungistatic Activity	478
	Bibliography	482

<i>Organism Index</i>	499
-----------------------	-----

<i>Subject Index</i>	503
----------------------	-----

	<i>Page</i>
<i>Chapter 10 Vitamin Requirements of Fungi</i>	<i>323</i>
1 Problems of Method	321
2 The Nature of Vitamin Requirements	325
3 The Metabolism of Vitamins	328
4 Fungi in the Bioassay of Vitamins	328
5 Thiamine	329
6 Biotin	333
7 Pyridoxine	335
8 Riboflavin	336
9 Nicotinic Acid	337
10 Inositol	339
11 Para aminobenzoic Acid	340
12 Pantothenic Acid	341
13 Pteroylglutamic Acid	342
14 The Vitamins B ₁₂	343
15 Vitamins K	344
16 Other Growth Factors	344
17 The Occurrence of Vitamin Deficiencies in the Fungi	345
18 Vitamins Taxonomy and Ecology	346
Bibliography	347
<i>11 Reproduction</i>	<i>356</i>
1 Light	357
2 Temperature	362
3 Water Relations	364
4 Oxygen Carbon Dioxide and Volatile Substances	365
5 Acidity	366
6 Carbon Nutrition	367
7 Nitrogen Nutrition	369
8 Mineral Nutrition	370
9 Vitamins and Unidentified Factors	371
10 Specific Reproductive Hormones	373
11 Metabolism and Reproduction	375
12 The Sporulation of Plant Pathogens	377
13 Conclusions	378
Bibliography	380
<i>12 Spore Germination</i>	<i>387</i>
1 Measurement	388
2 The Germination Process	389
3 Spore Metabolism	392
4 Maturation and Dormancy	394
5 Spore Longevity	398

1 Cultivation and Growth

The precise definition of growth in fungi depends on the method of measurement used. Most methods determine directly or indirectly the increase in mass of an inoculum after a known incubation in a complete medium. Increase in mass or in number of cells may therefore be used as an operational definition of growth.

In considering cultivation and growth we shall first concern ourselves with methods of cultivation and preservation of fungi, the dynamics of growth, dimorphism, and the measurement of growth. Then we turn to external factors—temperature, hydrogen ion concentration, oxygen, carbon dioxide, and water—as they affect growth. The same external conditions also influence spore germination (Chapter 12). Radiant energy effects are taken up in Chapter 13.

1 METHODS OF CULTIVATION

Cultures of fungi are usually maintained on conventional agar slants, and the agar petri dish has been employed in experimental work. Most quantitative studies, however, require a liquid medium which may be either surface (still) or submerged (aerated) culture.

The general technique of surface culture is too well known to require description here. It is not always realized that normal aerobic growth can be obtained only if the inoculum floats on the surface of the liquid. Many spores float naturally, for those that do not use of a bran cultivated inoculum (206) or inclusion of a small amount of agar

for *Ustilago zeae* and 56 mM per liter per hour for *Aspergillus niger* (97). Rates of aeration of this magnitude require an apparatus in which a sterile medium can be agitated and aerated. Various laboratory scale devices have been described (10 55 142 168 252) and some are commercially available. Theoretical and practical problems of design have been studied in several laboratories (9 49 68 97).

Growth on solid media other than agar is useful in special problems such materials include soil (224 260) sawdust (6) bran (206) and cotton (25). The use of plant materials sterilized by treatment with propylene oxide is particularly valuable in the induction of sporulation (Chapter 11) and may have wider application in the study of physiology.

The inoculum used for cultures is usually a population of spores from an agar slant. A common occurrence in the fungi is the replacement of the normal type by a variant arising in culture; this development has been analyzed especially with regard to sporulation. Other examples include the loss of biochemical capacities (63) change in virulence to a host plant (202) and change in assimilatory capacity (246). The ultimate recourse is to preservation of cultures in a dormant state (p. 4). Methods of obtaining single spore cultures of fungi are reviewed by Hildebrand (136); such cultures may be necessary in order to recover or purify the desired type. It must however be remembered that some spores are multinucleate (Chapter 12) and may contain genetically different nuclei.

Three physiological aspects of the inoculation process may be mentioned. First growth is more rapid with a heavy than with a light inoculum; in addition a heavy inoculation of a submerged culture usually results in the development of small colonies with less danger that oxygen diffusion into the colony will limit growth or respiration. Second the use of old spores of low viability is to be avoided; both the age and the conditions of growth of the inoculum, especially medium and temperature, should be standardized. For certain fungi the chemostat provides an easily standardized inoculum (221). Finally a large inoculum may be required in order to initiate growth on an unfavorable medium (235).

When rapid growth is a desideratum the inoculum may be pregerminated spores or mycelium (207). Mechanically macerated mycelium can be used as inoculum (233 259). This is often the method of choice for fungi which do not sporulate (3). The treatment to minimize injury should be as brief as possible.

Special methods of cultivating particular fungi are too numerous to mention here. Continuous flow apparatus may have some general

gum arabic or polyvinyl alcohol in the medium (334) may be necessary

Two major objections attach to the use of *surface culture for fungi*. The cells of the developed mat are not only visibly of different types—aerial and submerged growth—but also are exposed to different environmental conditions. Aerial hyphae are remote from the nutrient supply and are bathed in the atmosphere; the lower cells, by contrast, are in close contact with nutrients and with soluble metabolic products but suffer from a partially anaerobic environment (167). Examination of submerged hyphae in such a culture shows them to be highly vacuolated. The culture as a whole is probably always deficient in oxygen. This deficiency is shown by the favorable effect on sugar utilization of an increase in surface: volume ratio (208) and by the effect mentioned below of aeration on growth rate of most fungi.

The commonest and most useful laboratory-scale aerated culture apparatus is the shaker, first thoroughly studied by Kluwer and Perquin (161). Flasks are mounted on a platform which is either rotated rapidly or moved horizontally with a reciprocating motion. Problems of design are considered by Shu and Johnson (270), Paladino (230), and Chain and Gualandri (51).

Representatives of all the major groups of fungi and of the genus *Streptomyces* have been grown successfully in stirred or shake culture. The growth rate is usually at least double that of the same organism in surface culture (1, 61, 162, 286, 319, 335), although some fungi do not show this effect (14, 268) and some grow so poorly that the method cannot be used for them. One of the chief advantages, apart from more rapid growth, is that shaker-grown mycelium is usually homogeneous enough to be used directly in respirometric and other metabolic studies.

In shake culture each colony is exposed to a uniform environment in all spatial directions; hence the typical colony is a globose structure. Colonial morphology varies, however, with the species—even within a genus—and with the medium (49). Sporulation is usually, although not always suppressed—another factor of homogeneity. Microscopic characteristics of the mycelium are also affected by aeration and agitation (76, 83).

Although a shaken culture has a measurably higher content of dissolved oxygen and rate of oxygen diffusion than a still culture (283), for certain purposes a still higher rate of oxygen supply may be desirable (249, 252). This usually arises when a high concentration of a readily utilizable carbon source is employed; the peak oxygen demand under specified conditions is 16 mM oxygen per liter per hour.

spp can be successfully lyophilized (281) Haskins and Anastasiou (126) report successful use of a modification of the lyophil technique Too rapid rehydration of vacuum dried uredospores is apparently injurious (265a)

For sporulating fungi lyophilization appears to be the best single method until more information is available however preservation of many fungi will have to be on an experimental basis Lyophilization is comparatively easy and apparatus is commercially available Preliminary studies (197) indicate that spray drying of fungus spores may also be feasible

3 THE DYNAMICS OF GROWTH

Growth cannot be precisely defined a provisional operational definition is increase in either mass of cells or number of cells Mass—dry weight usually—in the filamentous fungi may be deceptive inasmuch as it can represent in part the accumulation of polysaccharides lipids or wall materials without any increase in living protoplasm Conversely in *Neurospora crassa* protein synthesis appears to continue for a period of time after weight increase has ceased (7)

The time course of growth of filamentous fungi in liquid shaken cultures is exemplified by the data of Figure 1 Similar curves are obtained with surface cultures (2 10, 218) Typical curves are characterized by three major phases with transitions between them

- 1 A phase of no apparent growth
- 2 A phase of rapid and approximately linear growth
- 3 A phase of no net growth or of autolysis and decline in dry weight

It should not however be assumed that all data fit this scheme In particular a two phase growth curve in which an initial growth period is succeeded by a second with a leveling off of growth intervening has been reported (30) The second phase may represent polysaccharide synthesis only without increase in other cell components or it may depend on a mobilization of nitrogen from older hyphae and use of it for new growth The observations of Morton and Proadbert (205) on *Scopulariopsis brevicaulis* support the second explanation at least in a medium relatively high in carbon it appears that nitrogen is reutilized after exhaustion of the available exogenous nitrogen

Probably by suitable adjustment of cultural factors especially nutrient concentrations a variety of growth curves very different from those of Figure 1 can be obtained Provision of calcium for example

interest these include devices in which only the solution is renewed (325) and those in which both nutrients and cell population are controlled (138)

Culture media for fungi are described by Fred and Waksman (101) Levine (176) and Lilly and Barnett (180) In general it has been too readily assumed that fungi need and tolerate high nutrient levels and high acidity this is true of many fungi but by no means of all For certain purposes concentrated media are of course useful but morphology physiology and reproduction are best studied in more dilute media than those often used

2 PRESERVATION

Interest in the preservation of fungi springs from the desire to avoid excessive labor in maintaining stock cultures and more important the wish to maintain cultures with a minimum of genetic change The problem is to maintain a viable inoculum in a non growing condition

Cultures and herbarium specimens often survive for surprisingly long periods after drying probably by virtue of the formation of long lived spores Zobl (344) reviews early records of up to 21 year survival in his own work he found that survival of dried cultures is common in ascomycetes and basidiomycetes but the Mucorales are relatively short lived The sclerotia of several fungi which may live for as long as 13 years (330) are less viable if stored dry (294) A few of the more leathery basidiomycete sporophores survive dry for a period of years more succulent types are less durable (41 43)

In the soil tube method (5 121) a fungus is transferred to moist sterile soil preferably brought to pH 6-7 with calcium carbonate and grows and sporulates until the soil dries out The limited data available indicate that although some fungi survive at least 5 years in soil tubes too many do not live that long and the method therefore is to be regarded as a special one not adapted for general use

Preservation of cultures under sterile mineral oil is relatively simple and has the advantage that sporulation is not essential (39 267) Limited tests indicate however that although many fungi survive at least 2 years others die out earlier (146 331 333)

Lyophilization of spores suspended in serum gelatin dextran or other colloidal materials appears to be a usable method although again not successful with some fungi e.g. the Entomophthorales and the dermatophytes (199 239 265 337) Simple vacuum drying without freezing is suitable for many fungi (244) The structure of the fungus is important thus oogonia but not sporangia or mycelium of *Pythium*

occurs only at hyphal tips (272 273 280) Interior cells of the mycelium do not normally contribute to the new growth directly although they supply nutrients to peripheral cells especially to aerial structures in still culture Growth in any one time period is therefore a function not of the total number of cells present as in bacteria and yeasts but of the number of hyphal tips and of the rate at which these tips are supplied nutrients

Some of the chemical events during the phase of rapid growth are illustrated in Figure 3 these include especially utilization of carbohydrate nitrogen and phosphate Metabolic products e.g acids may or may not appear in the medium at this time Characteristically respiratory activity is at a maximum (Chapter 7)

Few studies have considered critically the relation of reproduction to growth phases casual observation and the data of Robinson (248) on *Sporodinia grandis* indicate that in still culture spores usually develop at the end of the phase of rapid growth spores may however appear earlier (209)

The third and last phase is as shown in Figure 3 characterized usually by a decline in mycelial weight and in the appearance of nitrogen and phosphate in the medium The loss in weight varies from negligible to extreme a common pattern is a rapid loss of weight for a




Figure 2 The growth of *Neurospora crassa* as a function of time Curve 1 logarithmic plot curve 2 cube root plot curve 3 linear plot Retrawn from Emerson (86) by permission of the Williams and Wilkins Company




Figure 1 The growth of *Fusarium solani* in an aerated medium. Open circles dry weight, closed circles optical density.

to *Streptomyces fradiae* delays the onset of the autolytic phase (132).

The growth rate varies not only between species but between different isolates or clones of the same species; this has been found to be true in both liquid and agar cultures (71, 88, 89, 130).

The first phase, that of no apparent growth, presumably has two components: a genuine lag phase before spore germination and a phase in which growth is occurring but is undetectable by the methods used. The details of this growth period have not been studied in the fungi. In bacteria it appears that growth, as measured by protein and nucleic acid synthesis, begins immediately with no true lag period (298).

The second phase is that of rapid growth. As Emerson (86) has shown, the cube root of the dry weight increases linearly with time over a major portion of the period (Figure 2). It should be stressed, however, that this relation holds only if no external factor, e.g., oxygen or nutrient concentration, is limiting to growth. Conversely, a curve conforming to that of Figure 2 indicates that the organism is growing unrestrictedly.

The morphological basis of mycelial development is that growth

occurs only at hyphal tips (272 273 280) Interior cells of the mycelium do not normally contribute to the new growth directly although they supply nutrients to peripheral cells especially to aerial structures in still culture Growth in any one time period is therefore a function not of the total number of cells present as in bacteria and yeasts but of the number of hyphal tips and of the rate at which these tips are supplied nutrients

Some of the chemical events during the phase of rapid growth are illustrated in Figure 3 these include especially utilization of carbohydrate nitrogen and phosphate Metabolic products e.g. acids may or may not appear in the medium at this time Characteristically respiratory activity is at a maximum (Chapter 7)

Few studies have considered critically the relation of reproduction to growth phases casual observation and the data of Robinson (248) on *Sporodinia grandis* indicate that in still culture spores usually develop at the end of the phase of rapid growth spores may however appear earlier (209)

The third and last phase is as shown in Figure 3 characterized usually by a decline in mycelial weight and in the appearance of nitrogen and phosphate in the medium The loss in weight varies from negligible to extreme a common pattern is a rapid loss of weight for a




Figure 2 The growth of *Neurospora crassa* as a function of time. Curve 1 logarithmic plot; curve 2 cube root plot; curve 3 linear plot. Redrawn from Emerson (86) by permission of the Williams and Wilkins Company.




Figure 1 The growth of *Fusarium solani* in an aerated medium. Open circles dry weight; closed circles optical density.

to *Streptomyces fradiae* delays the onset of the autolytic phase (132).

The growth rate varies not only between species but between different isolates or clones of the same species; this has been found to be true in both liquid and agar cultures (71, 88, 89, 130).

The first phase, that of no apparent growth, presumably has two components: a genuine lag phase before spore germination and a phase in which growth is occurring but is undetectable by the methods used. The details of this growth period have not been studied in the fungi. In bacteria it appears that growth, as measured by protein and nucleic acid synthesis, begins immediately with no true lag period (298).

The second phase is that of rapid growth. As Emerson (86) has shown, the cube root of the dry weight increases linearly with time over a major portion of the period (Figure 2). It should be stressed, however, that this relation holds only if no external factor, e.g., oxygen or nutrient concentration, is limiting to growth. Conversely, a curve conforming to that of Figure 2 indicates that the organism is growing unrestrictedly.

The morphological basis of mycelial development is that growth

media and ammonia in high nitrogen media are the compounds most frequently involved

In better balanced and more dilute media the principal factor in cessation of growth is exhaustion of the carbohydrate supply (15 74 79) as shown by the data of Figure 3

The extent of autolytic breakdown once growth has ceased varies as mentioned above with species and cultural conditions Autolytic breakdown of *Aspergillus niger* mycelium is much more extensive in nitrate than in ammonium salt media possibly because of the alkalinity of the nitrate medium (15) However autolysis of a thermophilic *Streptomyces* sp is more rapid at acid than at neutral pH (158)

So much for growth in liquid media A fungus colony on an agar surface follows a different course Typically a period of little or no growth is succeeded by the establishment of a constant growth rate which is maintained unless toxic metabolites accumulate (36 93 308) The growth rate tube of Ryan et al (203) may be used to show this linear relation of extension to time the data of Figure 4 are typical Growth on soil should follow the same course if measured by horizontal spread only specialized rhizomorphs of *Armillaria mellea* grow several times as rapidly as unorganized mycelium (111)

Neurospora crassa and other fungi respond to certain toxicants e g




Figure 4 The linear growth of several fungi in growth rate tubes Curve 1 *Neurospora crassa* curve 2 *Rhizopus oryzae* curve 3 *R. nigricans* curve 4 *Sordaria fimicola* curve 5 *Fusarium oxysporum lycopersici* curve 6 *Monilia fruticola*




Figure 3 Growth of and changes in the medium by *Aspergillus oryzae* Redrawn from Crewther and Lennox (62) by permission of the *Australian Journal of Biological Sciences*

short time with no further change thereafter (151). Autolyzing mycelium of *Penicillium griseofulvum* undergoes an extensive breakdown of chitin carbohydrates and protein catalyzed by enzymes of the fungus (275). Other autolytic products include ammonia (71, 80, 160), free amino acids (71, 108), organic phosphorus compounds (200) and sulfur compounds (137, 245).

Cells at the end of the growth phase appear vacuolate; younger cells have a denser and more homogeneous protoplasm (49). Such old cells of course are found even during the growth period in regions remote from the growing hyphal tips.

From the rather limited data available the cessation of active growth appears to be determined by either of two factors. In concentrated media toxic metabolites can be shown to accumulate and these materials earlier called staling substances demonstrably retard growth (27, 37, 236). Probably organic acids in high-carbohydrate

sity of non mycelial growth phases is rendered more likely by the variety of stimuli which induce the phase in different fungi (Table 1)

Several theories reviewed elsewhere (152-262) have been proposed to explain dimorphism in the fungi pathogenic to animals. The most attractive concept is that synthesis of new protoplasm and cell division are separate processes. Filaments form when growth occurs under conditions not permitting rapid cell division. The studies of Nickerson and co workers (152-211-212-214-215-217) indicate that a supply of reduced organic sulphydryl compounds is essential for rapid cell division and hence for maintenance of the unicellular condition. At least in principle all the stimuli known to affect dimorphism in the pathogenic fungi could act through their effect on sulphydryl content of the cells. Whether this will apply to the non pathogenic fungi listed in Table 1 is not known.

The yeast phase of *Blastomyces dermatitidis* and *Paracoccidioides* (*Blastomyces*) *brasiliensis* is multinucleate that of some other dimorphic fungi uninucleate (82). The yeast and mycelial forms of *Blastomyces* spp. differ in their respiratory capacities (213).

Dimorphism in yeastlike fungi suggests of course that the true yeasts may have arisen by a more or less permanent acquisition of the unicellular habit. It is also perhaps legitimate to ask whether the filamentous habit in the true fungi and the algae is an expression of the same metabolic idiosyncrasy as the filamentous habit of *Blastomyces* and *Histoplasma*.

5 THE MEASUREMENT OF GROWTH

As mentioned earlier the meaning of growth is determined by the method of measurement chosen. Some methods are useful only for particular organisms or for special problems: no method is so general that it can be recommended for all.

The most widely used and within limits the most satisfactory measurement of growth is by determination of dry weight of the mycelium. The principal limitation is that weight may reflect accumulation of polysaccharides or other reserve materials rather than synthesis of new protoplasm. Oxidative assimilation (Chapter 7) is often very active in the fungi under conditions which restrict protein synthesis.

Various procedures may be used for the determination of dry weight. Often a coherent mycelium can be removed from a liquid medium, washed and dried in a tared container. More commonly filtration is required: tared filter paper protected from atmospheric moisture or porous discs (ceramic or sintered glass) held in a demountable suction

sorbose by a more colonial and less rapid growth on agar (293). The growing hyphal tips are killed and new branches then form and grow laterally (181).

4 DIMORPHISM

Several fungi pathogenic to man are found in infected tissues in a unicellular yeastlike form but when cultivated at room temperature grow out in a mycelial form. The term dimorphism strictly applies to these forms only but related phenomena in the normally filamentous fungi may provisionally be considered as in the same category. A dimorphic fungus is one therefore in which a reversible transformation from a mycelial to a non mycelial and unicellular growth type occurs.

Dimorphism is common in the yeasts (262) and unicellular bacteria may be caused experimentally to form filaments by for example magnesium deficiency (216-326). Some normally filamentous fungi e.g. smut fungi grow in culture as unicellular budding cells. The fungi listed in Table 1 include both the pathogenic dimorphic fungi

Table 1 Factors Affecting Non mycelial Growth in Fungi

Organism	Non mycelial Phase Favored by
<i>Ca. Aida albicans</i> <i>C. tropicalis</i>	Cysteine NH_4Cl glucose carbon source (152 211 214 217)
<i>Blastomyces dermatitidis</i>	High temperature (35-37 C) cultivation (177 256)
<i>Histoplasma capsulatum</i>	Cysteine, high temperature (255)
<i>Histoplasma farciminosum</i>	High CO_2 (40)
<i>Paracoccidioides brasiliensis</i>	High temperature (37 C) cultivation (82 250)
<i>Sporotrichum schenckii</i>	High CO_2 (81 254)
<i>Trichosporon capitatum</i>	Cysteine (217)
<i>Ceratostomella ulmi</i>	Aeration (75) low pH (26)
<i>Dematiaceae pullulans</i>	Low aeration (11) low carbohydrate (263)
<i>Fusarium oxysporum f. nicotianae</i>	Aeration (339)
<i>Mucor guilliermondii</i>	Low aeration (186)
<i>Ophiostoma multianellatum</i>	High (12 mg per liter) inositol (107)
<i>Ramanoa terricola</i>	Aeration (299)

and those filamentous fungi in which production of a unicellular growth phase can be induced reversibly. It should be stressed that in this second group the morphology of the unicellular phase may not be the same for all forms listed i.e. the available information does not discriminate between say conidia and hyphal fragments. The diver

We may conclude that the agar method is adequate for studies of some environmental factors e.g. temperature and (probably) composition of the atmosphere although even in these experiments the results are not always correlated with dry weight (91 134 145 250). It is decidedly not adequate for nutritional studies. Whether it should be used for essentially chemical problems like the effect of pH or toxic agents is still uncertain. The burden of proof that spread on agar is a fair index of growth rests on the investigator who chooses to use the method. Its value cannot be assumed.

6 TEMPERATURE AND GROWTH

Temperature affects growth: spore germination, reproduction and indeed all activities of the organism. Typical curves of growth as a function of temperature are shown in Figure 5. There is characteristically a linear portion in which growth increases directly with temperature, an optimum range which may be narrow or rather broad and a descending limb as the temperature becomes too high for growth. The curve as a whole is usually skewed to the right; an empirical method for converting this skewed curve to a straight line is described by Cohen and Yarwood (58).

The temperature growth curve tends to become more nearly symmetrical as the optimum becomes lower: fungi with optima of 22–24°C or less often have a much less skewed curve (53 98 143 171 311). Not all low temperature fungi, however, follow this pattern (113 164).

Fungi with high temperature optima generally have the typical skewed curve even more pronouncedly; in a few cases the high optimum is associated with a broader range of good growth but this is not the rule.

It should be noted that the concept of an optimum usually taken for granted is not in fact very clear. The response of *Phycomyces blakesleeenianus* to temperature (247) illustrates the principal ambiguities in the concept of optimum temperature. First growth is most rapid at 20–25°C but the maximum total amount of growth at limiting levels of thiamine is greatest at 10°. Second the effect of temperature on final dry weight can be almost wiped out if the nitrogen supply is made limiting.

Another example of the dependence of temperature characteristics on other factors has been elucidated by L. Fries (103). *Coprinus fimetarius* grows poorly at 44°C because of the failure of methionine biosynthesis to keep pace with other processes; if exogenous methionine is supplied growth at the elevated temperature is normal. The ap

assembly are practical filtration devices. Gelatinous growth which filters slowly may be separated by centrifugation. The use of solvents (16) to hasten drying is practicable of course only if the extracted lipids are negligible. Drying is usually effected by high temperature 70-80°C.

The dry weight of agar colonies although of dubious significance in most contexts has been determined by scraping or peeling off of the surface growth (91). Removal of agar with hot water results in a significant loss in dry weight (67).

Turbidimetric measurements of growth are suitable for those fungi which grow as single cells or very short dispersed hyphae. *Ustilago violacea* is of the first type and turbidity is correlated with dry weight (22). Methods have been evolved primarily for studies on bacteria (305-306) although most workers use a colorimeter in principle the nephelometer is better adapted to turbidity measurements. A turbidimetric method should of course always be calibrated against dry weight or other standard measure. Figure 1 illustrates the correlation.

Determination of growth by measurement of the total cellular nitrogen has been used only rarely in work with fungi (222). Although the *chromium* nitrogen complicates interpretation this is the best method for determination of growth defined as synthesis of protoplasm and deserves wider use.

Linear growth on agar is the least laborious method of estimating growth and has been correspondingly popular. The radial spread of a colony in a petri dish or linear advance in a growth rate tube (93-253) are most frequently determined. Growth is commonly expressed as the constant rate established after the initial period of slow growth.

Naturally the impurities present in agar rule out this method for certain problems. There are however more fundamental objections. Chaudhuri (56) found no response of colony diameter to even a four fold dilution of the nutrient medium. Nor does the rate of advance on agar always respond to necessary vitamins (106-179) or essential ions (238). Strain differences in growth on agar are not always reflected in dry weight measurements (133).

Chaudhuri (56) found on the other hand excellent correlation of radial growth on agar with dry weight and germ tube growth in temperature studies on *Fusicillium albo-atrum*. More recently Domsch (77) has confirmed the correlation in temperature response. Wood destroying fungi usually respond to temperature in agar culture about as they do in wood blocks (317). One crucial detail is often overlooked the depth of the medium influences growth and must be standardized (28-36-56).

We may conclude that the agar method is adequate for studies of some environmental factors e.g. temperature and (probably) composition of the atmosphere although even in these experiments the results are not always correlated with dry weight (91 134 145 250). It is decidedly not adequate for nutritional studies. Whether it should be used for essentially chemical problems like the effect of pH or toxic agents is still uncertain. The burden of proof that spread on agar is a fair index of growth rests on the investigator who chooses to use the method. Its value cannot be assumed.

6 TEMPERATURE AND GROWTH

Temperature affects growth spore germination reproduction and indeed all activities of the organism. Typical curves of growth as a function of temperature are shown in Figure 5. There is characteristically a linear portion in which growth increases directly with temperature an optimum range which may be narrow or rather broad and a descending limb as the temperature becomes too high for growth. The curve as a whole is usually skewed to the right; an empirical method for converting this skewed curve to a straight line is described by Cohen and Yarwood (58).

The temperature growth curve tends to become more nearly symmetrical as the optimum becomes lower: fungi with optima of 22–24°C or less often have a much less skewed curve (53 98 143 171 311). Not all low temperature fungi however follow this pattern (113 164).

Fungi with high temperature optima generally have the typical skewed curve even more pronouncedly; in a few cases the high optimum is associated with a broader range of good growth but this is not the rule.

It should be noted that the concept of an optimum usually taken for granted is not in fact very clear. The response of *Phycomyces blakesleeanus* to temperature (247) illustrates the principal ambiguities in the concept of optimum temperature. First growth is most rapid at 20–25°C but the maximum total amount of growth at limiting levels of thiamine is greatest at 10°. Second the effect of temperature on final dry weight can be almost wiped out if the nitrogen supply is made limiting.

Another example of the dependence of temperature characteristics on other factors has been elucidated by L. Fries (103). *Coprinus pmetarius* grows poorly at 44°C because of the failure of methionine biosynthesis to keep pace with other processes; if exogenous methionine is supplied growth at the elevated temperature is normal. The ap




Figure 5A Temperature and growth (dry weight) of fungi. Curve 1 *Cephalosporium cosantinii* from data of Treschow (311) curve 2 *Pullularia pullulans* from data of Rennerfelt (743) curve 3 *Coprinus* sp. from data of Johnson and Jones (151)




Figure 5B Temperature and growth (colony diameter on agar) of fungi. Curve 1 *Verticillium tricorpus redra n* from Isaac (144) by permission of the Cambridge University Press curve 2 *Polyporus vaporarius* from data of Gäumann (117) curve 3 *Pythium debaryanum* from data of Roth (700)

parent optimum temperature for radial growth of *Sclerotinia fructicola* is affected by pH (301)

It is clear that temperature optima and ranges reported are valid only under specified conditions of time medium and method of measurement There is no single temperature optimum of growth Nevertheless we shall continue to employ the concept of an optimum temperature on the grounds that it is empirically useful and that gross differences in apparent optima provide a basis for ecological studies

A given metabolic process—respiration antibiotic production or vitamin synthesis—does not of course necessarily respond to temperature in the same way as does growth Studies on penicillin production in particular support this generalization (49 227) as do studies of bacterial growth and metabolism (78)

Over the range in which growth is proportional to temperature the relation may be expressed by the value of Q_{10} defined as the ratio of the growth rate at a given temperature to that at a temperature 10°C lower The values found by Fawcett (92) in studies on four fungi are comparable to those reported for bacterial growth (298)

The shape of the temperature growth curve is somewhat affected by the time of incubation Growth at suboptimal temperatures is often so slow that it is not apparent at all at a time when cultures at more favorable temperatures are growing vigorously Growth at supra optimal temperature may begin rapidly but slow down or cease entirely after a period of time (34 223 278 286) Both of these factors of course affect the curve so that use of only one incubation period may introduce error

Most fungi make at least some growth over a 25 or 30-degree range of temperatures but narrower ranges are known e.g. in *Merulius lacrymans* (53) and *Capnodium salicinum* (98) Narrow ranges are often associated with low optima but the reverse is not true—many fungi with low optima grow over a normally wide range of temperature

Temperature optima for plant pathogenic fungi have been collected by Togashi (307) and are depicted graphically in Figure 6 Not all of these reports can be accepted and details of the distribution are unduly affected by popular incubator temperatures but it can be seen that most members of this ecological group have optima in the region $20-30^{\circ}\text{C}$ and about half have their optima between 26 and 30°C

The relation between temperature and habitat is not yet entirely clear Many fungi are probably world wide in distribution i.e. their growth is limited by factors other than temperature Species of *Allomyces* on the other hand are clearly confined naturally to the tropical




Figure 6 The optimum temperatures for growth of plant pathogenic fungi. Compiled by Dr C E Yarwood from Togashi (30).

and warmer temperate regions of the earth (84). Subtropical strains of a species may have a higher temperature optimum than strains from temperate regions (38). The distribution of *Phymatotrichum omnivorum* is determined by winter soil temperature (90). The few marine fungi studied so far have temperature optima for growth that are somewhat higher than the temperature of the sea (8). The occurrence and dominance of fungi in decomposing plant materials are largely determined by temperature (150-223).

Dermatophytic fungi grow best in culture at 25–35°C i.e. below the host body temperature (284). Fungi causing systemic mycoses grow well at 37° but optima have not been reported. It is significant that *Aspergillus fumigatus*, a thermotolerant species, can cause disease in warm blooded animals.

Some examples of high and low temperature optima are collected in

Table 2 Low and High Temperature Optima in the Fungi

Fungus	Criterion of Growth	Optimum C
<i>Botrytis cinerea</i> (115)	Dry weight	15-18
<i>Gnomonia</i> spp (135)	Linear extension	17-19
<i>Hendersoniella</i> sp (98)	Linear extension	18
<i>Herpotrichia nigra</i> (113)	Linear extension	15
<i>Mastigosporium album</i> (23)	Linear extension	15
<i>Phacidium infestans</i> (232)	Dry weight	15
<i>Phoma apicola</i> (17)	Linear extension	16-20
<i>Phycopsis</i> sp (98)	Linear extension	15
<i>Phytophthora primulae</i> (309)	Linear extension	15-20
<i>Actinomyces (Streptomyces)</i> sp (123)	Linear extension	35-37
<i>Aspergillus fumigatus</i> (243)	Dry weight	37
<i>Aspergillus nidulans</i> (223)	Linear extension	40
<i>Coprinus finetarius</i> (104)	Dry weight	35-40
<i>Lenites trabea</i> (53)	Linear extension	35
<i>Mucor pusillus</i> (218)	Linear extension	40-46
<i>Rhizopus chinensis</i> (330)	Linear extension	37.5-40.5
<i>Sclerotium (Rhizoctonia) bataticola</i> (224-310)	Linear extension	31-35
<i>Sepedonium</i> sp (223)	Linear extension	45

Table 2 It should be mentioned that not only may species within a genus differ widely (300-330) but strains or geographical isolates of the same species may respond differently (83a-143)

Surprisingly many fungi are able to grow although usually only slowly at 0 or slightly less. Plants under snow may be infected by the parasitic snow molds —*Typhula* spp. *Fusarium nivale* and an unidentified basidiomycete (32-65-242). *Phacidium infestans* cause of a disease of pines is able to grow in culture at -3° although its optimum temperature is about 15° (232).

A second ecological group of fungi tolerant of low temperature is commonly associated with the spoilage of refrigerated foods. Fungi of many different genera fall into this group and some especially strains of *Cladosporium* and *Sporotrichum* grow at temperatures well below zero e.g. -5 to -8 . The literature on these forms is reviewed by Berry and Magoon (19) and more briefly by Pehrson (232). Again although these fungi tolerate extremely low temperatures their optima are higher (122).

Survival of spores and mycelium at temperatures too low for growth is discussed briefly in Chapter 13.

Most fungi are unable to grow at $35-40^{\circ}\text{C}$. There are however two general types of exceptions to this rule. The first group is a




Figure 6 The optimum temperatures for growth of plant pathogenic fungi. Compiled by Dr C E Yarwood from Togashi (307)

and warmer temperate regions of the earth (84). Subtropical strains of a species may have a higher temperature optimum than strains from temperate regions (38). The distribution of *Phymatotrichum omnivorum* is determined by winter soil temperature (90). The few marine fungi studied so far have temperature optima for growth that are somewhat higher than the temperature of the sea (8). The occurrence and dominance of fungi in decomposing plant materials are largely determined by temperature (150-223).

Dermatophytic fungi grow best in culture at 25-35°C i.e. below the host body temperature (284). Fungi causing systemic mycoses grow well at 37° but optima have not been reported. It is significant that *Aspergillus fumigatus*, a thermotolerant species, can cause disease in warm blooded animals.

Some examples of high and low temperature optima are collected in

pathogen (155 156 302 322) However it is not certain that the relation is as simple as it appears and in some wilt diseases temperature affects the host (100 171) and probably other soil organisms which influence the disease (264)

For a few plant diseases it has been definitely shown that differences between temperature optima for disease and for growth of the pathogen occur and reflect the effect of temperature on the host This has been established particularly for the seedling blight of wheat and corn caused by *Gibberella saubinetii* (72 73) and a root rot of sweet clover caused by *Plenodomus meliloti* (258) and has been suggested for other diseases (114)

Finally there are some plant diseases in which the observed action of temperature appears to be exerted on the pathogen but on spore germination or sporulation rather than on mycelial growth (95 155 156) It is conceivable of course that pathogenicity is distinct from growth is affected by temperature but so far this has not been demonstrated

7 pH AND GROWTH

Under given conditions a fungus will grow maximally over a certain range of initial pH values of the medium and will fail to grow at high and low extremes However it must be borne in mind that pH is not a unitary factor the mechanism of its action differs at different concentrations of hydrogen ion For example one part of the pH-growth curve may reflect the effect of a low pH on enzyme systems another the effect of a high pH on metal solubilities Acidity may also affect the entry of essential vitamins (341) surface metabolic reactions entry into the cell of organic acids or the uptake of minerals (226)

It is not surprising therefore to find that almost any factor in the environment may change the shape of the pH growth curve Such factors include temperature (232 327) time of harvest (175 210) gross changes in the medium (187 340) growth factor supply (178) calcium and magnesium levels (314) and nitrogen source (140)

Furthermore pH is affected during growth by metabolic activities—raised by absorption of anions or production of ammonia from nitrogenous compounds lowered by formation of organic acids or absorption of cations These effects of growth on pH complicate results particularly in the poorly buffered media commonly employed It seems probable that many of the broad optima recorded

miscellaneous one examples of which appear in Table 2 it includes saprophytes—especially coprophilic forms—and several wood destroying fungi. The second group the true thermophiles is defined by the ability to grow at 50°C or higher and the inability to grow at temperatures below about 30°. Relatively few such fungi are known examples include *Thermomyces lanuginosus* (*Monotospora lanuginosa*) (135a 313) *Thermoascus aurantiacus* (201) *Penicillium duponti* (240) and *Chaetorium thermophile* (169). No study of the fundamental basis of this behavior in fungi has appeared.

Truly thermophilic forms also are found among the actinomycetes reviewed by Bernstein and Morton (18). The obligately thermophilic *Micromonospora vulgaris* is distinguished by the high resistance of its conidia to heat damage (87). Both fungi and actinomycetes are active in decomposition of compost at 50°C (320).

Killing of fungi by high temperature is considered elsewhere (Chapter 13) spores are the usual experimental material. It need only be noted here that mycelium is easily killed by elevated temperatures and that many fungi die slowly when held in culture at a temperature just above the maximum for growth.

Fluctuations in temperature induce two types of zonation the more common of which results from the presence of zones of reproductive structures (Chapter 11). In a second type of temperature induced zonation the zones are entirely mycelial this phenomenon occurs in *Cercospora beticola* (60) and in other fungi (20 124).

The data compiled by Togashi (307) indicate that the optimum temperature for spore germination is generally very close to that for growth in culture. There are instances however in which spore germination is less inhibited than growth by supraoptimal temperatures (185 330). Spore germination at temperatures too low for growth has been reported in *Rhizopus* spp (330) and is presumably common.

The relation between a fungus incited plant disease and temperature is rather more complex inasmuch as at least two organisms are necessarily involved. In general storage and transit rot diseases are most severe at the temperature which favors mycelial growth of the pathogen examples include the decay of sweet potato by *Rhizopus* spp (172 330) and storage rots caused by *Rhizoctonia solani* (170 332). Typically however the pathogen will grow in culture at temperatures below the minimum for infection (33).

The vascular wilts of plants caused by *Fusarium* spp often have the same pattern i.e. the disease optimum is close to that of the

in Table 3 reflect the ability of the fungus to raise or lower the pH of an initially unfavorable medium data on *Colletotrichum lindemuthianum* (173) illustrate this phenomenon

The values of Table 3 should therefore be viewed with some caution at best they offer a rough guide to choice of media and a possible indication of ecological preferences They are not to be regarded as absolute figures in any sense

A double pH optimum is frequently reported a minimum appearing between two maxima in the curve This type of response has been exhaustively analyzed in studies on *Coprinus* spp (102-104) It appears that the minimum merely reflects pH-dependent unavailability of one or more inorganic elements provision of iron zinc, and calcium in available forms eliminates the double optimum and replaces it with a broad single optimum zone

An organism may also be able to grow even quite well at a pH which does not permit the initiation of growth (189) The metabolic basis of this phenomenon is not clear again the artificiality of the simple pH growth curve is emphasized

In contrast to the bacteria and the actinomycetes fungi are relatively more able to invade acid environments although we should avoid the common assumption that they are restricted to such environments In culture the larger basidiomycetes are often unable to grow at an initial pH above 7.0 examples include most species of *Marasmius* (182) and *Tricholoma* (222) Correspondingly limited ecological studies of these fungi indicate that they are in general restricted to acid environments (333) However *Marasmius rotula* grows in culture and is found in nature on substrates of widely different pH (182-338) and several other fleshy basidiomycetes notably species of *Coprinus* actually require alkaline conditions for best growth (104-120-312)

Biological types of fungi may only roughly be characterized as to pH preference Most plant pathogens—exceptions are evident in Table 3—grow best in media with an initial pH of 5.0 to 6.5 Species of *Streptomyces* are marked by a relative intolerance of acidity very few can grow at a pH lower than 5.0 The same intolerance of acidity appears also in other groups e.g. the lower phycomycetes (61-85) The dermatophytic fungi perhaps because they are usually grown in complex media seem rather indifferent to pH over the range 4.0 to 10.0 (281-285)

Other biological activities of the fungi respond to pH but not necessarily in the same way as growth As a rule the pH range for sporulation is narrower than that for vegetative growth (Chapter

Table 3 pH Optima of Some Fungi and Actinomycetes*

Organism	Criterion of Growth	Optimum†
<i>Aphanomyces euteiches</i> (274)	Linear extension	4.5-6.5
<i>Aspergillus flavus</i> (28)	Linear extension	5.5-8.4
<i>A. niger</i> (28)	Linear extension	4.4-7.5
<i>Boletus variegatus</i> (198)	Dry weight	5.0
<i>Botryosporium</i> (?) sp. (149)	Dry weight	6.6-7.4
<i>Cercospora kikuchii</i> (70)	Dry weight	4.1
<i>Chalara quercina</i> (14)	Dry weight	4.5
<i>Chytridium</i> sp. (61)	Dry weight	5.2-7.5
<i>Colletotrichum hibisci</i> (314)	Linear extension	3.5-8.0‡
<i>Coprinus</i> sp. (154)	Dry weight	4.8-6.9
<i>Fomes annosus</i> (89)	Linear extension	4.6-4.9
<i>G. fraxineus</i> (203)	Linear extension	6.0-7.0
<i>Fusarium aqueductum</i> (229)	Dry weight	4.0-9.0
<i>F. saraitarum</i> (260)	Linear extension	6.3-7.0
<i>Geotrichum</i> sp. (229)	Dry weight	3.0
<i>Marasmius graminum</i> (182)	Dry weight	5.7-6.4
<i>Merulius confusus</i> (219)	Dry weight	4.0
<i>Mycogone perniciosus</i> (311)	Dry weight	6.7
<i>Neurospora crassa</i> (255)	Linear extension	4.3-6.5
<i>Ophiobolus graminis</i> (327)	Dry weight	4.9-7.4
<i>Penicillium expansum</i> (28)	Linear extension	4.4-7.5
<i>P. glaucum</i> (2)	Dry weight	5.0-6.5
<i>Phlyctothiza variabilis</i> (251)	Dry weight	7.2-7.6
<i>Phycomyces blakesleeana</i> (47)	Dry weight	3.5-4.6
<i>Pythium</i> sp. (52)	Dry weight	6.5
<i>Sepedonium</i> sp. (230)	Dry weight	7.0-8.5
<i>Septoria pepi</i> (266)	Linear extension	4.8-6.8
<i>Streptomyces coelicolor</i> (57)	Dry weight	6.9-7.7
<i>S. scabies</i> (318)	Dry weight	6.9
<i>Tricholoma nudum</i> (222)	Dry weight	5.0-6.0
<i>Trichophyton persicolor</i> (285)	Dry weight	6.5-7.0
<i>Trichosporon cutaneum</i> (229)	Dry weight	4.0-7.0
<i>Verticillium dahliae</i> (311)	Dry weight	5.3
<i>V. dactyloides</i> (311)	Dry weight	6.7-7.0

* For reasons discussed in the text these figures must be considered as approximations and as limited to the conditions of the particular experiment

† Defined for this table as the pH range tested in which the growth was within 10 per cent of the maximum growth

‡ With added calcium and magnesium in the medium

in Table 3 reflect the ability of the fungus to raise or lower the pH of an initially unfavorable medium. data on *Colletotrichum lindemuthianum* (173) illustrate this phenomenon.

The values of Table 3 should therefore be viewed with some caution at best they offer a rough guide to choice of media and a possible indication of ecological preferences. They are not to be regarded as absolute figures in any sense.

A double pH optimum is frequently reported a minimum appearing between two maxima in the curve. This type of response has been exhaustively analyzed in studies on *Coprinus* spp. (102-104). It appears that the minimum merely reflects pH-dependent unavailability of one or more inorganic elements. provision of iron, zinc, and calcium in available forms eliminates the double optimum and replaces it with a broad single optimum zone.

An organism may also be able to grow even quite well at a pH which does not permit the initiation of growth (189). The metabolic basis of this phenomenon is not clear again the artificiality of the simple pH growth curve is emphasized.

In contrast to the bacteria and the actinomycetes fungi are relatively more able to invade acid environments although we should avoid the common assumption that they are restricted to such environments. In culture the larger basidiomycetes are often unable to grow at an initial pH above 7.0. examples include most species of *Marasmius* (182) and *Tricholoma* (222). Correspondingly limited ecological studies of these fungi indicate that they are in general restricted to acid environments (338). However *Marasmius rotula* grows in culture and is found in nature on substrates of widely different pH (182-338) and several other fleshy basidiomycetes notably species of *Coprinus* actually require alkaline conditions for best growth (104-120-312).

Biological types of fungi may only roughly be characterized as to pH preference. Most plant pathogens—exceptions are evident in Table 3—grow best in media with an initial pH of 5.0 to 6.5. Species of *Streptomyces* are marked by a relative intolerance of acidity very few can grow at a pH lower than 5.0. The same intolerance of acidity appears also in other groups e.g. the lower phycmycetes (61-85). The dermatophytic fungi perhaps because they are usually grown in complex media seem rather indifferent to pH over the range 4.0 to 10.0 (284-285).

Other biological activities of the fungi respond to pH but not necessarily in the same way as growth. As a rule the pH range for sporulation is narrower than that for vegetative growth (Chapter

Table 3 pH Optima of Some Fungi and Actinomycetes*

Organism	Criterion of Growth	Optimum†
<i>Aphanomyces euteiches</i> (274)	Linear extension	4.5-6.5
<i>Aspergillus flavus</i> (28)	Linear extension	5.5-8.4
<i>A. niger</i> (28)	Linear extension	4.4-7.5
<i>Boletus variegatus</i> (198)	Dry weight	5.0
<i>Botryosporium</i> (?) sp (149)	Dry weight	6.6-7.4
<i>Cercospora kikuchii</i> (70)	Dry weight	4.1
<i>Chalara quercina</i> (14)	Dry weight	4.5
<i>Cladidium</i> sp (61)	Dry weight	5.2-7.5
<i>Colletotrichum hibisci</i> (314)	Linear extension	3.5-8.0‡
<i>Coprinus</i> sp (154)	Dry weight	4.8-6.9
<i>Fomes annosus</i> (89)	Linear extension	4.6-4.9
<i>F. fraxireus</i> (203)	Linear extension	6.0-7.0
<i>Fusarium aqueductum</i> (229)	Dry weight	4.0-9.0
<i>F. aurantiacum</i> (260)	Linear extension	6.3-7.0
<i>Geotrichum</i> sp (229)	Dry weight	3.0
<i>Marasmius graminum</i> (182)	Dry weight	5.7-6.4
<i>Merulius confluens</i> (219)	Dry weight	4.0
<i>Mycogone perniciosa</i> (311)	Dry weight	6.7
<i>Neurospora crassa</i> (253)	Linear extension	4.3-6.5
<i>Ophiobolus graminis</i> (327)	Dry weight	4.9-7.4
<i>Penicillium expansum</i> (28)	Linear extension	4.4-7.5
<i>P. glaucum</i> (2)	Dry weight	5.0-6.5
<i>Phlyctothiza variabilis</i> (251)	Dry weight	7.2-7.6
<i>Phyomyces diakesiceanus</i> (47)	Dry weight	3.5-4.6
<i>Pyrenopeziza</i> sp (52)	Dry weight	6.5
<i>Sepedonium</i> sp (230)	Dry weight	7.0-8.5
<i>Septoria pepis</i> (266)	Linear extension	4.8-6.8
<i>Streptomyces coelicolor</i> (57)	Dry weight	6.9-7.7
<i>S. s. abies</i> (318)	Dry weight	6.9
<i>Tricholoma nudum</i> (222)	Dry weight	5.0-6.0
<i>Trichophyton persicolor</i> (285)	Dry weight	6.5-7.0
<i>Trichosporon cutaneum</i> (229)	Dry weight	4.0-9.0
<i>Verticillium malthosum</i> (311)	Dry weight	5.3
<i>V. psalliotae</i> (311)	Dry weight	6.7-7.0

* For reasons discussed in the text these figures must be considered as approximations and as limited to the conditions of the particular experiment

† Defined for this table as the pH range tested in which the growth was within 10 per cent of the maximum growth

‡ With added calcium and magnesium in the medium

in Table 3 reflect the ability of the fungus to raise or lower the pH of an initially unfavorable medium data on *Colletotrichum lindemuthianum* (173) illustrate this phenomenon

The values of Table 3 should therefore be viewed with some caution at best they offer a rough guide to choice of media and a possible indication of ecological preferences They are not to be regarded as absolute figures in any sense

A double pH optimum is frequently reported a minimum appearing between two maxima in the curve This type of response has been exhaustively analyzed in studies on *Coprinus* spp (102-104) It appears that the minimum merely reflects pH-dependent unavailability of one or more inorganic elements provision of iron zinc, and calcium in available forms eliminates the double optimum and replaces it with a broad single optimum zone

An organism may also be able to grow even quite well at a pH which does not permit the initiation of growth (189) The metabolic basis of this phenomenon is not clear again the artificiality of the simple pH growth curve is emphasized

In contrast to the bacteria and the actinomycetes fungi are relatively more able to invade acid environments although we should avoid the common assumption that they are restricted to such environments In culture the larger basidiomycetes are often unable to grow at an initial pH above 7.0 examples include most species of *Marasmius* (182) and *Tricholoma* (222) Correspondingly limited ecological studies of these fungi indicate that they are in general restricted to acid environments (338) However *Marasmius rotula* grows in culture and is found in nature on substrates of widely different pH (182-338) and several other fleshy basidiomycetes notably species of *Coprinus* actually require alkaline conditions for best growth (104-120-312)

Biological types of fungi may only roughly be characterized as to pH preference Most plant pathogens—exceptions are evident in Table 3—grow best in media with an initial pH of 5.0 to 6.5 Species of *Streptomyces* are marked by a relative intolerance of acidity very few can grow at a pH lower than 5.0 The same intolerance of acidity appears also in other groups e.g. the lower phycomycetes (61-85) The dermatophytic fungi perhaps because they are usually grown in complex media seem rather indifferent to pH over the range 4.0 to 10.0 (284-285)

Other biological activities of the fungi respond to pH but not necessarily in the same way as growth As a rule the pH range for sporulation is narrower than that for vegetative growth (Chapter

Table 3 pH Optima of Some Fungi and Actinomycetes*

Organism	Criterion of Growth	Optimum†
<i>Aphanomyces euteiches</i> (274)	Linear extension	4.5-6.5
<i>Aspergillus flavus</i> (28)	Linear extension	5.5-8.4
<i>A. niger</i> (28)	Linear extension	4.4-7.5
<i>Boletus variegatus</i> (198)	Dry weight	5.0
<i>Botryosporium</i> (?) sp. (149)	Dry weight	6.6-7.4
<i>Cercospora kikuchii</i> (70)	Dry weight	4.1
<i>Chalara quercina</i> (14)	Dry weight	4.5
<i>Chytridium</i> sp. (61)	Dry weight	5.2-7.5
<i>Colletotrichum hibisci</i> (314)	Linear extension	3.5-8.0‡
<i>Coprinus</i> sp. (154)	Dry weight	4.8-6.9
<i>Fomes annosus</i> (89)	Linear extension	4.6-4.9
<i>F. fraxineus</i> (203)	Linear extension	6.0-7.0
<i>Fusarium aquaeductum</i> (229)	Dry weight	4.0-9.0
<i>F. aurantiacum</i> (260)	Linear extension	6.3-7.0
<i>Geotrichum</i> p. (229)	Dry weight	3.0
<i>Marasmius graminum</i> (182)	Dry weight	5.7-6.4
<i>Merulius confluens</i> (219)	Dry weight	4.0
<i>Mycogone perniciosa</i> (311)	Dry weight	6.7
<i>Neurospora crassa</i> (253)	Linear extension	4.3-6.5
<i>Ophiobolus graminis</i> (327)	Dry weight	4.0-7.4
<i>Penicillium expansum</i> (28)	Linear extension	4.4-7.5
<i>P. glaucum</i> (2)	Dry weight	5.0-6.5
<i>Polytorhiza variabilis</i> (251)	Dry weight	7.2-7.6
<i>Phycomyces blakesleeana</i> (47)	Dry weight	3.5-4.6
<i>Pythiopsis</i> sp. (52)	Dry weight	6.5
<i>Sepedonium</i> sp. (230)	Dry weight	7.0-8.5
<i>Septoria pepis</i> (266)	Linear extension	4.8-6.8
<i>Streptomyces coelicolor</i> (57)	Dry weight	6.9-7.7
<i>S. scabies</i> (318)	Dry weight	6.9
<i>Tricholoma nudum</i> (222)	Dry weight	5.0-6.0
<i>Trichophyton persicolor</i> (285)	Dry weight	6.5-7.0
<i>T. lasporum</i> cuartum (229)	Dry weight	4.0-9.0
<i>Verticillium nialhensis</i> (311)	Dry weight	5.3
<i>V. psalliotae</i> (311)	Dry weight	6.7-7.0

* For reasons discussed in the text these figures must be considered as approximations and as limited to the conditions of the particular experiment

† Defined for this table as the pH range tested in which the growth was within 10 per cent of the maximum growth

‡ With added calcium and magnesium in the medium

in Table 3 reflect the ability of the fungus to raise or lower the pH of an initially unfavorable medium data on *Colletotrichum lindemuthianum* (173) illustrate this phenomenon

The values of Table 3 should therefore be viewed with some caution at best they offer a rough guide to choice of media and a possible indication of ecological preferences They are not to be regarded as absolute figures in any sense

A double pH optimum is frequently reported a minimum appearing between two maxima in the curve This type of response has been exhaustively analyzed in studies on *Coprinus* spp (102-104) It appears that the minimum merely reflects pH-dependent unavailability of one or more inorganic elements provision of iron zinc and calcium in available forms eliminates the double optimum and replaces it with a broad single optimum zone

An organism may also be able to grow even quite well at a pH which does not permit the initiation of growth (189) The metabolic basis of this phenomenon is not clear again the artificiality of the simple pH-growth curve is emphasized

In contrast to the bacteria and the actinomycetes fungi are relatively more able to invade acid environments although we should avoid the common assumption that they are restricted to such environments In culture the larger basidiomycetes are often unable to grow at an initial pH above 7.0 examples include most species of *Marasmius* (182) and *Tricholoma* (222) Correspondingly limited ecological studies of these fungi indicate that they are in general restricted to acid environments (338) However *Marasmius rotula* grows in culture and is found in nature on substrates of widely different pH (182-338) and several other fleshy basidiomycetes notably species of *Coprinus* actually require alkaline conditions for best growth (104-120-312)

Biological types of fungi may only roughly be characterized as to pH preference Most plant pathogens—exceptions are evident in Table 3—grow best in media with an initial pH of 5.0 to 6.5 Species of *Streptomyces* are marked by a relative intolerance of acidity very few can grow at a pH lower than 5.0 The same intolerance of acidity appears also in other groups e.g. the lower phycomycetes (61-85) The dermatophytic fungi perhaps because they are usually grown in complex media seem rather indifferent to pH over the range 4.0 to 10.0 (284-285)

Other biological activities of the fungi respond to pH but not necessarily in the same way as growth As a rule the pH range for sporulation is narrower than that for vegetative growth (Chapter

11) The range for spore germination is similarly more narrow (185-291) this is to be expected inasmuch as the design of the spore germination experiment does not permit sufficient time for metabolic activities of the organism to modify the pH of the medium. Soil borne plant diseases may or may not show the same pH curve as do the fungi which cause them (96-260) here of course the effect of pH on the plant presumably exerts an important influence and the complex effect of pH on the soil itself may be even more decisive.

In most physiological studies it is necessary that the pH be controlled at least within limits. Periodic addition of alkali (51) and the use of solid calcium carbonate in the medium (99) are satisfactory for certain problems but in general it is necessary to employ a buffer system. Of these the most common and generally useful is the phosphate buffer. The usefulness of phosphate is limited however by the tolerance of the fungus to the ion this may be as low as 0.003 M (61) or as high as 0.20 M (66-151) and must be determined for each organism. Other buffer systems must be employed with caution acetate is often toxic (64-183-291) and citrate may be utilized by the fungus as a source of carbon with consequent displacement of the apparent pH optimum (64). Phthalate borate and citrate buffers are satisfactory for *Neurospora crassa* (273) α -hydroxybutyrate has been proposed as a non-utilizable buffer for studies on fungi (303). Equipment for automatic pH control has been described (50).

Physiological buffer systems can be devised for particular problems e.g. acetate as carbon source and ammonium ion as nitrogen source (188) or mixtures of ammonium and nitrate ions (141). In both these systems utilization of an anion is balanced by utilization of a cation and the pH is thereby held relatively constant.

In conclusion the pH growth curve has to say the least a limited usefulness. In designing experiments of this type the first question to ask is whether the information to be gained is worth the effort required. Once such an experiment is decided upon two rather elementary precautions should be observed. First the final pH of the medium after growth must be reported. Second the results will be more meaningful if buffer systems are used to control pH in the alkaline range addition of sodium hydroxide to control pH is followed by absorption of carbon dioxide from the air with a consequent false impression of the upper pH limit of growth.

The internal pH of the fungus cell—and the fungus cell is usually large enough so that the concept of pH may legitimately be applied—is not known with any precision. Indicator methods which have obvious uncertainties yield in most hands a value of about 6.0 (4).

193 219) close to the value for resting cells of yeast (59) Bunning (44) reports a lower internal pH below 5.0 for *Aspergillus niger*. Whether or not the internal pH responds to changes in the acidity of the external medium is uncertain there may be some effect (44 228) but it is to be presumed that any large change would result in injury and death. The principal effects of external pH are probably on permeability and other surface phenomena.

8 OXYGEN AND GROWTH

Fungi are commonly thought of as strictly aerobic, and this opinion is basically correct. However the quantitative relations of growth and oxygen supply vary considerably among different forms.

Visually observable growth of fungi occasionally occurs under nitrogen or in a liquid filled container (61 191 220). This limited growth probably depends on oxygen previously dissolved in the medium. *Blastocladia pringsheimii* grows well in tank nitrogen or carbon dioxide but fails to grow if oxygen is chemically absorbed by pyrogallol: the fungus has therefore a very low but real oxygen demand (85).

Achlya prolifera has been reported to grow in the presence of pyrogallol (134). However this report cannot be accepted without confirmation in view of the overwhelming evidence that other fungi and species of *Streptomyces* fail to grow at very low or zero oxygen pressures (69 94 119 127 166 192 195 204 234 237 257 292 342 343).

When growth on agar is the criterion it appears that most fungi grow as well at 20–40 mm oxygen pressure as at atmospheric (160 mm) (69 116 118 261 295). However the dry weight is usually affected at somewhat higher oxygen pressures: growth of *Ophiobolus graminis* is reduced at 105 mm oxygen (94). *Aspergillus oryzae* grows best at oxygen pressures higher than atmospheric (292). Contradictory data may be cited (12 166) but it seems that mass increase is more sensitive to oxygen deficit than is the more diffuse and thinner growth of a colony on agar.

In keeping with the relatively low oxygen requirement for growth fungi successfully colonize environments in which oxygen is limited e.g. relatively stagnant aquatic milieu. It is doubtful whether invasion of woody tissues is ever prevented by lack of oxygen (295) and the same is probably true of invasion by vascular pathogens and decay fungi. The number of fungi usually falls off at lower soil depths (31 45 147) but this may be caused more by high carbon dioxide

than by low oxygen (46) The adverse effect of flooding the soil on the growth and survival of fungi is however believed to reflect a deficit in oxygen (287 288 289)

Very high oxygen pressure may somewhat reduce growth or change the mycelial habit but does not usually have a lethal effect (159 328)

Many discussions of aerobiosis in the fungi have confused growth and fermentation Numerous fungi ferment sugars anaerobically (Chapter 7) this capacity has no essential bearing on growth and these forms should not be referred to as anaerobes The term an aerobe in fact cannot be applied to any fungus all require at least some oxygen and none has as yet been reported to grow better at low than at atmospheric oxygen pressure although reducing substances favor growth of *Armillaria mellea* (148) The vigorous anaerobic fermentation of some aquatic phycomycetes (Chapter 7) is coupled with a low oxygen demand and this relation is presumably a causal one

9 CARBON DIOXIDE AND GROWTH

Growth stimulation by carbon dioxide is considered later with carbon dioxide fixation (Chapter 7) High carbon dioxide pressures generally inhibit the growth of fungi but the level at which inhibition appears is quite variable Thus *Alternaria solani* is markedly inhibited by 38 mm of carbon dioxide (159) and *Coniophora cerebella* by 23 mm (345) but many other fungi are only slightly retarded in growth by carbon dioxide pressures of 150 mm or more (12 69 117 295) An isolate of *Zygorhynchus vulleminii* found only below the soil surface is retarded about 35 per cent by 152 mm of carbon dioxide while the surface soil inhabitant *Penicillium nigricans* is completely inhibited under the same conditions (16) Finally we may mention *Blastocladia pringsheimii* which grows well in an atmosphere of tank carbon dioxide and which requires carbon dioxide for the formation of resistant sporangia (85)

10 WATER AND GROWTH

The association of moisture and fungal growth is well known in relation to the deterioration of natural and manufactured products Fungi require relatively high moisture levels but most of the higher fungi can grow in the absence of liquid water

Two related types of study concern us here the effect of relative humidity on growth and the effect of osmotic pressure on growth. Both presumably measure the ability of the organism to draw water from its environment. Spore germination per se is considered elsewhere (Chapter 12) but it should be noted that experimental designs used in growth studies do not fully discriminate between spore germination and mycelial development.

Relative humidity effects must of course be determined by growing the organism on a substrate which is in equilibrium with the atmosphere. Agar and other solid substrates—bran, wood blocks, fabrics, etc.—have been used. Unfortunately quantitative determination of growth on such materials is difficult and the data are rather imprecise. This imprecision makes it hard to decide whether relative humidity itself or the moisture content of the equilibrated substrate is the more important factor in fungus growth. Data can be cited for each view (21, 277). Probably the amount of water in the substrate is the more important; thus if cotton and wool are incubated at 92 per cent relative humidity only the wool, which absorbs more water at this humidity than does cotton, supports growth of fungi (21).

The influence of moisture on fungal deterioration of economically important materials has been studied intensively and reviewed often (13, 184, 271, 277). Fungi tolerant of low humidity (85 to 90 per cent) include *Stereum frustulosum* (13), *Schizophyllum commune* (131) and *Aspergillus* spp. (163, 276). A few fungi attack substrates at as low as 65 per cent relative humidity (276) but most are limited to much higher humidities, 95 per cent or more.

Tolerance to low relative humidity appears as expected to be associated with tolerance to high osmotic pressure (131, 157); this relation is especially clear in spore germination experiments (Chapter 12). Species in which spore germination occurs at low humidity also tolerate a low humidity during growth (321, 323) and it is reasonable to infer that spore germination is more affected by marginal water supply than is mycelial growth (329).

Studies on the role of osmotic pressure are always complicated by possible non osmotic effects of the solute. Sodium chloride is distinctly inhibitory to some fungi (207); other complications are discussed by Brancato and Golding (29) and Stuart (290). Tolerance of high osmotic pressure is specific. Thus a mutant of *Neurospora crassa* is inhibited at 0.17 M glucose (139) while the other extreme species of *Aspergillus* can grow in media with an osmotic pressure of as high

as 100 atmospheres (28 282) Most fungi cease growth or are markedly inhibited when the concentration of soluble sugar exceeds about 2.0 M (109 129 159)

Obligate halophily—an absolute salt requirement—has been reported for several fungi (194 225 315 316) although at least some marine fungi do not require salt (8) Further quantitative studies on halophily in marine fungi would be of great interest

Protoplasm is presumably always more concentrated than the external solution and limited data support this assumption for the fungi (35) The indications are that the internal concentration as measured by freezing point depression or by plasmolysis methods increases as the external concentration is raised (195 225 231) In *Aspergillus niger* the relative amount of bound water is greater when the organism is grown in concentrated glucose (304) Ohtsuki (225) relying on the cryoscopic method reports an osmotic concentration in *Aspergillus glaucus* var *tonophilus* equivalent to approximately 250 atmospheres i.e. the freezing point of the expressed cell sap was -21°C


Thatcher (296 297) using plasmolytic methods found that the osmotic pressure of plant pathogenic fungi exceeds that of their hosts and may be a factor in parasitism Osmotic pressure of hyphae was found in this study to be 15.5–41.3 atmospheres that of haustoria of rust fungi to be 18.6–21.9 atmospheres

Fungi are more active at moderate than at high soil moisture (287) this is presumably to be ascribed to the effect of excessive soil water on aeration

11 BIOLUMINESCENCE

The phenomenon of bioluminescence in organisms generally is reviewed by Harvey (125) Wassink (324) critically reviews its occurrence in fungi and the biochemistry of the process is summarized by Johnson et al. (153) Bioluminescence occurs in at least 17 species of the basidiomycetes possibly in one ascomycete and has not been reported from phycomycetes (125 324) Within a species both luminescent and non luminescent strains occur in nature (21 42) in *Panus stipticus* luminescence is governed by a single gene (190) Luminescence occurs in sporophores rhizomorphs sclerotia and vegetative hyphae (42 125 241) As Robert Boyle discovered almost three centuries ago oxygen is necessary modern work shows that the oxygen requirement for fungal luminescence is much higher than that for bacterial (128) The reaction is temperature sensitive and is reversibly

Figure 7 Emission spectra of bio luminescence in *Photobacterium fischeri* (left) and *Armillaria mellea* (right) Redrawn from Spruit van der Burg (279) by permission of Elsevier Publishing Company Amsterdam



inhibited by narcotics (12) Active extracts have not as yet been prepared from fungi by analogy with other luminescent organisms we may safely assume that the process is enzymatic

Figure 7 compares the emission spectrum of the light produced by *Armillaria mellea* with that of *Photobacterium fischeri* Other fungi which have been examined—*Agaricus melleus* *Mycena* spp *Omphalia flava* *Pleurotus lunatellus* and *Polyporus hanelai*—have approximately the same maximum as *A. mellea* 520 to 530 mμ (125)

BIBLIOGRAPHY

- 1 Abe M T Yamano Y Kozu and M Kusumoto 1951 *J Agr Chem Soc Japan* 24 416-492
- 2 Acklin O and W Schneider 1978 *Biochem Z* 202 246-283
- 3 Andrus C F 1941 *Phytopathology* 31 566-67
- 4 Armstrong J I 1979 *Protoplasma* 8 229-260
- 5 Atkinson R C 1954 *Can J Botany* 39 673-678
- 6 Badcock E C 1949 *Trans Brit Mycol Soc* 23 200-203
- 7 Fallentine R and D G Stephens 1951 *J Cellular Comp Physiol* 37 369-387
- 8 Barghoorn E S and D H Linder 1944 *Ferrousia* 1 393-467
- 9 Bartholomew W H et al 1950a *Ind Eng Chem* 42 1801-1809
- 10 Bartholomew W H et al 1950b *Ind Eng Chem* 49 1827-1830
- 11 Bauer R 1938 *Z ntr Bakteriologie Parasitenk Abt II* 98 153-167
- 12 Bavendamm W 1928 *Centr Bakteriologie Parasitenk Abt II* 75 476-482 503-533
- 13 Bavendamm W and H Reichelt 1938 *Arch Mikrobiol* 9 485-544
- 14 Beckman C H J E Kuntz and A J Riker 1953 *Phytopathology* 43 441-447
- 15 Behr G 1930 *Arch Mikrobiol* 1 418-444
- 16 Benham R 1945 *Proc Soc Exp Biol Med* 58 199-201
- 17 Bennett C W 1971 *Mich State Univ Ag Exp Sta Tech Bull* 53 1-40
- 18 Bernstein A and H F Morton 1934 *J Bacteriol* 27 65-69
- 19 Berry J A and C A Magoon 1934 *Phytopathology* 24 780-796

- 20 Bisby G R 1925 *Mycologia* 17 89-97
- 21 Block S S 1953 *Appl Microbiol* 1 287-293
- 22 Blumer S 1937 *Arch Mikrobiol* 8 458-478
- 23 Bollard F G 1950 *Trans Brit Mycol Soc* 33 250-264
- 24 Bothe F 1935 *Arch Protistenk* 83 369-383
- 25 Bottcher E J and H J Conn 1949 *J Bacteriol* 41 137
- 26 Boudru M 1933 *Bull inst agron et ses recherches Gembloux* 2 310-346
(*Rev Appl Mycol* 13 335 1934)
- 27 Boyle C 1924 *Ann Botany* 98 113-133
- 28 Brancato F P and N S Golding 1953 *Mycologia* 45 848-864
- 29 Brancato F P and N S Golding 1954 *Mycologia* 46 442-456
- 30 Bretzlöff C W Jr 1954 *Am J Botany* 41 58-67
- 31 Briereley W B S T Jewson and M Brierley 1928 *Proc Intern Congr Soil Sci* 3 48-71
- 32 Broadfoot W C and M W Cormack 1941 *Phytopathology* 31 1058-1059
- 33 Brooks C and J S Cooley 1917 *J Agr Research* 8 159 163
- 34 Brooks C and J S Cooley 1921 *J Agr Research* 22 431-461
- 35 Brooks S C and M M Brooks 1941 *The Permeability of Living Cells* Berlin Zehlendorf Gebruder Borntraeger pp 39
- 36 Brown W 1923 *Ann Botany* 37 105-129
- 37 Brown W 1925 *Ann Botany* 39 373-408
- 38 Brown W and R K S Wood 1953 *Symposium Soc Gen Microbiol* 3 26-339
- 39 Buell C B and W H Weston 1947 *Am J Botany* 34 551-561
- 40 Bullen J J 1949 *J Pathol Bacteriol* 61 117-120
- 41 Buller A H R 1912 *Trans Brit Mycol Soc* 4 106-112
- 42 Buller A H R 1954 *Researches on Fungi* Vol 4 London Longmans Green and Co pp 611
- 43 Buller A H R and A T Cameron 1919 *Trans Roy Soc Can* 6 IV 73 18
- 44 Bunning L 1936 *Flora* 131 81-110
- 45 Burges A 1939 *Botanica Ser trimestral* 8 61-81
- 46 Burges A and E Fenton 1953 *Trans Brit Mycol Soc* 26 101-108
- 47 Burkholder P R and I McVeigh 1940 *Am J Botany* 27 634-640
- 48 Burkholder P R and E W Sinnott 1941 *Am J Botany* 32 424-431
- 49 Calam C I N Driver and R H Boyers 1951 *J Appl Chem (London)* 1 209-216
- 50 Callow D S and S J Lirt 1950 *J Gen Microbiol* 11 661-671
- 51 Cantino E C 1919a *Am J Botany* 36 91-112
- 52 Cantino E C 1949b *Am J Botany* 36 747-756
- 53 Cartwright A St G and W P K Findley 1934 *Ann Botany* 48 481-495
- 54 Chain E B and G Gualandri 1954 *Rend ist super sanità* (English ed) 17 5-60
- 55 Chain E B et al 1951 *Rend ist super sanità* (English ed) 17 61 80
- 56 Chaudhuri H 1923 *Ann Botany* 37 519-539
- 57 Cochrane V W and J F Conn 1947 *J Bacteriol* 51 213-218
- 58 Cohen M and C E Yarwood 1959 *Plant Physiol* 27 634-6 8
- 59 Conway E J and M Downey 1950 *Biochem J (London)* 47 355-360
- 60 Coons G H and F G Larmer 1930 *Papers Mich Acad Sci* 11 75-101
- 61 Crasemann J M 1954 *Am J Botany* 41 302-310
- 62 Crewther W G and F C Lennox 1955 *Australian J Biol Sci* 6 410-421
- 63 Cunningham K G and G G Freeman 1953 *Biochem J (London)* 53 318-332
- 64 Dags J and O Karkyte 1943 *Protoplasma* 39 127-151
- 65 Dahl A S 1934 *Phytopathology* 24 197-214

- 66 Darby R T and G R Mandels 1954 *Mycologia* 46 276-288
- 67 Day D and A Hervey 1916 *Plant Physiol* 21 233-236
- 68 Deindoeffer F and E L Gaden Jr 1955 *Appl Microbiol* 3 253-257
- 69 Denny F E 1933 *Contribs Boyce Thompson Inst* 5 9-10^o
- 70 Deutschmann F 1933 *Phytopathol Z* 20 297-310
- 71 De Vay J E 1954 *Phytopathology* 44 583-587
- 72 Dickson J C 1973 *J Agr Research* 23 837-870
- 73 Dickson J C S H Eckerson and K P Link 1973 *Proc Nat Acad Sci U S A* 9 434-439
- 74 Dietzel E H Behrenbruch and M Eucken 1910 *Arch Mikrobiol* 15 179-184
- 75 Dimond A E G H Plumb E M Stoddard and J G Horsfall 1919 *Conn (New Haven) Agr Exp Sta Bull* 531 1-69
- 76 Dion M A Carilli M Sermoniti and E B Chain 1954 *Rend ist super santid* (English ed) 17 187-205
- 77 Domsch K H 1955 *Arch Mikrobiol* 23 79-87
- 78 Dorn F L and O Rahn 1939 *Arch Mikrobiol* 10 6-1^o
- 79 Dox A W 1913 *J Biol Chem* 16 479-481
- 80 Dox A W and L Maynard 1912 *J Biol Chem* 1^o 227-231
- 81 Drouhet E and F Mariat 1909 *Ann inst Pasteur* 83 506-514
- 82 Drouhet E and R C Zapater 1914 *Ann inst Pasteur* 87 396-40
- 83 Duckworth R B and G C M Harris 1919 *Trans Brit Mycol Soc* 3^o 224-235
- 83a Edgington L A and J C Walker 1957 *Phytopathology* 47 594-598
- 84 Emerson R 1941 *Lloydia* 4 77-144
- 85 Emerson R and E C Cantino 1948 *Am J Botany* 35 17-171
- 86 Emerson S 1950 *J Bacteriol* 60 291-293
- 87 Erikson D 1952 *J Gen Microbiol* 6 96-94
- 88 Erwin D C 1954 *Phytopathology* 44 137-144
- 89 Etheridge D E 1955 *Can J Botany* 33 416-428
- 90 Ezekiel W N 1945 *Phytopathology* 35 296-301
- 91 Farries E H M and A F Bell 1930 *Ann Botany* 44 43-455
- 92 Fawcett H S 1941 *Univ Calif Publ Agr Sci* 4 183-3
- 93 Fawcett H S 1953 *Ann Appl Biol* 12 191-198
- 94 Fellows H 1978 *J Agr Resea ch* 57 349-355
- 95 Felton M W and J C Walker 1946 *J Agr Res arch* 7 69-81
- 96 Fikry A 1932 *Ann Botany* 46 29-70
- 97 Finn R K 1954 *Bacteriol Revs* 18 254-274
- 98 Fisher E E 1939 *Ann Botany* 3 399-46
- 99 Foster J W and J B Davis 1918 *J Bacteriol* 56 329-338
- 100 Foster R E and J C Walker 1917 *J Agr Research* 74 165-186
- 101 Fred E B and S A Waksman 1978 *Laboratory Manual of General Microbiology* New York McGraw Hill Book Co pp 145
- 102 Fries L 1945 *Arkiv Botan* 32 (10) 1-8
- 103 Fries L 1953 *Physiol Plantarum* 6 531-563
- 104 Fries L 1956 *Svensk Botan Tidskr* 50 47-96
- 105 Fries N 1938 *Symbolae Botan Upsalienses* 3 (2) 1-188
- 106 Fries N 1943 *Symbolae Botan Upsalienses* 7 (2) 5-73
- 107 Fries N 1950 *Arkiv Botan Ser 2* 1 271-287
- 108 Fromageot C M Jutis and P Tessier 1949 *Bull soc chim biol* 31 689-695
- 109 Fukuda Y S Takami and Y Ikeshoji 1954 *Japan J Botany* 14 67-90
- 110 Garner H H et al. 1953 *Am J Botany* 40 89-96
- 111 Carrett S D 1953 *Ann Botany* 17 63-79
- 112 Czummann E 1959 *Angew Botan* 21 59-69

- 113 Gäumann E C Roth and J Anliker 1934 *Z Pflanzenkrankh u Pflanzenschutz* 44 97-116
- 114 Gäumann E and E Hafziger 1943 *Phytopathol Z* 16 85-105
- 115 Gaumann E and U Nef 1947 *Ber schweiz botan Ges* 51 258-261
- 116 Golding N S 1940a *J Dairy Sci* 23 879-889
- 117 Golding N S 1940b *J Dairy Sci* 23 891-898
- 118 Golding N S 1945 *J Dairy Sci* 28 757-760
- 119 Gottheb D and H W Anderson 1947 *Bull Torrey Botan Club* 74 293-307
- 120 Grainger J 1946 *Trans Brit Mycol Soc* 23 52-64
- 121 Greene H C and E B Fred 1931 *Ind Eng Chem* 24 1201-1209
- 122 Haines R B 1930 *J Exp Biol* 8 319-389
- 123 Haines R B 1931 *J Exp Biol* 9 41-60
- 124 Hall M P 1933 *Ann Botany* 47 543-578
- 125 Hartey E V 1952 *Biotinamines* enco New York Academic Press pp 649
- 126 Haskins R H and J Anastasiou 1953 *Mycologia* 45 573-582
- 127 Haskins R H and W H Weston Jr 1950 *Am J Botany* 37 759-760
- 128 Hastings J W 1952 *J Cellular Comp Physiol* 39 1-30
- 129 Hawkins L A 1916 *J Agr Research* 7 253-260
- 130 Hawn E J and T C Vanterpool 1953 *Can J Botany* 31 699-710
- 131 Hayashi K 1944 *Japan J Botany* 14 91-98
- 132 Heim A H and H Lechevalier 1956 *Mycologia* 48 628-666
- 133 Held V M 1955 *Phytopathology* 45 39-42
- 134 Hemmi T and T Abe 1928 *Japan J Botany* 4 113-123
- 135 Henriksson L E and J F Morgan Jones 1951 *Svensk Botan Tidskr* 45 648-657
- 136a Hens en A 1957 *Arch Mikrobiol* 21 63-81
- 136 Hildebrand E M 1938 *Botan Rev* 4 627-664
- 137 Hockenfull D J D 1946 *Biochem J (London)* 40 337-343
- 138 Holsten B von A von Holsten and N Fries 1953 *Exp Cell Research* 5 550-555
- 139 Horowitz N H 1951 *Growth* 15 (suppl) 47 67
- 140 How J E 1910 *Ann Botany* 4 135-150
- 141 Hultin E and L Nordstrom 1949 *Acta Chem Scand* 3 1405-1417
- 142 Humfeld H 1947 *J Bacteriol* 54 689-696
- 143 Humphrey C J and P V Siggers 1933 *J Agr Research* 47 997-1008
- 144 Isaac I 1953 *Trans Brit Mycol Soc* 36 180-191
- 145 Jaarsveld A 1940 *De invloed van verschillende bodemschimmels op de virulentie van Rhizoctonia solani Kuhn Thesis Univ of Amsterdam 101 pp*
- 146 Jacquiot C 1951 *Rev mycol (Paris)* 16 21-29
- 147 Jeffreys E G F W Brian H G Hemming and D Lowe 1953 *J Gen Microbiol* 9 314-341
- 148 Jennison M W M D Newcomb and R Henderson 1955 *Mycologia* 47 275-304
- 149 Jensen H L 1930 *Trans Innec Soc N S Wales* 55 699-707
- 150 Jensen H L 1941 *Proc Innec Soc N S Wales* 66 216-226
- 151 Jermyn M A 1953 *Australian J Biol Sci* 6 48-69
- 152 Jillson O F and W J Nickerson 1948 *Mycologia* 40 369-385
- 153 Johnson F B H Eyring and M J Polissar 1954 *The Kinetic Basis of Molecular Biology* New York John Wiley and Sons pp 874
- 154 Johnson C T and A C Jones 1941 *Mycologia* 33 424-433
- 155 Jones L R 1954 *Am J Botany* 41 601-609
- 156 Jones L R J Johnson and J G Dickson 1926 *Wisconsin Univ Agr Exp Sta Research Bull* 71 1-144
- 157 Kaess G and W Schartz 1935 *Arch Mikrobiol* 6 708-714
- 158 Katznelson H 1940 *Soil Sci* 49 83-93

- 159 Klaus H 1941 *Phytopathol Z* 13 126-190
- 160 Klotz L J 1923 *Ann Missouri Botan Garden* 10 299-368
- 161 Kluyver A J and L H C Perquin 1933 *Biochem Z* 266 66-81
- 162 Koch O G and G A Dedie 1906 *Zentr Bakteriell Parasitenk Abt II* 109 17-21
- 163 Koehler B 1938 *J Agr Resea ch* 56 291-307
- 164 Koellreuter J 1900 *Phytopathol Z* 1 159-160
- 165 Koffler H R L Emerson D Perlman and R H Burris 1945 *J Bacteriol* 50 517-548
- 166 Kuhn F 1938 *Zentr Bakteriell Parasitenk Abt II* 98 430-444
- 167 Kusnetzow S J 1937 *Mikrobiologiya* 1 3-18 83
- 168 Lakata G D 1904 *Appl Microbiol* 2 2-4
- 169 La Touche C J 1900 *Trans Brit Mycol Soc* 33 94-104
- 170 Lauritzen J I 1929 *J Agr Research* 38 90-108
- 171 Lauritzen J I 1937 *J Agr Research* 44 861-917
- 172 Lauritzen J I and L L Harter 1925 *J Agr Research* 30 793-810
- 173 Leach J G 1923 *Minn Unn Agr Exp Sta Tech Bull* 14 1-41
- 174 Leach J G and T M Currence 1938 *Minn Unn Agr Exp Sta Tech Bull* 129 1-37
- 175 Lehman S G 1923 *Ann Missouri Botan Garden* 10 111-178
- 176 Levine M 1930 *A Compilation of Culture Media for the Cultivation of Microorganisms* Baltimore The Williams and Wilkins Co pp 969
- 177 Levine S and Z J Osdal 1946 *J Bacteriol* 52 686-694
- 178 Lilly V G and H L Barnett 1947a *Am J Botany* 34 131-138
- 179 Lilly V G and H L Barnett 1947b *Mycologia* 39 699-708
- 180 Lilly V G and H L Barnett 1951 *Physiology of the Fungi* New York McGraw Hill Book Co pp 464
- 181 Lilly V G and H L Barnett 1953 *West Va Univ Agr Exp Sta Bull* 367 1-8
- 182 Lindeberg G 1944 *Symbolae Botan Upsalienses* 8 (2) 1-183
- 183 Lindeberg G and M Korjus 1949 *Phycol Plantarum* 2 103-113
- 184 Lindgren R M 1947 *U S Dept Agr Tech Bull* 807 1-30
- 185 Ling L and J Y Yang 1944 *Ann Botany* 8 91-104
- 186 Luers H R Kühles and H Fink 1930 *Biochem Z* 217 53-78
- 187 Luz G 1934 *Phytopathol Z* 7 585-638
- 188 Machlis L 1953a *Am J Botany* 40 450-460
- 189 Machlis L 1953b *Am J Botany* 40 460-464
- 190 Macrae R 1947 *Can J Research C* 20 411-431
- 191 McVickar D L 1942 *Am J Botany* 29 372-380
- 192 Macy H 1909 *Minn Univ Agr Exp Sta Tech Bull* 64 1-86
- 193 Mahdihassan S 1939 *Biochem Z* 226 93-908
- 194 Malevich O A 1936 *Mikrobiologiya* 5 813-817
- 195 Matsumoto T 1921 *Ann Missouri Botan Garden* 8 1-62
- 196 Mayenburg O H von 1901 *Jahrb wiss Botan* 36 381-400
- 197 Mazur P and W H Weston 1906 *J Bacteriol* 71 207-66
- 198 Melin E 1905 *Untersuchungen über die Bedeutung der Baummykorrhiza eine ökologisch physiologische Studie* Jena Gustav Fischer pp 152
- 199 Meyer E 1905 *Mycologia* 47 664-667
- 200 Michel Durand E 1938 *Bull soc chim biol* 20 339-412
- 201 Miede H 1907 *Die Selbsthaltung des Heus* Jena Gustav Fischer pp 127
- 202 Miller J J 194 *Can J Resea ch C* 23 16-43
- 203 Montgomery H B S 1936 *Ann Appl Biol* 23 465-486
- 204 Moore E J 1937 *Phytopathology* 27 918-930
- 205 Morton A G and D Broadbent 1955 *J Gen Microbiol* 12 248-258
- 206 Moyer A J and R D Coghill 1945 *Arch Biochem* 7 161-183

- 207 Moyer A J and R D Coghill 1946 *J Bacteriol* 51 57-78
- 208 Moyer A J O E May and H T Herrick 1936 *Zentr Bakteriell Parasitenk Abt II* 9: 311-324
- 209 Muller F W 1911 *Ber schweiz botan Ges* 51 163-256
- 210 Neal D C 1927 *Ann Missouri Botanical Garden* 14 359-407
- 211 Nickerson W J 1951 *Trans N J Acad Sci II* 13 140-145
- 212 Nickerson W J and C W Chung 1954 *Am J Botany* 41 114-120
- 213 Nickerson W J and G A Edwards 1949 *J Gen Physiol* 33 41-55
- 214 Nickerson W J and Z Mankowski 1953a *Am J Botany* 40 584-591
- 215 Nickerson W J and Z Mankowski 1953b *J Infectious Diseases* 92 20-25
- 216 Nickerson W J and F G Sherman 1952 *J Bacteriol* 64 667-678
- 217 Nickerson W J and N J W Van Raj 1949 *Biochim et Biophys Acta* 3 461-475
- 218 Noack K 1912 *Jahrb wiss Botan* 51 593-616
- 219 Nord F F 1932 *Ergeb Enzymforsch* 1 77-112
- 220 Nord F F and L J Sciarini 1946 *Arch Biochem* 9 419-437
- 221 Nordbring Hertz B 1955 *Physiol Plantarum* 8 691-717
- 222 Norkrans B 1950 *Symbolae Botan Upsalienses* 11 (1) 5-126
- 223 Norman A G 1930 *Ann Appl Biol* 17 513-613
- 224 Norton D C 1953 *Phytopathology* 43 633-636
- 225 Ohtsuka T 1953 *Japan J Botany* 14 147-160
- 226 Overstreet R and L Jacobson 1952 *Ann Rev Plant Physiol* 3 189-206
- 227 Owen S P and M J Johnson 1955 *Appl Microbiol* 3 374-379
- 228 Owens R G 1955 *Contribs Boyce Thompson Inst* 18 175-144
- 229 Painter H A 1954 *J Gen Microbiol* 10 177-190
- 230 Paladino S 1954 *Rend ist super sanit* (English ed) 17 145-148
- 231 Pantanelli E 1901 *Jahrb wiss Botan* 40 303-367
- 232 Lehrson S O 1918 *Physiol Plantarum* 1 58-66
- 233 Perlman D 1948 *Am J Botany* 35 360-363
- 234 Pine L 1941 *J Bacteriol* 68 671-679
- 235 Pine L 1955 *J Bacteriol* 70 375-381
- 236 Pratt C A 1954 *Ann Botany* 38 563-595 599-615
- 237 Preston A and E I McLennan 1948 *Ann Botany* 12 53-64
- 238 Purdy L H Jr and R G Grogan 1954 *Phytopathology* 44 36-39
- 239 Raper K B and D F Alexander 1945 *Mycologia* 37 499-525
- 240 Raper K B and C Thom 1949 *A Manual of the Penicillia* Baltimore The Williams and Wilkins Co pp 815
- 241 Reitsma J 1932 *Phytopathol Z* 4 461-525
- 242 Remsberg R E 1940 *Mycologia* 32 59-96
- 243 Rennerfeldt E 1941 *Arch Mikrobiol* 12 19-40
- 244 Rhodes M 1950 *Trans Brit Mycol Soc* 33 35-39
- 245 Rippel A and G Behr 1936 *Arch Mikrobiol* 7 534-589
- 246 Robbins W J 1947 *Ann N Y Acad Sci* 49 75-86
- 247 Robbins W J and F Kavanagh 1944 *Bull Torrey Botanical Club* 71 1-10
- 248 Robinson R 1956 *Trans Brit Mycol Soc* 10 307-311
- 249 Robinson G N 1957 *J Gen Microbiol* 6 336-345
- 250 Roth C 1955 *Phytopathol Z* 8 1-110
- 251 Rothwell F M 1956 *Am J Botany* 43 28-30
- 252 Roxburgh J M J F T Spencer and H R Sallans 1954 *J Agr Food Chem* 2 1191-1121
- 253 Ryan F J C W Beadle and F L Tatum 1943 *Am J Botany* 30 781-799
- 254 Salvin S B 1947 *J Invest Dermatol* 9 315-320
- 255 Salvin S B 1949a *J Infectious Diseases* 81 275-283
- 256 Salvin S B 1949b *Mycologia* 41 311-319
- 257 Sanford G B 1956 *Phytopathology* 16 575-547

- 58 Sanford G B 1933 *Can J Research C S* 337-348
- 59 Savage G M and M J Vander Brook 1946 *J Bacteriol* 52 383-391
- 60 Schaffner E and K Meyer Hermann 1930 *Phytopathol Z* 9 99-166
- 61 Schaffer T G and B E Livingston 1937 *Am J Botany* 24 109-119
- 62 Scherr G H and R H Weaver 1953 *Bacteriol Revs* 17 51-92
- 63 Schostakowitsch W 1893 *Flora* 81 367-393
- 64 Schroeder W T and J C Walker 1942 *J Agr Research* 65 291-248
- 65 Sharp E L and F G Smith 1952 *Phytopathology* 42 263-264
- 66a Sharp E L and F G Smith 1957 *Phytopathology* 47 473-479
- 66b Shaw D E 1951 *Proc Linn Soc N S Wales* 76 7-23
- 67 Sherf A F 1943 *Phytopathology* 33 330-332
- 68 Shirakawa H S 1955 *Am J Botany* 42 379-384
- 69 Shu P 1953 *J Agr Food Chem* 1 1119-1123
- 70 Shu P and M J Johnson 1948 *Ind Eng Chem* 40 1202-1203
- 71 Siu R G H 1951 *Microbiological Decomposition of Cellulose* New York Reinhold Publ Corp pp 531
- 72 Smith J H 1923 *Ann Botany* 37 341-343
- 73 Smith J H 1924 *New Phytologist* 23 63-8
- 74 Smith P G and J C Walker 1941 *J Agr Research* 63 1-0
- 75 Smithies W R 1953 *Biochem J (London)* 5 346-3 0
- 76 Snow D 1945 *Ann Appl Biol* 32 40-44
- 77 Snow D M H G Crochson and N C Wright 1944 *Ann Appl Biol* 31 10 110
- 78 Sost H 1953 *Arch Protistenk* 100 341-361
- 79 Spruit van der Burg A 1950 *Biochim et Biophys Acta* 5 113 118
- 80 Stadler D R 1952 *J Cellular Comp Physiol* 39 449-454
- 81 Staffeldt F E and E L Sharp 1954 *Phytopathol* 44 213-21
- 82 Stare A 1942 *Arch Mikrobiol* 13 14-92
- 83 Starks O B and H Koffler 1949 *Science* 109 493-494
- 84 Stockdale P M 1932 *B of the Cambodge Phil Soc* 28 84-101
- 85 Stockdale P M 1933b *J Gen Microbiol* 8 434-441
- 86 Stoll C 1954 *Phytopathol Z* 22 235-274
- 87 Stover R H 1953 *Can J Botany* 31 603-637
- 88 Stover R H 1955 *Soil Sci* 80 391-411
- 89 Stover R H N C Thornton and V C Dunlap 1953 *Soil Sci Soc* 17 38
- 90 Stuart L S 1910 *J Agr Research* 1 29-36
- 91 Tamiya H 1957 *Acta Phytochim (Japan)* 3 51-173
- 92 Tamiya H 1959 *Acta Phytochim (Japan)* 4 297-29
- 93 Tatum E L R W Barratt and V M Cutler Jr 1949 *Science* 105 709-711
- 94 Taubenhaus J J and W N Ezekiel 1953 *Am J Botany* 23 10-12
- 95 Thacker D G and H M Good 1952 *Can J Botany* 30 413-48
- 96 Thatcher F S 1939 *Am J Botany* 26 449-4 8
- 97 Thatcher F S 1942 *Can J Research C* 20 313-311
- 98 Thumann K V 1955 *The Life of Bacteria* New York The Macmillan Co pp 775
- 99 Thirumalachar M J 1954 *Rend ist sup r sanita (English ed)* 17 206-212
- 100 Thom C and K B Raper 1941 *U S Dept Agr Misc Publ* 426 1-46
- 101 Tullford P E 1936 *Ohio Agr Exp Sta Bull* 567 1-1
- 102 Tisdale W B 1953 *J Agr Research* 71 55-86
- 103 Tobie W C and G B Ayres 1945 *J Bacteriol* 50 333-333
- 104 Todd G W and J Levitt 1951 *Plant Physiol* 26 331-336
- 105 Toennies G and D L Gallant 1948 *J Biol Chem* 174 451-463
- 106 Toennies G and D L Gallant 1949 *Cronth* 13 7-0
- 107 Togashi K 1949 *Biological Characters of Plant Pathogens Temperature Relations* Tokyo Meibundo pp 478

- 308 Tomkins R G 1932 *Trans Brit Mycol Soc* 17 150-153
- 309 Tomlinson J A 1959 *Trans Brit Mycol Soc* 33 221-233
- 310 Tompkins C M and M W Gardner 1933 *Hilgardia* 9 219-250
- 311 Treschow C 1941 *Dansk Botan Arkiv* 11 (1) 1-31
- 312 Treschow C 1944 *Dansk Botan Arkiv* 11 (6) 1-180
- 313 Tsilinsky P 1899 *Ann inst Pasteur* 13 500-505
- 314 Venning F D and B S Crandall 1931 *Phytopathology* 41 465-468
- 315 Vishniac H S 1955a *J Gen Microbiol* 12 453-463
- 316 Vishniac H S 1955b *Mycologia* 47 633-643
- 317 Wagener W W and R W Davidson 1954 *Botan Rev* 20 61-134
- 318 Waksman S A 1929 *Soil Sci* 14 61-79
- 319 Waksman S A A Schatz and H C Reilly 1946 *J Bacteriol* 51 753-759
- 320 Waksman S A W W Umbreit and T C Cordon 1939 *Soil Sci* 47 37-54
- 321 Walderdorff M G 1904 *Botan Arch* 6 81-110
- 322 Walker M N 1941 *Florida Univ Agr Exp Sta (Gainesville) Bull* 363 3-29
- 323 Walter H 1974 *Z Botan* 16 353-417
- 324 Wassink E C 1948 *Rec trav botan Néerl* 41 150-211
- 325 Wean R E and J E Young 1939 *Phytopathology* 29 893-898
- 326 Webb M 1918 *J Gen Microbiol* 2 275-287
- 327 Webb R W and H Fellows 1976 *J Agr Research* 33 813-872
- 328 Webley D M 1934 *J Gen Microbiol* 11 114-121
- 329 Wehmer C 1914 *Mycol Centr* 4 241-252 287-299
- 330 Weimer J L and L L Harter 1923 *J Agr Research* 21 1-40
- 331 Weiss F and B A Oteifa 1953 *Phytopathology* 43 407
- 332 Wellman F L 1939 *J Agr Research* 43 461-469
- 333 Wernham C C 1946 *Mycologia* 38 691-692
- 334 Wetter C 1934 *Arch Mikrobiol* 70 761-212
- 335 White I G 1935 *Am J Botany* 42 79-161
- 336 Wilhelm S 1935 *Phytopathology* 45 180-181
- 337 Wilkin G D and A Rhodes 1935 *J Gen Microbiol* 12 259-264
- 338 Wilkins W H J L Harley and G C Kent 1938 *Ann Appl Biol* 25 472-489
- 339 Wolf F T 1935 *Bull Torrey Botan Club* 82 343-354
- 340 Wolpert F S 1924 *Ann Missouri Botan Garden* 11 43-97
- 341 Wyss O V G Lilly and L H Leonian 1944 *Science* 99 18-19
- 342 Yoshii H 1933 *Bull Sci Fakultato Terkultura Kyusu Imp Univ (Japan)* 6 312-330
- 343 Yoshii H 1936 *Ann Phytopathol Soc Japan* 6 199-204
- 344 Zobl K H 1943 *Arch Mikrobiol* 13 191-206
- 345 Zycha H 1937 *Zentr Bakteriell Parasitenk Abt II* 97 202-244

2 The Composition of Fungus Cells

Under this head are considered those constituents which as a class if not individually are common to most or all fungi. Materials which are found only in some species are discussed if significant in the section on metabolism: thus sugar alcohols and carotenoids do not occur in the mycelium of all fungi and so are not included here but are considered in respectively Chapters 5 and 6. Known vitamins produced by fungi are reviewed in Chapter 10.

The theoretical significance of chemical composition is somewhat limited. It will be seen that within bounds set by the metabolic capacities of the species the composition varies widely with the environment. The quantities of fat, carbohydrate, ash, wall material and total nitrogen are all more or less responsive to the culture medium.

In composition the principal difference between fungi and bacteria is in the high chitin content of most fungi. Chitin is present in the bacteria (176); the green algae do not have chitin walls. Absence of chitin in the Oomycetes discussed below in turn suggests a polyphyletic origin of the true fungi. A second difference between fungi and bacteria is also quantitative rather than absolute: lipid accumulation is more common and attains higher levels in the fungi than in most bacteria. *Mycobacterium* and related forms however are exceptions to this rule and may have a rather high lipid content (130).

I THE MAJOR CONSTITUENTS OF THE CELL

Growing fungus cells are of course largely water early measurements indicate that 85-90 per cent of the fresh weight of mycelium and fleshy sporophores is water Presumably the more leathery sporophores and sclerotia have less water (25 197) The conidia of most fungi appear to have a relatively low water content e.g. 17.4 per cent in *Aspergillus oryzae* (171) 25 per cent in *Monilia fructicola* and 6 per cent in *Penicillium digitatum* (195) It is probable as suggested by Yarwood (195) that most of the water in these spores is hygroscopic Their water content is largely therefore a function of the environmental humidity this may explain some of the rather high values reported in the early literature

Species of *Erysiphe* are unusual in that the conidia have a high water content (195 196) With this is associated the ability to germinate at low relative humidity (Chapter 12)

The bound water of *Aspergillus niger* mycelium defined as water which is removed by heat but not by *in vacuo* drying over a desiccant at moderate temperatures amounts to 71-84 per cent of the dry weight when the fungus is cultivated on 5 per cent glucose at higher glucose concentrations the bound water rises to as much as 30 per cent of the dry matter (182)

The elementary composition of fungi has been studied chiefly with respect to carbon and nitrogen The carbon content of a number of fungi studied by Heck (78) was 40-44 per cent of the dry weight Lunck and Allison (128) report a somewhat wider range—43 to 53 per cent *Aspergillus oryzae* mycelium is about 49 per cent carbon (173) and *A. niger* mycelium about 46 per cent (129) but other species of *Aspergillus* may be as high as 55 per cent (194) Most fungi are 10-50 per cent carbon and the value is relatively little affected by cultural conditions (173 194) or age (126)

Bacteria and viruses in general have carbon contents of the same order as fungi (130) The mycelium of *Streptomyces griseus* has slightly less carbon—about 38 per cent of the dry weight (58)

In contrast to the relative uniformity of carbon content the total nitrogen of fungal mycelium is quite variable both the kind and the concentration of the nitrogen source influence the composition

The major constituents of the mycelium of several fungi are listed in Table 1 These data cannot be considered typical of all fungi since it is known that at least four of the fractions—carbohydrate protein lipid and ash—are strongly affected by cultural conditions

Table 1 Major Constituents of Cultivated Fungi and Actinomycetes*

Organism	Carbohydrate	Chitin	Protein	Lipid	Ash
<i>Aspergillus oryzae</i> (spores) (171)	9.00	—	22.75†	2.20	5.33
<i>Gliocladium roseum</i> (157)	34.4	8.4	21.2‡	22.4	—
<i>Rhizopus japonicus</i> (99)	—	—	38.8†	9.72	5.52
<i>Penicillium chrysogenum</i> (157)	60.0	10.6	16.4‡	1.4	—
<i>Penicillium griseofulvum</i> (165)	45.0	5.5	12.0§	34.5	—
<i>Phymatotrichum omnivorum</i> (47)	32.42	—	23.94†	4.49	6.24

* As per cent of dry weight

† Estimated from total nitrogen only

‡ Estimated from total nitrogen after subtraction of chitin nitrogen

§ Determined in a chitin free fraction

and differ in amount from one species to another. As discussed below those protein values of Table 1 which are based only on the total nitrogen are almost certainly high.

2. MINERAL CONSTITUENTS OF THE CFL

The ash content of fungi varies with the medium used and presumably with species. Some figures reported for total ash as per cent of dry weight are: *Rhizopus japonicus* 5.52 (99), *Aspergillus niger* 2.39 (129), *A. oryzae* 7.19 (78), *Coprinus radians* 4.97 (78), *Penicillium notatum* 28.3–34.0 (50). Fruiting bodies collected in the field may be as low as 1.08 or as high as 29.9 per cent ash; most have 5–10 per cent (197).

The composition of the ash of a few fungi is summarized in Table 2. Although no figures should be considered typical, it is clear that the most abundant minerals are phosphorus and potassium. Probably none of the analyses of Table 2 is complete, and inspection raises some obvious problems, e.g. the accumulation of silicon in *Rhizopus japonicus* and aluminum in *Penicillium notatum*.

The total ash content and the concentration of individual elements are influenced by the environment and by the stage of development. In *Aspergillus niger* the total ash of conidia is about three times that of mycelium; the increase is confined, however, to a few of the elements, particularly phosphorus, potassium and magnesium (5, 139). Autolysis is accompanied by loss of phosphorus (61) and by a decline in total ash (100). Among the environmental effects, the simplest is the increase in the internal concentration of an element in the mycelium as its external concentration is raised (62, 139, 144, 161). The total ash and the concentration of individual elements are also influ

Table 2 The Mineral Constituents of Fungi*

Mineral	<i>Rhizopus japonicus</i> (99)	<i>Penicillium notatum</i> (50)	<i>Trametes suataeolens</i> (198)	<i>Morchella esculenta</i> (94)
K	27.3	4.52	31.52	63.07
Na	12.9	7.89	0.46	0.38
Ca	—	26.72	22.57	1.75
Mg	1.42	8.56	6.07	1.77
P	28.7	25.01	4.64	26.13
S	0.28	2.17	19.06	1.78
Si	21.2	—	—	0.63
Cl	—	0.5	0.52	1.37
Al	—	20.62	0.26	—
Fe	1.84	4.52	—	2.00
Zn	0.18	—	—	—

* As per cent of total ash

enced probably indirectly through effects on growth or pH by the type of nitrogen source (61, 143, 194) and by the concentration of nitrogen (78). A low nitrogen supply may be expected to act directly on phosphorus content by restriction of the synthesis of nucleic acids (63).

It is assumed that only phosphorus, sulfur, and chlorine occur in fungi in organic combination; an attempt to isolate organic compounds of magnesium failed (143). Phosphorus is of course present in several types of organic molecules: sugar phosphates and adenosine phosphates (22, 92, 113), phosphoproteins (107), nucleic acids (63), and phospholipids (p. 47). None of these has been investigated thoroughly in the fungi.

The inorganic phosphate of fungal mycelium includes orthophosphate and inorganic condensed phosphates (151). Pyrophosphate occurs in *Aspergillus niger* (106) but most attention has been directed to the metaphosphates, polymers of the general formula $(\text{NaPO}_3)_n$. Volutin, a common visible intracellular inclusion, is probably metaphosphate (85). Metaphosphates have been found in cells of *Aspergillus niger* (96) and possibly *Neurospora crassa* (83). The metaphosphate of *A. niger* has a molecular weight of 6000–7000 (86). Fungal enzymes which hydrolyze these condensed inorganic phosphates are considered briefly in Chapter 9. Utilization of the bond energy of these compounds to drive synthetic reactions, although not impossible in principle, has not as yet been demonstrated.

3 THE CELL WALL

The most serious defect in our knowledge of cell wall composition in the filamentous fungi is the fact that no pure wall material is yet available. The major sources of error are the inclusion of extraneous substances in the preparation and the loss of wall material during purification. We do not have therefore for any fungus even a qualitative picture of the chemical components of the wall. This section consequently can only review work in which one or the other major constituent has been studied. When we speak of a chitin wall, we mean a wall with detectable chitin, not necessarily or even probably a wall which is exclusively chitin. Morphologically in fact the wall may be a multiple structure with layers of different composition (15-46).

In most fungi the basic material of the cell wall is chitin, a polymer of N-acetylglucosamine, identical chemically with the chitin of arthropods. The history of the study of chitin in fungi is reviewed by Schmidt (152) and by Frey (56). Much confusion exists however because of the use of methods which are not critical. The most acceptable method of qualitative identification of chitin is determination of the X-ray diffraction pattern (56, 66, 79, 87, 91). However if chitin is present only in small amounts the X-ray method may not reveal it. Other methods—microchemical tests, isolation of derivatives from a hydrolyzate and determination of the nitrogen in a purified preparation—are usually in agreement with X-ray results but are not sufficiently discriminating to stand by themselves. Microchemical methods for chitin and cellulose are particularly unsatisfactory (56).

The quantitative determination of chitin requires purification usually by the method of Scholl (153) and determination of the nitrogen of the product. In a survey of 25 fungi, chitin contents from 2.6 to 26.2 per cent of the dry weight have been reported (18); other quantitative data are found in Table 1 above and in the paper of Schmidt (152). The possibility that some protein nitrogen from resistant proteins is included in these analyses cannot be ignored.

On the basis of microchemical tests for chitin, von Wettstein (186) proposed that large taxonomic groups of the fungi differ in the constitution of the cell wall. Specifically he found chitin in the Zygomycetes and in the higher fungi—ascomycetes and basidiomycetes, representative Myxomycetes, Oomycetes and Monoblepharidales were found not to form chitin. This supports a polyphyletic origin of the phycomycetes.

Table 3 Cell Wall Materials of Fungi*

Organism	Classification	Chitin	Cellulose
<i>Pythium debaryanum</i>	Oomycetes	Absent	Present
<i>Phytophthora</i> spp	Oomycetes	Absent	Present
<i>Saprolegnia lathronica</i>	Oomycetes	Absent	Present
<i>Achlya caroliniana</i>	Oomycetes	Absent	Present
<i>Lagenidium giganteum</i>	Oomycetes	Absent	Present
<i>Allomyces arbuscula</i>	Blastocladales	Present	Absent
<i>Blastocladiella variabilis</i>	Blastocladales	Present	Absent
<i>Phycomyces blakesleeanae</i>	Zygomycetes	Present	Absent
<i>Mucor</i> spp	Zygomycetes	Present	Absent
<i>Rhizopus bovinus</i>	Zygomycetes	Present	Absent
<i>Absidia glauca</i>	Zygomycetes	Present	Absent
<i>Basidiobolus ranarum</i>	Zygomycetes	Present	Absent
<i>Eremascus fertilis</i>	Endomycetaceae†	Present	Absent
<i>Endomyces despiens</i>	Endomycetaceae†	Present	Absent
<i>Endomycopsis fibuliger</i>	Endomycetaceae†	Present	Absent
<i>Saccharomyces</i> spp	Saccharomycetaceae†	Absent	Absent
<i>Scheuosaccharomyces</i> spp	Saccharomycetaceae†	Absent	Absent
<i>Zygosaccharomyces vini</i>	Saccharomycetaceae†	Absent	Absent
<i>Pichia membranifaciens</i>	Saccharomycetaceae†	Absent	Absent
<i>Ashbya gossypii</i>	Spermophthoraceae†	Absent	Absent
<i>Eremothecium ashbyii</i>	Spermophthoraceae†	Absent	Absent
<i>Torula</i> spp	Pseudosaccharomycetaceae†	Absent	Absent
<i>Rhodotorula rubra</i>	Pseudosaccharomycetaceae†	Absent	Absent
<i>Mycoderma vini</i>	Pseudosaccharomycetaceae†	Absent	Absent
<i>Aspergillus oryzae</i>	Ascomycetes†	Present	Absent
<i>Nectria cinnabarina</i>	Ascomycetes	Present	Absent
<i>Sclerotinia fructicola</i>	Ascomycetes	Present	Absent
<i>Glenomyces farfugifolius</i>	Ascomycetes	Present	Absent
<i>Boletus edulis</i>	Basidiomycetes	Present	Absent
<i>Cantharellus cibarius</i>	Basidiomycetes	Present	Absent
<i>Coprinus atramentarius</i>	Basidiomycetes	Present	Absent
<i>Hypholoma fasciculare</i>	Basidiomycetes	Present	Absent
<i>Ustilago violacea</i> (budding sporidia)	Basidiomycetes	Absent	Absent
<i>Blastomyces dermatitidis</i>	Fungi Imperfecti	Present	Absent
<i>Epidermophyton floccosum</i>	Fungi Imperfecti	Present	Absent
<i>Histoplasma capsulatum</i>	Fungi Imperfecti	Present	Absent
<i>Paracoccidioides brasiliensis</i>	Fungi Imperfecti	Present	Absent
<i>Sabouraudia audouinii</i>	Fungi Imperfecti	Present	Absent
<i>Sporobolomyces roseus</i>	Fungi Imperfecti	Present	Absent
<i>Sporotrichum schenckii</i>	Fungi Imperfecti	Present	Absent
<i>Trichoderma viride</i>	Fungi Imperfecti	Present	Absent
<i>Trichophyton ferrugineum</i>	Fungi Imperfecti	Present	Absent

* Limited to data from X ray diffraction studies of Blank (16-17) Frey (56) and Gonell (66)

† Classification of Gaumann (60)

‡ Assuming a perfect stage in the ascomycetes

Although several reports since 1921 have challenged von Wettstein's generalizations, more recent work with the X-ray diffraction method has confirmed them. These results are summarized in Table 3. From this compilation it can be seen that chitin is found in most of the true fungi except for the Oomycetes which are characterized by cellulose. The Chytridiales have not been studied by modern methods; individual species have been claimed to form chitin (111-186).

Two points need to be made in connection with the yeasts. The X-ray diffraction data of Table 3 indicate that they divide on family lines only the mycelial yeasts having chitin. The principal cell wall material of the unicellular yeasts appears to be a polyglucose distinct from cellulose (8) with a second membrane containing a mannan-protein complex (45-119). However, microchemical tests suggest that there is a small amount of chitin in unicellular yeasts; glucosamine can be isolated (45-145). The amount present in these yeasts is probably so small as to escape detection by the X-ray diffraction method.

The amount of chitin produced is lower in a medium with nitrate than in one with ammonium ion as the nitrogen source (7-152); this may be a pH effect. Autolytic loss of chitin is said to occur in nitrate but not in ammonium media (7).

Although it is probable that other materials are more or less closely associated with chitin in the cell wall, unambiguous evidence of such association is lacking. Pectic materials, protein, lipids, cellulose, callose, and minerals have all been suggested (47-103, 104-178-181). The cellulose problem is considered by Castle (23) and by Farr (46). In *Herpomyces stylopygae*, parasitic on insects, histochemical studies indicate the presence in the basal shield of a chitin-protein layer and a process of sclerotization similar to that in the development of insect cuticle may take place (141).

Chitin appears not to occur in the actinomycetes (57a); staining reactions of *Streptomyces* spp. are the basis of a suggestion that a lipid layer is associated, perhaps loosely, with the spore wall (43).

The sheath surrounding the stalk of *Dictyostelium discoideum* is composed of cellulose laid down extracellularly (137).

Callose, described by Mangin (105) and by Thomas (179), appears to be an acidic polyglucose; its relation to the cell wall itself is uncertain and cannot at present be accepted.

A material identified as lignin occurs in relatively large amounts in several fungi (128-177). The identification rests on solubility characteristics alone, and it seems very doubtful that the substance is chemically the same as the lignin of higher plants (69-155-165).

1 CELLULAR CARBOHYDRATES

In this section are included polysaccharides and related compounds exclusive of chitin just discussed. Some of these may be found later to be associated with the cell wall and others appear in the medium as well as the mycelium. Characteristically cells grown on high-carbohydrate media are rich in polysaccharides; for example the mycelium of *Aspergillus niger* may be 25 per cent polysaccharide (148).

The first fungal polysaccharide to be studied was that of *Penicillium glaucum* spores named spore starch by Cramer (29). In the period from 1894 to about 1925 several polysaccharides were described but with few exceptions the criteria of identification were inadequate and the nature of the materials remains in doubt. These early studies have been reviewed by Dox and Neidig (36) and by Schmidt (149). It need only be pointed out here that none of the presumed identifications with the starch of higher plants is based on sufficient evidence.

Fungal polysaccharides of known or partially known constitution are listed in Table 4. Most are formed by species of *Penicillium* and *Aspergillus* but this may only mirror the more intensive study which has been devoted to these genera. The most common sugar component is glucose; the next most common galactose.

Galactocarolose is distinguished by two unusual features: galactose is in the furanose form and the galactose units are joined by a 1,5 glycosidic bond (76). It has been speculated that the galactofuranose structure may be metabolically related to the tetrionic acid derivatives also formed by *Penicillium charlesii* (Chapter 6).

The glycogen of fungi is chemically identical with or very similar to that of animals (10) and occurs also in yeast (118). The fraction appearing as glycogen in routine analyses usually comprises about 5 per cent of the mycelial or spore dry weight (140-171) but in the sclerotia of *Phymatotrichum omnivorum* the glycogen content is 36.7 per cent (40).

Polysaccharides of which the constitution is not fully known include especially the hemicelluloses defined by their solubility and ease of hydrolysis. The crude hemicellulose fraction of *Aspergillus niger* constitutes 12 to 31 per cent of the dry matter of the mycelium and is like several other constituents of the mycelium profoundly affected by the supply of metals in the medium (129-156). Lentosans have been isolated as about 1 per cent of the dry matter of fungi (35-150) but the possibility that these materials are in fact hemicelluloses can

Table 4 Polysaccharides of Fungi and Actinomycetes*

Polysaccharide	Hydrolysis Product(s)	Organism
Glycogen	Glucose	<i>Aspergillus niger</i> (171) <i>Phymatostichum omnivorum</i> (40 41 42)
Sclerotiose	Glucose	<i>Penicillium sclerotiorum</i> (2)
Compactose	Glucose	<i>Penicillium brevis-compactum</i> (65)
Nigeran (mycodextran)	Glucose	<i>Aspergillus niger</i> (6 34) <i>Penicillium</i> spp ‡ (36 185)
Polyglucose	Glucose	<i>Aspergillus fischeri</i> (116 117)
Polyglucose	Glucose	<i>Penicillium charlesii</i> (14)
Acidic polyglucose	Glucose	<i>Penicillium luteum</i> (55)
Capsular polyglucose	Glucose	<i>Torulopsis</i> spp (101 102)
Luteic acid	Beta-glucose malonic acid	<i>Penicillium luteum</i> (3)
Varianose	Glucose galactose†	<i>Penicillium varians</i> (75)
Capreolinose	Glucose mannose galactose malonic acid	<i>Penicillium capreolinum</i> (26)
Polyuronide (?)	Glucose galacturonic acid†	<i>Coecidioides immittis</i> (72)
Levan	Fructose	<i>Aspergillus sydowii</i> (95)
Polyfructose	Fructose	<i>Streptomyces</i> sp (183)
Polygalactose	Galactose	<i>Penicillium charlesii</i> (27)
Rugulose	Galactose	<i>Penicillium rugulosus</i> (76)
Mycogalactan	Galactose	<i>Aspergillus niger</i> (37)
Neutral polygalactose	Galactose	<i>Penicillium luteum</i> (55)
Galactocarolose	Galactose	<i>Penicillium charlesii</i> (76)
Mannocarolose	Mannose	<i>Penicillium charlesii</i> (73-75)
Capsular polysaccharide	Xylose mannose glucuronic acid	<i>Cryptococcus (Torulopsis) neoformans</i> (38 44)
Capsular pentosan	Xylose	<i>Torulopsis</i> spp (101 102)

* Including only those polysaccharides for which some information on structure is available

† All hydrolysis products not identified

‡ Identification uncertain

not be excluded Mannans are common in yeasts but not in fungi (59)

5 NITROGENOUS CONSTITUENTS

The total amount of nitrogen in mycelium is usually determined by the Kjeldahl method which is only approximately complete Heck

(78) found the total nitrogen of cultivated fungi to range from 2.27 per cent of the dry weight in *Coprinus radians* to 5.13 per cent in *Trichoderma lignorum*, sporophores from nature had as little as 1.56 per cent nitrogen. These data are fairly typical although fungi grown in culture may be as high as 7.6 per cent nitrogen (115-132). Different species of *Tricholoma* vary from 3.1 to 7.7 per cent nitrogen (114).

Measurements of total nitrogen are however not highly significant by reason of the susceptibility of the value to environmental influences and its variation with developmental status. Characteristically young mycelium is higher in nitrogen than old (7) during autolysis both proteins and other nitrogen compounds are liberated into the medium (Chapter I). In *Aspergillus niger* the onset of sporulation is coincident with a decline in mycelial protein (168).

The most important single influence on the total nitrogen content is the concentration of nitrogen in the medium: an increase in the amount of available nitrogen is accompanied by a significant—as much as threefold—increase in cell nitrogen (78, 81, 166, 173).

The nitrogen source used also exerts an influence on the total nitrogen of the mycelium: in general the nitrogen content is higher when ammonium salts are used than when nitrates provide the nitrogen (93, 170, 173, 191). The cause of this effect is uncertain: either the low pH incident upon utilization of ammonium ion reduces autolytic loss of nitrogen or the nitrate is simply less available to the cell for synthesis.

Fractionation of the total nitrogen of the cell reveals that only about 60–70 percent is in proteins (67, 165); still lower values have been reported (99). Determination of cellular protein by calculation from total nitrogen is therefore inadmissible in principle although most protein determinations have been made in this way.

A few specific proteins have been described (20, 29, 112, 125) but no general or systematic study of fungal proteins has been undertaken. Michel Durand (107) found indications of phosphoproteins in *Aspergillus niger*. Qualitatively the amino acids of fungal proteins are much the same as those of other proteins: surveys of the amino acids in hydrolyzates of *Aspergillus flavus* (127), *Penicillium chrysogenum* (135), *Ustilago zeae* (32) and *Venturia inaequalis* (124) have been made by chromatography of the acid hydrolysate and indicate a general similarity among these diverse fungi with minor qualitative differences.

Quantitative analyses of the amino acids in hydrolysates of cellular material of fungi and actinomycetes (168, 191) similarly do not reveal

any striking differences between these and higher organisms when all proteins are hydrolysed together

The biological utilization of fungal proteins has been studied in two contexts. Mycelium added to the soil is decomposed by the microflora about as rapidly as are other organic materials (78). Norman (115) could find no evidence of a decay resistant nitrogenous fraction in mycelium (89). In animal nutrition the utilizability of fungus proteins is generally low, i.e. they do not support growth of the rat (49-52, 162, 163, 189). The nutritional value of fungal proteins is reviewed by Thatcher (175); the mycelium of some fungi is toxic to laboratory animals, but the underlying mechanism—whether an amino acid deficiency or a toxic substance—is not known.

The non-protein nitrogen of the fungus cell consists of chitin, already discussed, nucleic acids, free amino acids and miscellaneous nitrogen compounds of low molecular weight. The nucleic acid of *Penicillium* spp. accounts for 4 per cent of the total nitrogen and about 1 per cent of the dry weight (1, 165). These findings cast some doubt on earlier claims reviewed by Akashi (1) of high nucleic acid concentrations in fungi. Spores of *Penicillium* spp. contain 158-592 μg ribonucleic acid per billion (33). The conidia of *Aspergillus nidulans* contain measurable deoxyribonucleic acid and the amount is as in other organisms, twice as high in diploid as in haploid cells (77).

Free amino acids occur in fungus mycelium (31, 68, 109, 120, 127, 158, 190) and are also found in the culture medium (32, 109, 135).

The other nitrogenous constituents known in fungi include many different and unrelated compounds, several of which are discussed in Chapter 8. Nitrogen bases and amines have been found (90, 171, 197) possibly arising by degradation of larger molecules during analysis. Cyclic choline sulfate has been isolated from fungi (31, 167, 192). Urea appears to be a frequent constituent of the higher basidiomycetes (88) and constitutes 3.9 per cent of the total nitrogen of *Aspergillus fischeri* mycelium (67). Allantoic acid has been reported in appreciable amounts in the sporophores of Agaricales (54) and may be one precursor of urea (Chapter 8).

6 CELL LIPIDS

The total lipid content of fungus mycelium, usually expressed as a percentage of the dry weight, is so strongly affected by cultural conditions and age that any set of figures is misleading. The effect of these conditions is brought out especially well in a survey by Walker

and co workers (110 160 188) some of the known environmental factors determining fat content are discussed in Chapter 6. High fat values—20 per cent or more of the dry weight—are found among species and strains of *Aspergillus* *Endomyces* *Fusarium* *Mucor* *Oospora* *Penicillium* and *Torula* (10 18 110 141 152 160 161 185 188) and in individual species of other genera (147 157 188). The highest fat contents reported are about 50 per cent of the dry weight (10 110). Consideration of common saprophytic fungi do not have unusually high fat contents but the sclerotia of *Claviceps purpurea* contain 43 per cent lipids (151) and uredospores of *Puccinia graminis tritici* are high in fat (Chapter 12).

Some variation in apparent lipid content arises from the use of different methods of extraction these are discussed with reference to the fungi by Peck (122) and by Bernhauer and others (11).

In Table 5 are collected a few partial analyses of fungal lipids. The

Table 5 Lipids of Fungi

Organism	Iodine Number	Acid Number	Phospho- tides*	Unsapon- ifiable Matter*	Sterols*
<i>Aspergillus niger</i> (15)	95†	71.2	—	12.0	14†
<i>Aspergillus syd. az.</i> (169)	114.4	43.4	Present	8.2	5.4
<i>Blastomyces dermatitidis</i> (123)	106†	45.3	24.3	8.0	—
<i>Fusarium graminearum</i> (108)	84.7	—	—	2.1	—
<i>Penicillium javanicum</i> (184)	84.0	10.6	—	2.0	—
<i>Phycomyces blakesleeanae</i> (9)	—	—	9.9	5.3	4.0†

* As per cent of total lipid

† Ergosterol

most striking feature is the usually very high proportion of free fatty acids up to 88 per cent of the readily extractible lipids may be free acids (158). However it should be noted that the content of free acid is markedly affected by cultural conditions. In *Penicillium javanicum* the acid number rises from 10.6 to 50.7 as the glucose level of the medium is increased from 20 to 40 per cent (185). In general any factor which increases fat formation also increases the relative amount of free fatty acids (138). It should also be noted that *Aspergillus nidulans* lipids are very low in free acid (161) as are sclerotia of *Claviceps purpurea* (197). It seems possible that lipase action during

preparation of the mycelium may contribute to the high apparent acid value of fungal lipids

The fatty acid composition of some fungal lipids is summarized in Table 6. These and other data (10, 53, 97) show that the 16 and 18 carbon acids are the most abundant and that with few exceptions the unsaturated acids—especially oleic and linoleic—predominate. The low Reichert Meissl number of fungal lipids reflects the relative absence of the lower fatty acids (10).

The antibiotic activity of a strain of *Penicillium crustosum* has been traced to the presence of linoleic acid in its mycelium (142).

A number of fatty acids which are not known to be constituents of fats have been isolated from fungi. Three of these—agaric, spicularic and mineoluteic acids—are long chain hydroxy tricarboxy acids similar to some of the lichen acids (136). *Actinomyces* (*Streptomyces*) *flavus* forms a wax, the dodecyl ester of 5 keto octadecanoic acid (82) and 6 keto octadecanoic acid has been isolated from sporophores of *Lactarius* spp. (21). The ustilic acids identified as di and trihydroxy hexadecanoic acids occur as constituents of the glucolipid ustilagic acid, a metabolic product of *Ustilago zeae* (98).

The phosphatides determined by their solubility are usually less than 10 per cent of the total lipid but in *Blastomyces dermatitidis* may amount to as much as one third of the lipid fraction (122). In *Aspergillus* spp. the phosphatides constitute at least 0.15 to 0.7 per cent of the dry weight (138, 193).

Excluding earlier studies reviewed by Czapek (30) which employed inadequate methods, well characterized phosphatides are a lecithin and a cephalin from *Aspergillus sydowii* (193) and a lecithin and a cephalin not identical with these from *Blastomyces dermatitidis* (122). These have been characterized by their physical properties and their hydrolysis products; structures have not been determined. In addition a cephalin fraction from *Neurospora crassa* has been described and evidently comprises more than one compound since both serine and ethanolamine are present (39). The phospholipid fraction of *Penicillium chrysogenum* yields inositol phosphate (81) and an inositol phospholipid occurs in *Neurospora crassa* (57). Similarly the positive Molisch reaction of the acetone insoluble lipids of several fungi indicates that a carbohydrate containing phospholipid may be present (122).

The mycelium of *Aspergillus sydowii* contains at least 0.1–0.4 per cent of a cerebrin (19) this partially characterized nitrogenous lipid is probably identical with that isolated earlier from yeast *Claviceps purpurea* and from the sporophores of several basidiomycetes. The

Table 6 The Fatty Acid Composition of Fungal Lipids*

Acid	<i>Aspergillus nidulans</i> (161)	<i>A. sydowii</i> (169)	<i>Penicillium japonicum</i> (184)	<i>Penicillium</i> sp. (80)	<i>Phycomyces blakesleeana</i> (9)	<i>Penicillium lilactinum</i> (159)
Myristic (dodecanoic)	0.7	—	—	3.5	—	0.1
Palmitic (tetradecanoic)	20.9	10.9	23.4	40.5	23.7	32.3
Stearic (octadecanoic)	15.9	13.6	9.4	7.9	4.7	9.4
Arachidic (eicosanoic)	{ 1.4 }	—	—	1.1	—	1.4
Behenic (docosanoic)		—	—	—	1.7	—
Lignoceric (tetracosanoic)		1.1	0.8	—	2.1	—
Hexacosanoic	—	—	—	—	1.3	—
Hexadecenoic	1.2	—	—	7.9 (?)	—	3.4
Oleic (octadecenoic)	40.3	36.7	34.6	19.0	29.6	38.6
Linoleic (octadecadienoic)	17.0	20.0	31.8	20.1	25.8	13.4
Linolenic (octadecatrenoic)	0.2	—	—	—	3.4	—
Tetracosenoic	—	—	—	—	2.9	—
Hexacosenoic†	—	—	—	—	4.9	—

* As per cent of the total fatty acids after saponification

† Undenitrified as to structure

occurrence and the chemistry of the cerebriins are reviewed by Celmer and Carter (24)

Virtually all fungi appear to form sterols. The amount in mycelium is usually about 1 per cent or less of the dry weight. Comprehensive surveys have been made by Pruess and others (132-134) the sterol content was found to be affected by the growth medium the highest level attained being 1.70 per cent of the dry weight in *Paeclomyces varioti*. Different isolates assigned to *Aspergillus niger* vary in sterol content from 0.23 to 1.16 per cent (12). A more recent survey (4) finds that the sterol content of fungi ranges from zero in *Colletotrichum graminicolum* to 2.2 per cent of the dry weight in *Penicillium westlingii*.

The principal sterol of fungi is ergosterol, a dihydro derivative fungisterol occurs in ergot (174). Ergosterol occurs as the palmitate in *Penicillium* spp. (121) and *Aspergillus fumigatus* (187). Other sterols have been only tentatively identified (70, 71, 146).

The metabolism of exogenous steroids and the composition of the carotenoid pigments are considered elsewhere (Chapter 6).

BIBLIOGRAPHY

- 1 Akai S. 1939. *J Biochem (Tokyo)* 9, 21-29.
- 2 Alberici V. J. T. P. Curtin and D. Reilly. 1913. *Biochem J (London)* 37, 243-246.
- 3 Anderson C. G. W. N. Haworth H. Raistrick and M. Stacey. 1939. *Biochem J (London)* 33, 272-279.
- 4 Appleton G. S. R. J. Kiebel and W. J. Payne. 19, *Appl Microbiol* 3, 49-51.
- 5 Bajak V. S. P. Damle and P. S. Krishnan. 1954. *Arch Biochem Biophys* 50, 451-460.
- 6 Barker S. A. E. J. Bourne and M. Stacey. 1953. *J Chem Soc (London)* 1953, 3084-3090.
- 7 Behr G. 1930. *Arch Mikrobiol* 1, 418-441.
- 8 Bell D. J. and D. H. Northcote. 1950. *J Chem Soc (London)* 1950, 1944-1947.
- 9 Bernhard K. and H. Albrecht. 1948. *Helv Chim Acta* 31, 977-988.
- 10 Bernlauer K. 1913. *Ergeb Enzymforsch* 9, 297-360.
- 11 Bernhauer K. I. Muller and E. Theile. 1950. *Biochem Z* 320, 94-298.
- 12 Bernhauer K. and G. Patz. 1953. *Biochem Z* 380, 388-393.
- 13 Bernhauer K. and G. Posselt. 1937. *Biochem Z* 291, 215-230.
- 14 B. Kinshaw J. H. J. H. V. Charles and H. Raistrick. 1931. *Phil Trans Roy Soc London Ser B* 220, 1-56.
- 15 Blackwell E. 1910. *Trans Brit Mycol Soc* 24, 68-86.
- 16 Bla. k. T. 1953. *Biochim et Biophys Acta* 10, 110-113.
- 17 Blank F. 1954. *Can J Microbiol* 1, 1-5.
- 18 Blumenthal H. J. and S. Roseman. 1957. *J Bacteriol* 64, 222-224.
- 19 Bohonos N. and W. H. Peterson. 1943. *J Biol Chem* 149, 293-300.

- 20 Bohonos N D W Woolley and W H Peterson 1942 *Arch Biochem* 1 319-324
- 21 Bougalt J and C Charoux 1911 *Compt rend (Paris)* 153 519-523
- 22 Buston H W and A H Khan 1956 *J Gen Microbiol* 14 625-629
- 23 Castle E S 1945 *Am J Botany* 32 148-151
- 24 Ceimer W D and H E Carter 1959 *Physiol Pers* 37 167-196
- 25 Chatfield C and G Adams 1940 *U S Dept Agr Circ* 549 1-91
- 26 Clutterbuck P W 1936 *J Soc Chem Ind (London)* 55 55T-61T
- 27 Clutterbuck P W W N Haworth H Raistrick G Smith and M Stacey 1934 *Biochem J (London)* 28 94-110
- 28 Clutterbuck P W R Lovell and H Raistrick 1932 *Biochem J (London)* 26 1907-1918
- 29 Cramer E 1891 *Arch Hyg* 20 191-210
- 29a Cummins C S and K Harris 1938 *J Gen Microbiol* 18 173-189
- 30 Czapek F 1922 *Prochemie der Pflanzen* 3d ed Vol 1 Jena Gustav Fischer pp 898
- 31 DeFlines J 1932 *J Am Chem Soc* 54 1676-1677
- 32 DeLay J E 1934 *Phytopathology* 24 583-587
- 33 Dirks J 1932 *Biochim et Biophys Acta* 8 191-201
- 34 Dox A W 1915 *J Biol Chem* 20 83-84
- 35 Dox A W and R E Neidig 1911 *J Biol Chem* 9 267-269
- 36 Dox A W and R F Neidig 1914a *J Biol Chem* 18 167-175
- 37 Dox A W and R E Neidig 1914b *J Biol Chem* 19 23-237
- 38 Drouhet E C Segretain and J P Aubert 1930 *Ann Inst Pasteur* 79 891-900
- 39 Ellman G I and H K Mitchell 1934 *J Am Chem Soc* 56 4028-4030
- 40 Ergle D R 1947 *J Am Chem Soc* 69 2061-2062
- 41 Ergle D R 1948 *Phytopathology* 28 147-151
- 42 Ergle D R and L M Blank 1947 *Phytopathology* 37 153-161
- 43 Erikson D 1947 *J Gen Microbiol* 1 39-44
- 44 Evans E E and R J Theriault 1953 *J Bacteriol* 65 571-577
- 45 Falcone G and W J Nickerson 1956 *Science* 124 272-273
- 46 Farr W K 1934 *Trans N Y Acad Sci II* 16 209-214
- 47 Farr W K and S H Eckerson 1934 *Contribs Boyce Thompson Inst* 6 189-203
- 48 Fink H H Hahn and W Hoerburger 1937 *Chemiker Ztg* 61 689 693 725-726 744-747
- 49 Fink H and A Hock 1941 *Z Naturforsch* 2(b) 187-203
- 50 Fink H J Schlie and U Ruge 1933a *Z physiol Chem Hoppe Seyler's* 207 231-235
- 51 Fink H J Schlie and U Ruge 1933b *Z physiol Chem Hoppe Seyler's* 203 264-267
- 52 Fink H J Schlie and U Ruge 1933c *Z physiol Chem Hoppe Seyler's* 205 173-178
- 53 Fosse J 1918 *Arch Biochem* 16 161-168
- 54 Fosse R and A Brunel 1933 *Compt rend (Paris)* 197 289-290
- 55 Freeman G G and C S Macpherson 1949 *Biochem J (London)* 43 149-185
- 56 Frey R 1930 *Ber Schweiz botan Ges* 60 199-230
- 57 Fuller R C and E L Tatum 1936 *Am J Botany* 43 361-363
- 58 Garner H H et al 1935 *Am J Botany* 40 289-296
- 59 Gartuly Janke R 1940 *Zentr Bakteriell Parasitenk Abt II* 107 361-363
- 60 Gaumann E 1949 *Die Pilze* Basel Verlag Birkhauser pp 58
- 61 Gayet J 1918a *Bull soc chim biol* 50 483-496
- 62 Gayet J 1918b *Bull soc chim biol* 50 549-547

- 63 Gayet J 1919a *Bull soc chim biol* 31 796-800
- 64 Gayet J 1919b *Bull soc chim biol* 31 1046-1051
- 65 Godin P 1953 *Biochim et Biophys Acta* 10 528-532
- 66 Gonell H W 1976 *Z physiol Chem Hoppe Seyler's* 152 18-30
- 67 Gorica H J W H Peterson and H Steenbock 1934 *Biochem J (London)* 28 504-511
- 68 Gordon M P Numerof and S C Pan 1951 *J Am Chem Soc* 73 4037-4038
- 69 Gottlieb S and M J Pelczar Jr 1951 *Bacteriol Revs* 15 55-77
- 70 Gruber W and C Broske 1951 *Monatsh Chem* 82 25-258
- 71 Guider J M T G Halsall and E R H Jones 1951 *J Chem Soc (London)* 1951 4471-4475
- 72 Hassid W A F E Baker and R M McCready 1953 *J Biol Chem* 149 303-311
- 73 Haworth W N E L Hurst F Isherwood and J K N Jones 1939 *J Chem Soc (London)* 1939 1818-1880
- 74 Haworth W N H Raistrick and M Stacey 1935a *Biochem J (London)* 29 612-621
- 75 Haworth W N H Raistrick and M Stacey 1935b *Biochem J (London)* 29 2668-678
- 76 Haworth W N H Raistrick and M Stacey 1937 *Biochem J (London)* 31 610-644
- 77 Heagy F C and J A Roper 1952 *Nature (London)* 170 713-714
- 78 Heck A F 1959 *Soil Sci* 27 1-46
- 79 Hevly A N J 1936 *Protoplasma* 23 377-396
- 80 Hilditch T P et al 1944 *J Soc Chem Ind (London)* 63 117-114
- 81 Hilpert R S G Friesen and W Rosske 1937 *Biochem Z* 283 193-197
- 82 Hirata Y and K Nakanishi 1949 *Bull Chem Soc Japan* 22 121-127
- 83 Houlahan M B and H K Mitchell 1948 *Arch Biochem* 19 25-261
- 84 Imai Y 1950 *J Japan Biochem Soc* 29 19-196 (*Chem Abstr* 45 9191 1951)
- 85 Ingelman B 1950 In J B Sumner and K Myrbeck (eds) *The Enzymes* Vol 1 (1) p 511-516 New York Academic Press
- 86 Ingelman B and H Malmgren 1950 *Acta Chem Scand* 4 48-486
- 87 Iterson G van Jr K H Meyer and W Lotmar 1956 *Proc trav chim* 5 61-63
- 88 Iwanoff N N 1957 *Z physiol Chem Hoppe Seyler's* 150 214-218
- 89 Jensen H L 1957 *J Agr Sci* 29 1-2
- 90 Keil W and H Bartmann 1953 *Biochem Z* 280 58-60
- 91 Khouvinc Y 1957 *Compt rend (Paris)* 19 396-397
- 92 Kita D A and W H Peterson 1953 *J Biol Chem* 203 861-868
- 93 Klotz L J 1953 *Ann Missouri Bot n Garden* 10 299-368
- 94 König J 1950 *Chemie der Nahrungs- und Genussmittel sowie der Gebrauchsgegenstände* Vol 2 Berlin Julius Springer pp 937
- 95 Kopeloff N I Kopeloff and C J Welcome 1950 *J Biol Chem* 43 171-187
- 96 Krishnan P S and V Bajaj 1953 *Arch Biochem Biophys* 47 39-53
- 97 Krocker E H F M Strong and W H Peterson 1955 *J Am Chem Soc* 77 34-356
- 98 Lemieux R U 1953 *Can J Chem* 31 396-417
- 99 Lim H 1953 *J Fac Ag Hokkaido Imp Univ (Sapporo Japan)* 37 16-209
- 100 Luz G 1934 *Phytopathol Z* 7 58-638
- 101 Mager J 1947 *Biochem J (London)* 41 603-609
- 102 Mager J and M Aschner 1947 *J Bacteriol* 53 283-29

- 103 Mangin L 1890 *Compt rend (Paris)* 110 644-647
- 104 Mangin L 1899 *J de Botan* 13 209-216 276-287 307-316 339-348 371-378
- 105 Mangin I 1910 *Compt rend (Paris)* 151 279-283
- 106 Mann T 1914 *Biochem J (London)* 38 345-351
- 107 Michel Durand E 1938 *Bull soc chim biol* 20 399-419
- 108 Mull R P and F F Nord 1911 *Arch Biochem* 5 283-290
- 109 Murray H C and F P Zscheile 1936 *Phytopathology* 46 363-366
- 110 Murray S M Woodbine and T K Walker 1933 *J Exp Botany* 4 251-256
- 111 Nabel K 1939 *Arch Mikrobiol* 10 515-541
- 112 Nelson C I 1933 *J Agr Research* 46 183-186
- 113 Nord F F H Hofstetter and E Dammann 1937 *Biochem Z* 993 231-255
- 114 Norkrans B 1900 *Symbolae Botan Upsalienses* 11 (1) 5-126
- 115 Norman A G 1935 *Ann Appl Biol* 20 146-164
- 116 Norman A G and W H Peterson 1932 *Biochem J (London)* 26 1946-1953
- 117 Norman A G W H Peterson and R C Houtz 1932 *Biochem J (London)* 26 1934-1945
- 118 Northcote D H 1933 *Biochem J (London)* 53 348-352
- 119 Northcote D H and R W Horne 1952 *Biochem J (London)* 51 231-236
- 120 Okazaki K S Koyama and O Tamemasa 1955 *J Pharm Soc Japan* 75 1090-1093
- 121 Oxford A F and H Raistrick 1933 *Biochem J (London)* 27 1176-1180
- 122 Peck R L 1947 In W J Nickerson (ed) *Biology of Pathogenic Fungi* p 167-188 Waltham Mass Chronica Botanica Co
- 123 Peck R L and C R Haulser 1938 *J Am Chem Soc* 60 2 99-2603
- 124 Pelletier R L and G W Keitt 1954 *Am J Botany* 41 362-371
- 125 Petermann M L M G Hamilton and H C Reilly 1932 *Arch Biochem Biophys* 3, 117-130
- 126 Peterson W H E B Fred and E G Schmidt 1922 *J Biol Chem A* 19-34
- 127 Pillai N C and K S Srinivasan 1936 *J Gen Microbiol* 14 248-255
- 128 Pinck I A and F E Allison 1914 *Soil Sci* 5, 155-161
- 129 Porges N 1932 *Botan Caz* 91 191-203
- 130 Porter J R 1916 *Bacterial Chemistry and Physiology* New York John Wiley and Sons pp 1073
- 131 Trill E A P R Wenck and W H Peterson 1933 *Biochem J (London)* 29 21-33
- 132 Pruess L M E C Fehinger and W H Peterson 1931 *Zentr Bakteriell Parasitenk Abt II* 89 310-317
- 133 Pruess L M H J Gorkica H C Greene and W H Peterson 1932 *Biochem Z* 246 401-413
- 134 Pruess L M W H Peterson H Steenbock and E B Fred 1931 *J Biol Chem* 90 369-384
- 135 Pyle A J H 1954 *J Gen Microbiol* 11 191-194
- 136 Raistrick H 1930 *Proc Roy Soc (London) B* 136 481-508
- 137 Raper A B and D I Fennell 1932 *Bull Torrey Botan Club* 79 25-51
- 138 Raveux R 1948 *Bull soc chim biol* 30 357-366
- 139 Rennerfeldt E 1934 *Planta* 22 221-239
- 140 Reuter C 1912 *Z physiol Chem Hoppe Seyler's* 78 167-215
- 141 Richards A G 1934 *Sci nce* 190 761-763
- 142 Riley R F and D K Miller 1918 *Arch Biochem* 18 13-20
- 143 Rippel A and C Behr 1930 *Arch Mikrobiol* 1 271-276
- 144 Rippel A and C Behr 1934 *Arch Mikrobiol* 5 561-577

- 143 Rolofsen P A and I Hoette 1931 *Antonie van Leeuwenhoek J Microbiol Serol* 17 297-315
- 146 Saito A 1953 *J Fermentation Technol (Japan)* 31 140-141 141-143 3 8-3 3 (*Chem Abstr* 48 5275-5276 1954)
- 147 Satina S and A F Blakeslee 1928 *Proc Nat Acad Sci U S A* 14 308-316
- 148 Schmidt C F Jr 1935 *J Biol Chem* 110 511-520
- 149 Schmidt D 1925 *Biochem Z* 158 223-222
- 150 Schmidt E G W H Peterson and E B Fred 1923 *Soil Sci* 15 479-488
- 151 Schmidt G 1931 In W D McElroy and B Glass (eds) *Phosphorus Metabolism* Vol 1 p 443-475 Baltimore Johns Hopkins Press
- 152 Schmidt M 1936 *Arch Mikrobiol* 7 241-260
- 153 Scholl E 1908 *Monatsh Chem* 29 1025-1037
- 154 Schönborn W 1932 *Arch Mikrobiol* 2 408-431
- 155 Schubert W J and F F Nord 1900 *J Am Chem Soc* 22 5337-5338
- 156 Schulz G 1937 *Planta* 27 196-218
- 157 Shu P and J A Thorn 1952 *Can J Botany* 30 2 26-36
- 158 Simonart P and K Y Chow 1934 *Antonie van Leeuwenhoek J Microbiol Serol* 20 174-180 210-216
- 159 Singh J S Shah and T K Walker 1956 *Biochem J (London)* 62 222-224
- 160 Singh J and T K Walker 1956 *Biochem J (London)* 62 286-289
- 161 Singh J T K Walker and M L Meara 1955 *Biochem J (London)* 61 85-88
- 162 Skinner C E 1934 *J Bacteriol* 28 99-106
- 163 Skinner J T W H Peterson and H Steenbock 1933 *Biochem Z* 267 169-178
- 164 Smith A M 1936 *J Soc Chem Ind (London)* 55 217T-221T
- 165 Smithies W R 1952 *Biochem J (London)* 51 259-264
- 166 Steinberg R A and J D Bowling 1939 *J Agr Research* 58 717-732
- 167 Stevens C M and P Vohra 1955 *J Am Chem Soc* 77 4935-4936
- 168 Stokes J L and M Guinness 1946 *J Bacteriol* 52 195-207
- 169 Strong F M and W H Peterson 1931 *J Am Chem Soc* 53 925-935
- 170 Subramaniam C V and K V Srinivasa Pai 1953 *Proc Indian Acad Sci* 37B 149-157
- 171 Sumi M 1928 *Biochem Z* 195 161-174
- 172 Takata F 1929 *J Soc Chem Ind Japan* 32 497-510 544-547 (*Chem Abstr* 23 4748 1929)
- 173 Tamura H 1912 *Advances in Enzymol* 2 183-238
- 174 Tanret C 1908 *Compt rend (Paris)* 147 75-77
- 175 Thatcher F S 1954 *Ann Rev Microbiol* 8 449-472
- 176 Thymann K V 1955 *The Life of Bacteria* New York The Macmillan Company pp 775
- 177 Thorn C and M Phillips 1932 *J Wash Acad Sci* 22 237-239
- 178 Thomas R C 1928 *Am J Botany* 15 537-547
- 179 Thomas R C 1930 *Am J Botany* 17 779-788
- 180 Thomas R C 1942 *Ohio J Sc* 42 60-6
- 181 Thomas R C 1943 *Ohio J Sc* 43 135-138
- 182 Todd G W and J Levitt 1931 *Plant Physiol* 26 331-336
- 183 Veibel S 1938 *Biochem J (London)* 32 1949-1953
- 184 Ward G E and G S Jamieson 1934 *J Am Chem Soc* 56 973-975
- 185 Ward G E L B Lockwood O E May and H T Herrick 1935 *Ind Eng Chem* 27 318-322
- 186 Wettstein F von 1921 *Sitzber Akad Wiss Wien Math naturw Kl Abt I* 130 3-20
- 187 Wieland P and V Prelog 1947 *Helv Chim Acta* 30 1028-1030

- 188 Woodbine M M E Gregory and T K Walker 1931 *J Exp Botany* 9
204-211
- 189 Woodman H E and R E Evans 1917 *J Agr Sci* 37 81-93
- 190 Woolley D W and W H Peterson 1936 *J Biol Chem* 114 85-90
- 191 Woolley D W and W H Peterson 1937a *J Biol Chem* 121 507-510
- 192 Woolley D W and W H Peterson 1937b *J Biol Chem* 121 213-218
- 193 Woolley D W F M Strong W H Peterson and E A Prill 1935 *J Am
Chem Soc* 57 2589-2591
- 194 Yamagata S 1934 *Acta Phytochim (Japan)* 8 107-116
- 195 Yarwood C E 1930 *Am J Botany* 37 636-639
- 196 Yarwood C E 1932 *Mycologia* 41 506-520
- 197 Zellner J 1907 *Chemie der höheren Pilze* Leipzig W Enchmann pp
257
- 198 Zellner J 1908 *Monatsh Chem* 29 45-51

3. Carbon Nutrition

1 PROBLEMS OF METHOD

Carbon compounds serve two essential functions in the metabolism of fungi as of all heterotrophic organisms. They supply in the first place the carbon needed for the synthesis of the compounds which go to make up the living cell—proteins, nucleic acids, cell wall materials, reserve foods, and so on—in a typical fungus about 50 per cent of the dry weight is carbon (Chapter 2). Second, the sole source of appreciable amounts of energy is the oxidation of carbon compounds, which may account for half or more of the carbon supplied to a culture.

Knowledge of carbon nutrition, therefore, is fundamental to an understanding of the physiology of the fungi. Much valuable work has been done in this area, but it will be apparent that the data required for confident generalization are usually still lacking, and that much of our knowledge is based on experiments which are open to criticism on the ground of method.

The study of the carbon nutrition of fungi, or of other microorganisms, involves general problems of two sorts: conceptual and methodological.

The concept of utilization may have more than one operational meaning. The commonest usage describes an experiment in which a small inoculum is placed in a medium containing one major source of carbon and all other materials necessary for growth; the organism

must initiate and maintain growth at the expense of the test compound. A second more narrow meaning of utilization is the ability of viable active mycelium to make further growth at the expense of a test compound. Here the design usually involves providing a small amount of a utilizable carbon source and measuring the additional growth if any obtained from a second compound. It is often found that materials which are not by themselves satisfactory carbon sources are used in this kind of experiment: examples in particular fungi include lactose (135) sugar alcohols (150) and cellobiose (142). As discussed later a compound which is used only in association with another carbon source is probably dependent for its metabolism on the induced synthesis of an adaptive enzyme system or upon the occurrence of mutants possessing the enzyme system.

So long as these two experimental designs are not confused each has its place. Confusion however does arise if an author proposes to use the first design—one compound as the sole source of carbon—but includes other carbon sources in the medium either as complex extracts or as amino acids. Unless the additional carbon is negligible and this must be shown by control experiments without carbohydrate there is the possibility that the experiment measures utilization in its second meaning rather than the one intended. Fortunately most fungi can be grown in a synthetic medium with an inorganic nitrogen source and with pure growth factors at levels which do not contribute significant amounts of carbon.

There are several methodological errors which are common enough to deserve comment. A cluster of such errors is associated with acidity. The initial pH should be determined after sterilization to guard against the possibility that compounds in the medium are broken down by autoclaving. Some compounds particularly organic acids may not enter the cell at the pH chosen in which case it is necessary to determine whether or not the organism can tolerate a pH at which penetration is more likely. Finally, the organism itself may produce organic acids which in a poorly buffered medium lower the pH so far as to restrict growth. Media may be buffered with phosphate for most organisms; for those fungi which cannot tolerate the level of phosphate required for effective buffer action (45) the pH may be controlled by addition of sterile solid calcium carbonate after autoclaving or by inclusion of an indicator and periodic adjustment of the pH with sterile alkali (58). The only way in which these problems can be detected in a particular instance is to determine the culture pH at frequent intervals and to exercise judgment in the interpretation of growth results which may be affected by pH changes.

It is possible that metabolites other than acids may accumulate in inhibitory concentrations for example the accumulation of ethanol has been suggested (181) as the explanation of thiamine toxicity to *Rhizopus suinus*

Another set of problems is associated with contamination of supposedly pure chemicals with growth active substances. There is evident in the literature an unjustified confidence in the label placed on a chemical by the manufacturer. Even compounds rated as reagent quality or the equivalent may be contaminated with biologically significant amounts of other substances particularly inorganic ions and organic growth factors. Such contamination is particularly likely in a compound derived from natural sources and most of the important carbon and organic nitrogen sources are so derived. In a highly purified medium one source of carbon may appear superior to another merely because it is contaminated with a required inorganic element in which the basal medium is partially deficient. More commonly the fungus has an unrecognized vitamin requirement which is met by the less pure of a series of carbon compounds thus the early and anomalous finding that *Diplodia macrospora* grows with starch maltose or sucrose but not with glucose or fructose is now known to reflect contamination of the first group of compounds with biotin required by the fungus as a growth factor (199-241). Growth factors for fungi have been found in many carbon compounds e.g. soluble starch (172) glucose (121) galactose (44) and saponin (19).

Growth inhibition by contaminants is less commonly encountered but has been reported from the ions present in crude natural materials (153) and from an unidentified alcohol soluble contaminant of glucose and galactose (44).

It should not be necessary to point out that such materials as cellulose and chitin as they occur in nature may be contaminated with organic materials in sufficient amount to provide the carbon source for visible growth of fungi. It is not for example justified to claim that an organism can utilize chitin merely on the evidence that it grows on the exoskeleton of an insect particularly since insect cuticle is less than 50 per cent chitin (239).

Breakdown of sugars during autoclaving constitutes another serious and often unrecognized source of error in studies of carbon nutrition. Growth inhibition from sugar breakdown products occurs in both bacteria and fungi (9-116-193). Breakdown is more extensive if the sugar is in contact during autoclaving with phosphate (59) or with amino acids (86-125). Ketohexoses e.g. fructose and sorbose break down (to 5-hydroxymethylfurfural) more rapidly than do the aldo

hexoses (138) and furfural formation from autoclaved xylose is sufficient to inhibit bacterial growth (120). Autoclaving of carbohydrate in the medium has been reported also to cause the formation of growth promoting substances for bacteria (86) and fungi (114).

Oligosaccharides may be partially hydrolyzed during autoclaving especially at low pH (7-22) and probably many claims of slight growth on oligosaccharides reflect merely the partial hydrolysis of a non-utilizable sugar to a utilizable monosaccharide.

It follows that sugars and other carbon sources should never be autoclaved in a medium. As a general rule they may be autoclaved separately and then added aseptically to the cooled basal medium. The expedient of intermittent sterilization is no better than autoclaving (136). The ultimate recourse is to filter sterilization and questionable results with a possibly labile sugar should always be checked with this method.

It is usual in carbon nutrition experiments to harvest all cultures after a predetermined, often arbitrarily chosen, incubation period. No objection can be made to this procedure if the only purpose of the experiment is to find a practicable growth medium for a specific purpose. If however it is intended to yield meaningful information on the intrinsic utilizability of compounds, some error may be introduced. One compound may be used with a very long lag, probably adaptive in nature (p. 86); the classification of this compound as utilizable or non-utilizable will depend on the length of the incubation period chosen. On the other hand, if time is allowed for slow utilization of compounds, growth on rapidly utilized substrates may have been completed and followed by extensive autolysis before the cultures are harvested. Ideally both these difficulties may be met by planning periodic harvests; comparison of compounds can then be made on the basis of two criteria: maximum growth and the time required to reach maximum growth. The data of Lilly and Barnett (116) show for many fungi the gain in understanding which flows from the use of more than one incubation period.

Some additional information and considerably greater confidence in interpretation may be gained by analysis of the medium at the time of harvest for residual carbon source. This procedure affords an additional check upon growth evidence, especially if growth is limited.

Differences in results between different laboratories are often casually explained as an expression of strain differences or as the result of variation. There is of course no doubt that strains of a given species may differ in their nutritional requirements and there are authenticated cases of metabolic changes resulting from spontaneous genetic

change. Nevertheless in the absence of adequate evidence these phenomena should not be accepted as sufficient explanation of the failure of one laboratory to confirm the work of another. It is clear that extensive recourse to such explanation would mean that no study could be confirmed or contradicted by later evidence.

From this review of methodological problems a number of recommendations may be formulated. Although not of equal importance and not necessarily applicable to all cases they should be considered carefully in designing experiments on the carbon nutrition of fungi.

1 Complex natural products which supply carbon should if possible not be included in the basal medium.

2 A control series with only the basal medium should be included in every experiment. Failure to include such a control makes it almost impossible to detect limited utilization of the test carbon compound since it is not known to what extent impurities in the basal medium support growth.

3 Adequate replication of cultures is necessary; the number to be determined by the variance found in pilot experiments. Probably for most fungi a minimum of four replicates for each harvest is required.

4 The carbon source should always be autoclaved separately from the basal medium and added aseptically to it. If it is found to be labile to heat, it should be sterilized by filtration.

5 The pH of the medium should be determined electrometrically after all ingredients are added and periodically during growth.

6 The inoculum should be of minimum size to avoid carry-over of materials from the parent culture. In doubtful cases serial subculture on the same medium may be necessary. All cultures should be inoculated with the same amount (mass or number) of cells.

7 The measurement of growth must be objective and quantitative, e.g. by dry weight, total nitrogen or turbidity (Chapter I).

8 For fungi which have rapid rates of growth and autolysis or which grow adaptively on some substrates it may be necessary to construct a growth curve for each carbon source or at least to make more than one harvest.

9 Analysis of the spent medium for residual carbon source is highly desirable although not always practicable.

10 Possible interactions between carbon and nitrogen sources should be considered; in some cases it may be necessary to study all putative carbon sources in media with different nitrogen sources.

11 When applicable the distinction between initiation of growth

and maintenance of preformed growth should be a factor in the interpretation of results

12 Reasonable precautions should be taken against complications arising from (a) contamination of the carbon source with impurities which affect growth or (b) the formation of toxic metabolites during growth. Hydrogen ion is the most common toxic metabolite and consequently the most important precaution in this connection is the periodic determination of the pH of the culture during the growth period

The most frequent and perhaps the least excusable deficiency in experiments on carbon nutrition is failure to meet the criterion of objective and quantitative measurement of growth. As a rule it is wise to disregard results which are reported in qualitative terms only although the complete absence of growth in such experiments may on occasion be significant. This stricture applies with almost equal force to the evaluation of growth on the basis of linear extension of a colony on agar: this type of growth measurement has its uses (Chapter 1) but is not sufficiently accurate to assess the relative utilizability of different carbon sources

2 MONOSACCHARIDES AND THEIR DERIVATIVES

Hexoses Of the 6-carbon sugars the hexoses D-glucose is biologically the most important and is utilized for growth by virtually all cultivable fungi. A few forms have been reported not to grow with glucose: the only convincing evidence so far presented is the qualitative observation of Schade (175) that *Leptomitius lacteus* uses none of the common hexoses but grows with acetate or other fatty acids as the sole source of carbon. This pattern resembles that of the acetate flagellates e.g. *Polytomella caeca* in which it has been suggested (122) that enzymes necessary for the phosphorylation of glucose are lacking. A similar deficiency in *Pseudomonas putrefaciens* has the same effect (106).

For the great majority of fungi and actinomycetes D-fructose and D-mannose are equivalent to glucose for growth. Occasionally as in *Stachybatrys atra* (99) one of these may appear superior to glucose for unknown reasons. Particularly among the lower fungi there appear to be numerous exceptions to the general rule that fructose and mannose are equivalent to glucose. In conventional experiments of limited duration it is found that either or both of these sugars fail to support the growth of several members of the Chytridiales (2, 45) and


Blastocladales (91-123) Less exhaustive studies suggest that the same is true of some species of the Saprolegniales (17-231) Of several reports of failure of higher fungi to utilize fructose that of Shirakawa (184) is the most convincing although the fructose was autoclaved in the medium Several isolates of *Streptomyces* fail to grow in an agar medium with fructose as carbon source (14)

Considerable light has recently been thrown on the non utilization of mannose and fructose through the work of Sistrom and Machlis (185) on *Allomyces macrogynus* one of the species of the Blastocladales previously found not to grow appreciably on these sugars in experiments of conventional duration It is now clear that *A. macrogynus* can grow on both mannose and fructose with a very long period of incubation the final growth is comparable to that on glucose In addition provision of a small amount of glucose to a medium in which the main carbon source is mannose or fructose allows normal growth with no delay Both of these observations are consistent with the hypothesis that this fungus—and by implication perhaps others—utilizes fructose and mannose only under conditions which permit or encourage the formation of an adaptive or induced enzyme although other observations remain to be reconciled with this hypothesis It would seem therefore that utilization of these monosaccharides is dependent on an enzyme system which is constitutive in some organisms and inducible in others strains of a given species may fall into different groups in this regard

Data from a comprehensive study of 57 fungal species mostly of the Fungi Imperfecti (116) show that relatively few—about one fifth of the sample—are completely unable to grow with D galactose as carbon source in a medium containing asparagine and salts Many especially in the genus *Fusarium* grow with galactose as well as with glucose Other fungi e.g. *Helminthosporium sativum* reach the same final dry weight as with glucose but reach it slowly suggesting that here too an adaptive mechanism is at work and that the inclusion of asparagine in the medium may have biased the results by providing a substrate able to support the formation of induced enzymes Finally for many fungi galactose is clearly utilized but does not even in a long time period support the same level of growth as does glucose We may conclude from this and other work that galactose is used by most fungi but is not usually so good a source of carbon as glucose A substantial number of fungi are unable to utilize it and it may even be toxic to a few (90-194) The ability or inability to utilize galactose bears no apparent relation to taxonomic position

On comparative grounds it seems probable that the ability of an

organism to use fructose mannose or galactose depends upon its ability to convert the sugar in question into a phosphorylated derivative of glucose able to enter the main respiratory pathways. The metabolic relationships of these four sugars in yeast and animal cells are shown in the diagram (67)



From this diagram it may be suggested that in adaptive growth on fructose mannose or galactose the enzyme actually lacking in the cell and induced by the environment is the specific kinase responsible for the phosphorylation of the free sugar. However no data are available and other possibilities and other pathways are of course conceivable. The occurrence of mannose 1 phosphate in the mycelium of *Penicillium chrysogenum* (105) for example suggests a pathway of mannose utilization similar to that of galactose in the scheme outlined above.

Studies on the economic coefficient (p. 89) and similar indices of growth indicate that poorly utilizable monosaccharides are such because they are not attacked or do not penetrate the cell rather than because once attacked they do not provide energy or usable metabolic products. This is especially clear from experiments with *Aspergillus niger* for which the small amount of a poor carbon source which disappears from the medium is as efficient in supplying carbon for growth as an equivalent amount of glucose (209, 209). The sugars which do not support growth adequately remain that is in the medium. Whether this is the result of specific enzymatic deficiencies as suggested above or whether it is to be attributed to slow penetration of the sugar remains to be determined. The two possibilities are not necessarily mutually exclusive if penetration of sugars depends as it may on a surface enzymatic reaction. It is perhaps significant that the yeast cell is impermeable to sorbose galactose and arabinose (171) all of which are generally poor sources of carbon for microorganisms.

1. Sorbose a ketohexose differing from fructose in the configuration of carbon atom 5 supports normal growth of a few fungi and actinomycetes (14, 116, 116). Although unequivocal data are not available it appears not to be utilizable by most forms and it is definitely toxic to

some The toxicity is evidenced by death of the hyphal tips followed by branching of the mycelium below the killed portion (116) Fungi e.g. *Neurospora crassa* which normally spread widely on an agar plate grow as small colonies in a medium with sorbose (204) The fundamental basis of sorbose toxicity is not yet known although often more pronounced in a maltose medium toxicity to some fungi is equally well expressed in glucose media so that interference with maltose utilization cannot be the explanation Sorbose inhibition occurs only at relatively high concentrations and is strongly affected by temperature (116) A possible line of approach to the problem of its mode of action may be found in the hypothesis that sorbose interferes with a respiratory pathway organisms dependent on this pathway would then be inhibited completely fungi which have alternate pathways would be less or not at all affected

Heptoses The naturally occurring 7 carbon sugars have not been tested as carbon sources for fungi except for the report (195) that *Aspergillus niger* does not grow with D-mannoheptulose

Pentoses D-Xylose is the most generally utilizable of the pentoses and has even been reported to be superior to glucose for some organisms (12 150 216) Some fungi and actinomycetes grow poorly or not at all with xylose and species within the same genus may differ markedly (195 201) It must be remembered that xylose is known to be broken down to furfural during autoclaving for which reason all reports of its non utilizability should be regarded with some reserve until confirmed with filter sterilized xylose *Phymatotrichum omnivorum* makes better growth with alcohol sterilized than with autoclaved xylose (18) but the poor growth of *Venturia inaequalis* strains on agar with xylose is not improved by filter sterilization of the sugar (113)

Both isomers of arabinose occur in nature and are commercially available Probably in those publications in which the isomer used is not specified the L isomer which is more easily available was used *Penicillium urticae* utilizes D-arabinose much less well than it does L-arabinose (57) In most studies L-arabinose is inferior to xylose and glucose as a carbon source (116a) In a medium containing a second carbon source however several fungi e.g. *Endoconidiophora adiposa* utilize L-arabinose with a long incubation period apparently adaptively (116)

Relatively few studies on rhamnose and methyl pentose have been made but the indications are that it is used by only a few organisms including *Stachybotrys atra* (99) and *Streptomyces coelicolor* (39)

Although *Penicillium urticae* grows with D-ribose as carbon source (57) it and the other naturally occurring pentoses and methyl pentoses have not been investigated thoroughly. Early studies not all satisfactory in their methods on pentose utilization by fungi are reviewed by Hawkins (80) and by Peterson et al (155).

It seems in conclusion that xylose is often a good source of carbon for fungi and that contrary findings should be examined closely. The pathway of utilization of xylose is not known; it seems likely that it enters the phosphogluconate oxidation pathway (Chapter 7) as pentose phosphate. L-Arabinose is used by several fungi, probably not by most. The other pentoses and methyl pentoses have been investigated rather infrequently, but they appear in general to be poor sources of carbon or not utilizable at all.

Other Monosaccharides A single study of a 4 carbon sugar (195) indicates that D-threose is not utilized by *Aspergillus niger*.

The L isomers of the naturally occurring aldohexoses and of xylose do not support growth of fungi; nor do those monosaccharides which do not occur in nature (99, 195, 201). Acetylation or methylation of the ring carbons of glucose yields derivatives which are not utilized by representative fungi (187), although *Aspergillus niger* has been reported to make a limited growth after adaptation with pentaacetyl glucose (8). *Aspergillus niger* also grows with 2-deoxyglucose as carbon source (8).

Polyhydric Alcohols Reduction of the sugars yields sugar alcohols; e.g. D-sorbitol arises by reduction of D-glucose, D-mannitol by reduction of D-fructose or D-mannose. In addition to the hexitols derived from the hexose sugars, tetritols, pentitols, and heptitols related respectively to 4, 5, and 7 carbon sugars occur in nature. Glycerol may be considered as a reduction product of a 3 carbon sugar.

Mannitol is the most generally utilizable of the hexitols and is equivalent to glucose for many fungi and actinomycetes (39, 100, 144, 180). However, many fungi with generally wide substrate ranges grow only poorly or not at all with mannitol as sole carbon source (65, 129, 152). In the lower phycomycetes a few forms grow poorly with mannitol (17, 32); most do not utilize it at all.

Sorbitol and dulcitol are unavailable to most fungi and actinomycetes. From the limited data it appears that sorbitol, the reduction product of glucose, is more generally used than is dulcitol, which is related chemically to galactose. It may be significant that many organisms able to utilize dulcitol are also able to grow more or less well

with galactose (17 180 201) Erythritol is apparently only rarely utilizable although the data again are extremely limited

Glycerol is often the best carbon source of all for actinomycetes (40 43 221) In the fungi the ability to utilize glycerol appears to be specific thus among fairly omnivorous saprophytes *Aspergillus oryzae* and *A. niger* make only poor growth (195 201) but *Memnoniella echinata* grows well with glycerol as sole carbon source (150)

Although the data are insufficient for generalization it may tentatively be concluded that the only polyhydric alcohols available to many fungi are glycerol and mannitol These may be expected on oxidation to yield respectively fructose and glyceraldehyde or dihydroxyacetone all of which can after phosphorylation enter known respiratory pathways

Sugar Acids Oxidation of the aldehyde carbon of an aldohexose yields an acid D-gluconic acid from D-glucose is the only one to have been investigated as a source of carbon for fungi As might be expected from the fact that many strains of *Aspergillus niger* form and then consume gluconic acid during sugar metabolism (Chapter 6) it is a good source of carbon for *A. niger* (46) and *A. oryzae* (201) although the strain of *A. niger* studied by Steinberg (195) could not grow with gluconate as sole carbon source Several other organisms grow on gluconate (40 99 121) but others e.g. *Chalara quercina* do not utilize it (12) Growth with gluconate is probably often limited by an unfavorably high pH resulting from utilization of the neutralized acid

Utilization of 2 ketogluconic acid by fungi is like that of gluconic acid highly specific (50) and is probably a function of the respiratory systems available

A single report (87) indicates without quantitative data that *Aspergillus niger* grows with D-glucuronic or D-galacturonic acids as sole carbon source

D-Saccharic acid the dibasic acid resulting from the oxidation of carbon atoms 1 and 6 of D-glucose has been reported a good source of carbon for *Aspergillus niger* (46) but a relatively poor source for *A. oryzae* (201)

Glycosides Limited studies of the methylglucosides indicate as would be expected from the relative ubiquity of β glucosidases (Chapter 5) that β methylglucoside is much more easily utilizable than the α -isomer (51 180) Amygdalin which is hydrolyzed to gentiobiose and mandelonitrile is utilized by several members of the Saprolegnia

ceae (17) in general it may be expected that the ability of an organism to use a given β glycoside will depend on its ability to use the sugar which is split off from it. The cyanogenetic glycosides however may not be utilisable by reason of the toxicity of the cyanide liberated on hydrolysis (171).

3 OLIGOSACCHARIDES

Chemically the oligosaccharides are glycosides composed of two or more monosaccharides and are characterized by the nature of the component sugars, the number of units, and the type of glycosidic bond. Those which are important in the nutrition of fungi include five disaccharides—maltose, cellobiose, trehalose, sucrose, and lactose—and one trisaccharide, raffinose, which is made up of galactose, glucose, and fructose units. In addition to data on specific sugars, two general problems are relevant at this point: (1) the enzymatic basis of disaccharide utilization, and (2) adaptive growth with oligosaccharides.

Maltose occurs in nature as a product of starch hydrolysis and is utilized by virtually all fungi which have been tested, even those which have a limited substrate range. A few fungi are known on the basis of quantitative studies to grow poorly or not at all with maltose: *Clytridium* sp. (15), *Polychytrium aggregatum* (2), *Entomophthora* (*Empusa*) sp. (242), and *Penicillium digitatum* (65, 244). Surprisingly, the two chytrids mentioned make at least some growth with starch; the other fungi use neither starch nor maltose. Maltose is more easily hydrolyzed by autoclaving than are sucrose or lactose (136), but even with this possible source of error it seems that it is very generally available to fungi.

Cellobiose, like maltose, is composed of two glucose residues, but the linkage is of the β configuration. The data are more limited than those for maltose, but indicate that it is almost as widely utilisable as maltose. Indeed, two of the fungi mentioned above, which do not grow with maltose, do with cellobiose. *Trichosoma flavobrunneum* is among the small group of tested fungi which fail to utilize cellobiose, but other species of the same genus grow well with it as sole carbon source (142). The utilisability of cellobiose reflects the general occurrence of β glucosidases in fungi (Chapter 5).

Trehalose, a glucose-containing disaccharide produced by fungi and a few higher plants (Chapter 5), appears from the few studies which have been made to be generally available, even to rather fastidious fungi (10, 99, 201, 212).

Sucrose, the characteristic sugar of higher plants, is generally a good

source of carbon for fungi but is not so nearly universally available as maltose. Thus of 57 fungi studied by Lilly and Barnett (116) only 1 were completely unable to utilize sucrose but several grew slowly with it. Among the major groups of fungi failure to utilize sucrose appears to be particularly common in the Chytridiales (Table 1) and the Mucorales (174) less common in other groups. Actinomycetes rarely if ever utilize sucrose (15, 221).

Lactose the β -galactosidoglucose of mammalian milk is used by far fewer fungi than any of the disaccharides so far mentioned and may be characterized as a poor carbon source (61, 116). At least some actinomycetes grow with lactose (40, 51, 215). Adaptation to lactose occurs among fungi (p. 87) and whether or not an organism is reported as able to utilize the sugar often therefore depends on the length of the incubation period chosen.

Melibiose an α -glycoside has not been investigated often; the few data available (Table 1) indicate that it is not utilizable by many fungi. Data for gentiobiose are similarly too limited to allow generalization. Growth on amygdalin the β -glycoside of gentiobiose by aquatic fungi (17) presumably indicates gentiobiose utilization.

Raffinose attacked at one linkage yields galactose and sucrose; attacked at the other yields fructose and melibiose. From the limited data of Table 1 and the more extensive studies of Lilly and Barnett (116) it would appear that the majority of fungi utilize raffinose but that non-utilization is common. Some doubt is raised with regard to raffinose utilization by the finding (18) that raffinose supports growth of *Phymatotrichum omnivorum* if autoclaved but not if sterilized by alcohol treatment; a reinvestigation of this problem by comparing filter-sterilized to heat-sterilized raffinose may show that raffinose is broken down by autoclaving at least to the extent that some simple sugars are available to initiate growth and to permit therefore the formation of adaptive enzymes which then hydrolyze the unchanged raffinose.

The utilization of an oligosaccharide is believed to be preceded by and dependent on its conversion to hexoses or possibly hexose phosphates. The evidence for this generalization may be summarized:

1. Fungi that can use a disaccharide possess enzymes which hydrolyze it (17, 79). An organism which grows on sucrose for example proves on examination to form sucrase. Formation of the enzyme is not however a sufficient condition for utilization—a mutant of *Neurospora crassa* produces lactase (β -galactosidase) but does not grow with lactose as sole carbon source (112).

Table 1 Carbon Nutrition of Some Fungi and Actinomycetes*

Carbon Source	<i>Chytridium</i> sp (45)	<i>Macrochytrium</i> <i>botrydoides</i> (45)	<i>Polychytrium</i> <i>aggregatum</i> (2)	<i>Allomyces</i> <i>javanicus</i> (94)
None	4	1	0	0
Glucose	100	100	100	100
Fructose	101	1	0	0
Mannose	101	120		103
Galactose	99	23	0	0
Sorbose	8	1		
Xylose	7	103	0	0
L Arabinose	4	1	0	
D Arabinose				
Ribose				
Rhamnose	4	1		
Mannitol			0	
Sorbitol				
Dulcitol	4	2		
Adonitol				
Erythritol				
Glycerol	17	9		0
Maltose	10	116	0	103
Sucrose	9	4	0	0
Cellobiose	100	137	0	
Lactose	3	4	0	0
Trehalose				
Melibiose	5	2		
Raffinose	5	2	0	0
Glycogen	94	124		
Starch	50	106	+	+
Dextrin				
Inulin	3	3		
Cellulose	3	+	0	
Formate				
Acetate	0	<3		0
Propionate				
Oxalate				
Lactate				0
Succinate	0	<3		0
Fumarate				0
Malate				
Tartrate	0	<3		
Citrate	0	<3		
Glucuronate				

* All data adjusted by taking growth on glucose as 100. Organisms arranged in order: phytomyces, ascomycetes, basidiomycetes, Fungi Imperfecti, actinomycetes. Tr = trace of growth, + = visible growth, not measured.

Table 1 (continued)

Carbon Source	<i>Allomyces</i> <i>ja anicus</i> <i>macrogyms</i> (123)	<i>Blastocladia</i> <i>prinsheimii</i> (32)	<i>Brevilegnia</i> <i>gracilis</i> (17)	<i>Saprolegnia</i> <i>delica</i> (17)
None	0	5	0	0
Glucose	100	100	100	100
Fructose	0	88	81	102
Mannose	0	97	94	0
Galactose	0	7	86	0
Sorbose	0	7		
Xylose	0	6	0	0
L Arabinose	0	7	0	0
D Arabinose				
Ribose				
Rhamnose			0	0
Mannitol			52	0
Sorbitol			89	29
Dulcitol			0	0
Adonitol				
Erythritol			0	0
Glycerol		39	86	14
Maltose	107	101	88	107
Sucrose	0	99	83	14
Cellobiose	0			
Lactose	0	6	61	0
Trehalose				
Melibiose	0			
Raffinose			76	0
Glycogen	50		76	93
Starch	54	95	76	100
Dextrin	98	87	86	86
Inulin	0		40	0
Cellulose		6		
Formate		5		
Acetate	37	5		
Propionate				
Oxalate				
Lactate				
Succinate	0	9		
Fumarate				
Malate		5		
Tartrate		5		
Citrate	0	6		
Gluconate				

Table 1 (continued)

Carbon Source	<i>Pytophthora cactorum</i> (129)	<i>Pythium</i> <i>sp</i> (33)	<i>Chalara quercina</i> (12)	<i>Coprinus</i> <i>sp</i> (100)
None	0	3	2	34
Glucose	100	100	100	100
Fructose	50	104	119	107
Mannose	38		96	127
Galactose	63	10	115	76
Sorbose				
Xylose	42	64	215	52
L Arabinose	55	7	46	56
D Arabinose				
Ribose				
Rhamnose	58		13	
Mannitol	63	±10	13	111
Sorbitol	92			
Dulcitol	55		2	
Adonitol				
Erythritol	52		2	
Glycerol	102	5	15	
Maltose	125	99	133	105
Sucrose	133	101	94	84
Cellobiose		101	119	
Lactose	83	36	4	72
Trehalose				
Melibiose		5	27	
Raffinose	63		100	96
Glycogen				
Starch	125	98	6	112
Dextrin	63		217	120
Inulin	83		33	77
Cellulose		+	23	
Formate		0		
Acetate		0		
Propionate				
Oxalate		0		
Lactate		0		
Succinate		0	15	
Fumarate				
Malate		0		
Tartrate		0		
Citrate		0		
Gluconate		0	2	

Table 1 (continued)

Carbon Source	<i>Psalliota bispora</i> (216)	<i>Stereum gaucapatum</i> (82)	<i>Ustilago violacea</i> (19)	<i>Fusarium oxysporum f. nicotianae</i> (243)
None		12	5	
Glucose	100	100	100	100
Fructose	95	104	99	104
Mannose		104		109
Galactose	73	89	63	78
Sorbose				59
Xylose	132	123	3	135
L Arabinose	109	43	5	68
D Arabinose				14
Ribose				74
Rhamnose		41	7	39
Mannitol	20		93	41
Sorbitol			90	24
Dulcitol				27
Adonitol				7
Erythritol				
Glycerol		24	24	48
Maltose	73	160	95	138
Sucrose	68	158	88	113
Cellobiose			78	108
Lactose			11	19
Trehalose				111
Melibiose				23
Raffinose		126	8	120
Glycogen		112		
Starch	20	160	10	59
Dextrin			25	105
Inulin	23	70	9	
Cellulose				
Formate				
Acetate	11			
Propionate				
Oxalate	100			
Lactate				
Succinate				
Fumarate				
Malate	116			
Tartrate	32			
Citrate				
Gluconate				

Table 1 (continued)

Carbon Source	<i>Pytophthora cactorum</i> (129)	<i>Pythium sp</i> (33)	<i>Chalara quercina</i> (12)	<i>Coprinus sp</i> (100)
None	0	3	2	34
Glucose	100	100	100	100
Fructose	50	104	119	107
Mannose	38		96	127
Galactose	63	10	115	76
Sorbose				
Xylose	42	64	215	52
L Arabinose	55	7	46	56
D Arabinose				
Ribose				
Rhamnose	58		13	
Mannitol	63	±10	13	111
Sorbitol	92			
Dulcitol	55		2	
Adonitol				
Erythritol	52		2	
Glycerol	102	5	15	
Maltose	125	99	133	105
Sucrose	133	101	94	84
Cellobiose		101	119	
Lactose	83	36	4	72
Trehalose				
Melibiose		5	27	
Raffinose	63		100	96
Glycogen				
Starch	125	98	6	112
Dextrin	63		217	120
Inulin	83		33	77
Cellulose		+	23	
Formate		0		
Acetate		0		
Propionate				
Oxalate		0		
Lactate		0		
Succinate		0	15	
Fumarate				
Malate		0		
Tartrate		0		
Citrate		0		
Gluconate		0	2	

Table 1 (continued)

Carbon Source	<i>Stachybotrys atra</i> (92)	<i>Streptomyces coelicolor</i> (40)
None		17
Glucose	100	100
Fructose	241	79
Mannose	66	100
Galactose	128	84
Sorbose	0	26
Xylose	94	143
L Arabinose	25	65
D Arabinose	22	
Ribose	9	
Rhamnose	91	
Mannitol	100	82
Sorbitol	0	27
Dulcitol	28	20
Adonitol	25	
Erythritol	0	
Glycerol	166	135
Maltose	131	62
Sucrose	78	34
Cellobiose	72	81
Lactose	84	
Trehalose	59	90
Melibiose		
Raffinose	56	
Glycogen		
Starch	66	107
Dextrin	53	32
Inulin	113	
Cellulose	+	
Formate	0	
Acetate	22	33
Propionate	tr	
Oxalate	tr	
Lactate	tr	60
Succinate	tr	47
Fumarate		69
Malate	3	60
Tartrate	0	20
Citrate	tr	24
Gluconate	9	82

Table 1 (continued)

Carbon Source	<i>Memnoniella echinata</i> (150)	<i>Penicillium digitatum</i> (65)	<i>Sclerotium delphinii</i> (151)
None			
Glucose	100	100	100
Fructose	110	108	113
Mannose	90	108	
Galactose	105	99	
Sorbose			
Xylose	145	102	76
L Arabinose	105	95	98
D Arabinose			
Ribose			
Rhamnose		65	
Mannitol	90		
Sorbitol	50		
Dulcitol			
Adonitol			
Erythritol			46
Glycerol	85		104
Maltose	105	21	152
Sucrose	105	92	147
Cellobiose	105	101	
Lactose	120	64	
Trehalose	45		
Melibiose			
Raffinose	100		
Glycogen			
Starch	170	2	195
Dextrin	140		
Inulin		1	191
Cellulose			
Formate	45		
Acetate	35		43
Propionate	40		
Oxalate	25		
Lactate			
Succinate		2	
Fumarate		1	
Malate		21	
Tartrate	10		
Citrate		28	
Gluconate			

Direct utilization may of course occur in the fungi as it does in bacteria (53) but the evidence is not yet at hand

The second general problem in oligosaccharide utilization is that of adaptive growth. Operationally nutrition experiments often show that a given organism grows on a disaccharide only after a long incubation period but that once growth is initiated the final crop is as heavy as that produced with glucose or other easily utilizable carbohydrates. The study of Lilly and Barnett (116) contains many examples of this phenomenon which as discussed later is probably to be ascribed to the formation of an induced enzyme system.

4 POLYSACCHARIDES

Polysaccharides as the name implies are polymers of sugars or sugar derivatives. Although the number of units may be as small as 10 the natural polysaccharides have as a rule very high molecular weights and contain up to several thousand saccharide units. They can be divided into two broad classes: reserve or nutrient polysaccharides (e.g. starch, glycogen and inulin) and structural polysaccharides (e.g. cellulose, pectic materials and chitin). The structural polysaccharides found in cell walls and other extraprotoplasmic structures are generally insoluble linear polymers; the reserve polysaccharides contain a large percentage of branched chain structures and are usually more soluble (235). Many polysaccharides are too poorly known to be fitted into either category.

Starches and Related Polysaccharides Starch, the principal reserve polysaccharide of plants (actually the starches of different plants are somewhat different) is an excellent carbon source for most fungi and actinomycetes even for rather fastidious forms. A few fungi are known on acceptable evidence not to grow with starch as the sole carbon source. They include representatives of widely different taxonomic and ecological groups: *Ustilago violacea* (180), *Rhizophlyctis rosea* (160), *Psalliota bispora* (*P. campestris*) (216), *Tricholoma imbricatum* (142), *Entomophthora (Empusa) spp.* (242), *Penicillium digitatum* (65) and *Chalara quercina* (12). The failure of *Penicillium digitatum* to grow with starch finds its explanation in an inability to form amylase (88).

As would be expected from the fact that starch is hydrolyzed primarily to maltose, really good growth on starch is accompanied by the ability to grow well on maltose (142) although a *Chytridium* sp. unable to utilize maltose makes limited growth on starch (45).




Figure 1 The growth of *Schizothecium longicollie* on disaccharides and on equivalent mixtures of their constituent hexoses. From tabular data of Lilly and Barnett (116)

2 The rate of growth of a lactase-rich mycelium on lactose is demonstrably higher than that of a lactase-poor mycelium (112)

3 Fungi which cannot use a disaccharide have been shown to use its constituent monosaccharides (Figure 1). That is the only bar to utilization appears to be inability to convert the compound sugar to hexoses

It is often held that direct utilization of oligosaccharides without prior splitting of the glycosidic bond occurs in fungi. Data cited in support of this claim include (1) quantitative superiority of a disaccharide to its constituent monosaccharides (78-82), (2) differences in metabolic processes during disaccharide and monosaccharide utilization (52-115) and (3) utilization of a disaccharide by intact cells at a pH at which the corresponding hydrolytic enzyme is inactive *in vitro* (141). All of these phenomena, however, can be explained without recourse to the hypothesis of direct utilization; some of the problems as they have been studied in yeast are reviewed by Hestrin (83-84).

in wood (85 169) and in other plant materials (143 179) the extensive early literature on hemicellulose breakdown in soil is reviewed by Waksman and Diehm (225)

Purified pentosans are attacked by most fungi and actinomycetes which have been tested (80 99 124 216 225)

Cellulose Cellulose the principal structural polysaccharide of plants is by far the largest reservoir of biologically utilizable carbon on the surface of the earth and its decomposition in which fungi probably play a major role is of great ecological importance Structurally it is a linear polymer of β -glucose the configuration being that of a $1,4\beta$ glucoside The number of units in the polymer is high probably individual linear chains are connected by cross linkages Complete hydrolysis yields glucose partial hydrolysis the disaccharide cellobiose

The literature on the utilization of cellulose by fungi is very large beginning with the observations of the pioneer plant pathologists of the nineteenth century Unfortunately frequent reliance on inadequate criteria makes it difficult to evaluate this work and much of it must be considered of doubtful validity The early literature is reviewed by McBeth and Scales (123) Otto (147) and Waksman (223) The monograph of Siu (186) especially concerned with fabric deterioration offers the most complete survey available of more recent work

Several methods have been used in attempts to assay the ability of fungi to grow on or destroy cellulose Some of these e.g. observation of the erosion of cotton fibers (56 147) and measurement of the loss in breaking strength of rotted cotton fabric (73 214) are adequate for their purpose but do not measure growth directly Similarly measurement of gas exchange during growth on cellulose (55 107) determines an activity which is not necessarily a function of growth

Visual estimate of growth on filter paper or other cellulosic materials is by itself of no value as an index of growth or cellulolytic capacity (126) Growth on reprecipitated cellulose in agar (123) is equally unreliable probably because of chemical changes occurring in the preparation of the cellulose (143)

The quantitative methods which seem most promising are the following

- 1 Measurement of growth on or hydrolysis of carboxymethylcellulose a soluble cellulose derivative (96) However the correlation of this property with cellulolytic capacity is uncertain (163)

- 2 Chemical determination of the disappearance of cellulose from complex natural materials This has been applied extensively to

Most investigators have used soluble starch a partially depolymerized derivative prepared by acid treatment rather than native starch. Raw starch unless the granule structure is destroyed by autoclaving will not support the growth of *Phymatotrichum omnivorum* (18) and indeed is not hydrolyzed by purified amylase. This raises of course the question of the utilization of plant starches in nature by plant pathogens and by saprophytes acting on unmodified plant materials.

Starch is often a better substrate for growth than glucose although no such case has been studied in detail; this effect presumably results either from contamination of the starch with growth factors or from the fact that utilization of a slowly hydrolyzable compound is accompanied by less accumulation of acids than utilization of glucose.

Dextrins which are chemically or enzymatically modified starches of uncertain structure are generally satisfactory sources of carbon for fungi. Quantitative differences between starch and dextrins as carbon sources are frequently reported but cannot easily be interpreted until the chemistry of these polysaccharides is better known. *Chalara quercina* unable to utilize starch grows very well on dextrin (12).

Inulin a polymer of D-fructose possibly containing a small amount of D-glucose is found in several of the Compositae e.g. *Dahlia*. It is not hydrolyzed by enzymes which act on starch or on sucrose. Limited data suggest that it is a good carbon source for many but not all fungi. It should however be noted that inulin like all fructans is susceptible to hydrolysis by heat and in careful work it is necessary to use filter sterilization (18).

Unidentified bacterial fructans are probably utilized by *Streptomyces coelicolor* and related species (127).

Glycogens the reserve polysaccharides of animals and some thallophytes are glucose polysaccharides quite similar to starch or more precisely to the amylopectin fraction of starch. Glycogens have not been tested extensively in the nutrition of fungi but the data suggest that they are quite generally available to fungi of widely different ecological groups (17, 45, 82, 201). Those fungi unable to use starch can be expected not to use the glycogens since so far as is known the same enzymes attack both polysaccharides.

Hemicelluloses The hemicelluloses comprise a group of insoluble polysaccharides of plant cell walls. The group as a whole is not well defined and there is much confusion in the literature. The hemicellulose fraction as usually isolated includes pentosans, hexosans and polyuronides; the main constituent is probably a pentosan or mannan (235). Fungi are active in the breakdown of hemicelluloses.

Although early work reviewed by Waksman (224) suggested that most species of the Actinomycetales are able to attack cellulose the evidence must be considered inadequate. It is certain that some species of *Streptomyces* and at least one of *Micromonospora* decompose cellulose (56 93 143 162 222) but whether this is true of most species and whether the actinomycetes as a group play an important role in the decomposition of cellulose in nature remain to be determined.

Generalizations on the distribution of cellulolytic capacity among the different taxonomic groups are: it should be remembered based on limited data not only in the sense that often only a few isolates of a taxon have been tested but also in the sense that the conditions of test may not be favorable for all potentially cellulose utilizing forms. The utilization of cellulose is affected by such factors as the available nitrogen (108) the temperature (126) the acidity (162) and the presence of other carbon compounds (85 142). It seems likely therefore that cellulose is used by more fungi than present information would indicate.

The widespread occurrence of cellulolytic fungi in soil and the fact that addition of cellulosic materials to soil results in an increase in the number of fungi isolated have led most workers to believe that at least under aerobic conditions and at moderate temperatures the fungi play a more important part in the initial decomposition of cellulose in soil than do the cellulolytic bacteria. The procedures of soil microbiology are however so inexact and the total soil microflora is so poorly known that this supposition attractive as it is is not adequately supported by the evidence.

The fact of occurrence in nature on a cellulosic substrate even very frequent occurrence is distinctly not acceptable evidence of ability to utilize cellulose. Thus *Pullularia pullulans* is one of the organisms most frequently isolated from exposed cotton cloth but it is not cellulolytic (186).

Chitin Chitin the structural polysaccharide of higher invertebrates is also a constituent of many fungi (Chapter 2). Chemically it is a linear polymer of N-acetyl-D-glucosamine the units being linked by β 1 4 glycosidic bonds. Like cellulose it is stable and difficult to purify without alteration of structure.

Fungi are often found growing on the exoskeletons of living or dead Arthropoda. Although it is often assumed perhaps correctly that these fungi attack chitin this evidence by itself cannot establish the point. Visible growth on or erosion of chitin has been reported in studies on soil fungi and actinomycetes (189) and on aquatic fungi

wood (10 31 178) straw (143 222) litter (117) and other materials (68 198) It is the appropriate method for the practical problem involved in the decomposition of cellulose in nature but suffers of course from the inherent difficulty of all growth measurements in complex materials that the cellulose may not be the primary growth substrate

3 Determination of the loss in weight of cellulosic materials exposed to the test organism (162 233) The principal complication is that the mycelial weight is included with the residual cellulose alkali treatment (191) can reduce but not eliminate this error

4 Determination of total or protein nitrogen in the insoluble material left after growth of a test organism on cellulose in a medium with a soluble nitrogen source (142) This is without doubt the most accurate and reliable method for the measurement of real growth of a fungus on an insoluble non nitrogenous substrate such as cellulose Unfortunately the method has been little used and cannot as yet be compared to others in detail

The methods which have been used do not always agree thus *Aspergillus versicolor* strains appear to utilize the cellulose of straw (143) but apparently do not cause measurable deterioration of cotton fabric (126) A complete list of the fungi reported by various methods as able to attack cellulose has been compiled (186) It is apparent that certain genera of the Fungi Imperfecti and the ascomycetes are particularly rich in species able to decompose cellulose In the basidiomycetes the wood rotting and litter-decomposing fungi are generally although not universally able to decompose cellulose genera particularly involved include *Armillaria Clitocybe Collybia Corticium Forres Marasmius Mycena Polyporus Polystictus Schizophyllum* and *Stereum* (31 118 164) It must be noted however that the species within a given genus are not all necessarily alike in their response to cellulose This has been especially well established in *Aspergillus* (238) and *Tricholoma* (142) In some species strain differences in the utilization of carboxymethylcellulose have been reported (9b)

Among the phycomycetes many members of the Chytridiales utilize cellulose (45 233) Limited data indicate that none of the Blastocladales is cellulolytic (34) In the Saprolegniales members of several genera apparently produce an enzyme hydrolyzing filter paper cellulose (16 173) as do species of *Phytophthora* in the *Peronosporales* (128) this may be taken as *prima facie* evidence for their ability to grow on cellulose None of the Mucorales so far studied is cellulolytic (126 147 164 236)

essential ions in solution or limit growth by removing ions (183) which of these two opposite effects will occur is determined by the solubilities of the chelate complexes and the metallic salts at the culture pH chosen

The non fatty organic acids which are most often utilized by fungi are those e.g. citrate succinate and malate which are products of glucose metabolism. Most of the data are from studies on common saprophytes and it appears that utilization is highly specific. Thus *Penicillium janczewskii* grows well on succinate (23) succinate is a relatively poor carbon source for *P. javanicum* (121) and is not utilized at all by *P. digitatum* (65). Because of the likelihood that cells may be impermeable to acids especially hydroxy acids nutritional data have no necessary relevance to the acid metabolism of the cell.

Tartaric acid the utilization of which by a fungus is so important in the history of both crystallography and microbiology (148) is in general not a good carbon source for fungi although it is utilized as well as glucose by *Memnoniella echinata* (150). The growth of some common fungi with tartrate has been surveyed by Stadtman et al. (190) an adaptive tartrate-decomposing system of *Aspergillus versicolor* has been described (10). The dextrorotatory isomer is used preferentially (70).

Oxalic acid is used by few microorganisms among which are *Proactinomyces* sp. and a strain of *Aspergillus oryzae* (137) *Psalliota bispora* (216) and a few bacteria (104). Possibly the insolubility of calcium and magnesium oxalates results in a removal of these ions from solution and a consequent limitation of growth in oxalic acid media.

Malonic acid is used although it is not a favorable carbon source by a few fungi and actinomycetes (23 92 99).

Inclusive studies of polyfunctional organic acids in nutrition have been reported by Camp (30) Tamiya (201) and Steinberg (190).

The lower fatty acids—formic acetic propionic etc.—are as a rule poorly utilizable or even toxic. Propionic acid in fact is employed as a fungistatic agent. Some fungi however do make at least some growth with these acids particularly with acetate on a carbon basis acetate is equivalent to glucose for *Allomyces javanicus* (124). Qualitative data (175) indicate that *Leptomitia lacteus* can grow with any of the lower organic acids including some that are almost uniformly toxic to the higher fungi. Probably it will be found that the lower fungi utilize fatty acids better than the more extensively investigated higher forms. The utilization of acetate by *Penicillium chrysogenum* is strongly pH dependent (90).

(245) It must however be concluded that there is no quantitative and entirely satisfactory evidence for the utilization of chitin or of N-acetylglucosamine. The work just mentioned suggests that an investigation of this problem in the fungi would be rewarding. Chitin splitting enzymes must be present in those fungi which autolyze extensively and an exocellular chitinase is formed by *Streptomyces* sp and some fungi (Chapter 5).

Pectic Substances The pectic substances are polysaccharides of plant cell walls usually isolated as a mixture of which the major constituent is a polymer of methylated galacturonic acids (235). Very little attention has been devoted to these materials as carbon sources although pectin has been reported to provide carbon for the growth of *Sclerotinia cinerea* (240) and of *Psalliota bispora* (216). From the frequent occurrence in fungi of enzymes acting upon the pectic substances (Chapter 5) and from the employment of pectin as the carbon source in experiments on enzyme production it is probably safe to assume that they are utilized in nature by many fungi both parasitic and saprophytic.

5 ORGANIC ACIDS

Organic acids are generally poorer sources of carbon for fungi than are the carbohydrates. Apart from what might be called intrinsic utilizability however three factors interfere with the utilization of the organic acids in particular cases. First cells are often impermeable to organic acids at physiological pH levels. Impermeability is more often invoked than demonstrated but there is no doubt that it is a factor in the utilization of acids. Second the utilization of neutralized organic acids causes a rise of the culture pH which may interfere with growth. This phenomenon explains the repeated observation that growth with organic acids is better if nitrogen is supplied as an ammonium salt (115, 165, 207). Utilization of the ammonium cation tends to drive the pH down partially or wholly counteracting the alkalinizing effect of acid utilization. In addition it appears that the utilization of ammonium ion is more rapid if the medium is neutral or alkaline (131). It has been argued (24) that a more direct reaction exists between organic acids and ammonium ion but this appears unlikely and is not necessary in order to explain the observed phenomena.

A third and probably only occasionally important factor is the chelation of inorganic ions by certain of the organic acids e.g. citrate and tartrate. This may however either promote growth by keeping

essential ions in solution or limit growth by removing ions (183) which of these two opposite effects will occur is determined by the solubilities of the chelate complexes and the metallic salts at the culture pH chosen

The non fatty organic acids which are most often utilized by fungi are those e.g. citrate succinate and malate which are products of glucose metabolism. Most of the data are from studies on common saprophytes and it appears that utilization is highly specific. Thus *Penicillium janczewskii* grows well on succinate (23) succinate is a relatively poor carbon source for *I. javanicum* (121) and is not utilized at all by *P. digitatum* (65). Because of the likelihood that cells may be impermeable to acids especially hydroxy acids nutritional data have no necessary relevance to the acid metabolism of the cell.

Tartaric acid the utilization of which by a fungus is so important in the history of both crystallography and microbiology (148) is in general not a good carbon source for fungi although it is utilized as well as glucose by *Memnoniella echinata* (150). The growth of some common fungi with tartrate has been surveyed by Staduman et al. (190) an adaptive tartrate-decomposing system of *Aspergillus versicolor* has been described (10). The dextrorotatory isomer is used preferentially (70).

Oxalic acid is used by few microorganisms among which are *Proactinomyces* sp. and a strain of *Aspergillus oryzae* (137) *Psalliota bispore* (216) and a few bacteria (104). Possibly the insolubility of calcium and magnesium oxalates results in a removal of these ions from solution and a consequent limitation of growth in oxalic acid media.

Malonic acid is used although it is not a favorable carbon source by a few fungi and actinomycetes (23, 92, 99).

Inclusive studies of polyfunctional organic acids in nutrition have been reported by Camp (30), Tamura (201) and Steinberg (190).

The lower fatty acids—formic, acetic, propionic etc.—are as a rule poorly utilizable or even toxic. Propionic acid in fact is employed as a fungistatic agent. Some fungi however do make at least some growth with these acids particularly with acetate on a carbon basis acetate is equivalent to glucose for *Allomyces javanicus* (124). Qualitative data (175) indicate that *Leptomitopsis lacteus* can grow with any of the lower organic acids including some that are almost uniformly toxic to the higher fungi. Probably it will be found that the lower fungi utilize fatty acids better than the more extensively investigated higher forms. The utilization of acetate by *Penicillium chrysogenum* is strongly pH dependent (95).

(2 45) It must however be concluded that there is no quantitative and entirely satisfactory evidence for the utilization of chitin or of N acetylglucosamine. The work just mentioned suggests that an investigation of this problem in the fungi would be rewarding. Chitin splitting enzymes must be present in those fungi which autolyze extensively and an exocellular chitinase is formed by *Streptomyces* sp and some fungi (Chapter 5).

Pectic Substances The pectic substances are polysaccharides of plant cell walls usually isolated as a mixture of which the major constituent is a polymer of methvlated galacturonic acids (235). Very little attention has been devoted to these materials as carbon sources although pectin has been reported to provide carbon for the growth of *Sclerotinia cinerea* (240) and of *Psalliotia bispora* (216). From the frequent occurrence in fungi of enzymes acting upon the pectic substances (Chapter 5) and from the employment of pectin as the carbon source in experiments on enzyme production it is probably safe to assume that they are utilized in nature by many fungi both parasitic and saprophytic.

5 ORGANIC ACIDS

Organic acids are generally poorer sources of carbon for fungi than are the carbohydrates. Apart from what might be called intrinsic utilizability however three factors interfere with the utilization of the organic acids in particular cases. First cells are often impermeable to organic acids at physiological pH levels. Impermeability is more often invoked than demonstrated but there is no doubt that it is a factor in the utilization of acids. Second the utilization of neutralized organic acids causes a rise of the culture pH which may interfere with growth. This phenomenon explains the repeated observation that growth with organic acids is better if nitrogen is supplied as an ammonium salt (115 165 207). Utilization of the ammonium cation tends to drive the pH down partially or wholly counteracting the alkalinizing effect of acid utilization. In addition it appears that the utilization of ammonium ion is more rapid if the medium is neutral or alkaline (134). It has been argued (24) that a more direct reaction exists between organic acids and ammonium ion but this appears unlikely and is not necessary in order to explain the observed phenomena.

A third and probably only occasionally important factor is the chelation of inorganic ions by certain of the organic acids e.g. citrate and tartrate. This may however either promote growth by keeping

essential ions in solution or limit growth by removing ions (183) which of these two opposite effects will occur is determined by the solubilities of the chelate complexes and the metallic salts at the culture pH chosen

The non fatty organic acids which are most often utilized by fungi are those e.g. citrate succinate and malate which are products of glucose metabolism. Most of the data are from studies on common saprophytes and it appears that utilization is highly specific. Thus *Penicillium janczewskii* grows well on succinate (23) succinate is a relatively poor carbon source for *P. javanicum* (121) and is not utilized at all by *P. digitatum* (65). Because of the likelihood that cells may be impermeable to acids especially hydroxy acids nutritional data have no necessary relevance to the acid metabolism of the cell.

Tartaric acid the utilization of which by a fungus is so important in the history of both crystallography and microbiology (148) is in general not a good carbon source for fungi although it is utilized as well as glucose by *Memoniella echinata* (150). The growth of some common fungi with tartrate has been surveyed by Stridman et al. (190) an adaptive tartrate decomposing system of *Aspergillus versicolor* has been described (10). The dextrorotatory isomer is used preferentially (70).

Oxalic acid is used by few microorganisms among which are *Proactinomyces* sp. and a strain of *Aspergillus oryzae* (137) *Psalliota bispora* (216) and a few bacteria (104). Possibly the insolubility of calcium and magnesium oxalates results in a removal of these ions from solution and a consequent limitation of growth in oxalic acid media.

Malonic acid is used although it is not a favorable carbon source by a few fungi and actinomycetes (23 92 99).

Inclusive studies of polyfunctional organic acids in nutrition have been reported by Camp (30) Tamura (201) and Steinberg (195).

The lower fatty acids—formic acetic propionic etc.—are as a rule poorly utilizable or even toxic. Propionic acid in fact is employed as a fungistatic agent. Some fungi however do make at least some growth with these acids particularly with acetate on a carbon basis acetate is equivalent to glucose for *Allomyces javanicus* (124). Qualitative data (175) indicate that *Leptomitus lacteus* can grow with any of the lower organic acids including some that are almost uniformly toxic to the higher fungi. Probably it will be found that the lower fungi utilize fatty acids better than the more extensively investigated higher forms. The utilization of acetate by *Penicillium chrysogenum* is strongly pH dependent (95).

Formic acid has been used as substrate in metabolic studies and apparently supports at least limited growth (6 15 38 76) but is never a good source of carbon. Its low nutritional value however does not exclude it or its derivatives from an active role in metabolism.

6 THE LIPIDS

Perhaps because of the technical difficulty in the use of water insoluble substrates the lipids have been little studied. However it seems clear that many fungi are able to grow on crude or purified natural fats and oils (62 89 162a). Similar observations incidental to metabolic studies indicate that some of but not all the higher fatty acids support the growth of *Mucor mucedo* (197) and *Penicillium glaucum* (212 213). The quantitative data of Tausson (205) are reproduced in Table 2 although the absence of controls somewhat reduces their sig-

Table 2 The Utilization of Fats by *Aspergillus flavus**

Experiment	Carbon Source	Dry Weight mg	Utilization per cent
1	Cocoa butter	508.0	30.8
	Coconut oil	459.2	—
2	Oleic acid	338.8	—
	Stearic acid	577.8	34.0
	Palmitic acid	510.2	25.0
	Tristearin	266.8	16.5
	Tripalmitin	258.4	41.4

* From Tausson (205) by permission of the *Biochemische Zeitschrift*

nificance it is clear that both fats and fatty acids are utilized and support growth in a medium containing no other source of carbon.

It has been observed that cholesterol and other steroids support the growth of *Proactinomyces* spp. (217 218) and that some fungi make visible growth with ergosterol as carbon source (177). Earlier work on crude steroids suffers from the probability of contamination with other carbon compounds.

Numerous fungi and actinomycetes grow on crude or purified hydrocarbons. The evidence reviewed by ZoBell (246 247) is largely observational only. *Proactinomyces opacus* grows on and oxidizes dodecane and tetradecane (231). Actinomycetes as a group appear to be more active than fungi in attack on paraffins, other hydrocarbons and steroids (60 98 219).

7 OTHER CARBON SOURCES

Pure Compounds The lower monohydric aliphatic alcohols—methanol ethanol etc—are not generally utilized although *Aspergillus niger* makes a limited growth with both methanol (6) and ethanol (201) The most extensive study of the alcohols is that of Stahl and Lessen (192) who found that *Aspergillus versicolor* grows poorly with ethanol and not at all with the other primary alcohols from methyl through undecyl (C_1 through C_{11}) Alcohols with 12 to 18 carbon atoms support some growth of *A. versicolor* and there appears to be an optimum chain length of 13 to 15

Amino acids of course provide carbon as well as nitrogen and many can serve as sole sources of carbon (139-203) In general if enough amino acid carbon is supplied for growth the accumulation of ammonia raises the pH to an unfavorably high level so that the amino acids alone cannot compare with the carbohydrates in the amount of growth supported

Enrichment culture methods have been used successfully in the isolation of an actinomycete able to use urethane as sole source of carbon (176) *Proactinomyces* spp make visible growth with several aromatic compounds e.g. phenol aniline and nitrobenzene as sole sources of carbon (133)

Natural Products Lignin is a complex amorphous polymer of high molecular weight the units of which are aromatic in nature but neither the chemical nor the physical structure is known precisely It is a plant product and is particularly abundant in wood

There is good evidence based on the analysis of plant materials before and after decomposition by pure cultures that some fungi are able to decompose lignin *in situ* i.e. in association with other substances of the plant Evidence on the fungi is limited however to the litter-decomposing and the wood destroying basidiomycetes The lignin of litter is decomposed actively by species of *Marasmius* *Collybia* *Mycena* *Clavaria* and other soil inhabiting Hymenomycetes (118) Most of the species studied use both the lignin and the cellulose of litter but a few appear to utilize only the lignin (75-118)

The wood rotting basidiomycetes have long been differentiated into two groups (63-64) the brown rot fungi e.g. *Merulius domesticus* attacking primarily the cellulose of wood and the white rot fungi e.g. *Polyporus annosus* which attack the lignin more rapidly Bavendamm (11) differentiated these two groups by their reaction on a tannic acid

medium the white rot fungi forming a colored zone (the oxidase reaction). Numerous studies on this reaction summarized by Gottlieb and Pelczar (72) indicate that the enzyme responsible is a polyphenol oxidase (Chapter 6) generally present in but not limited to the lignin utilizing fungi. The separation into brown and white rots although useful is not absolute—most white rot fungi use cellulose at one stage or another of the decomposition process (31-81). The brown rot fungi however do not attack lignin appreciably (35) except perhaps in the advanced stages of wood decomposition (169).

In considering the utilization of lignin as the sole source of carbon i.e. apart from other plant materials we are reminded of the dependence of nutritional studies on adequate knowledge of the chemistry of the substrate. Most of the early work reviewed by Waksman and Hutchings (226) is of doubtful validity since it appears that the lignin used was contaminated with carbohydrates and protein degradation products (21-72). A more recent method of preparation employing mild procedures is believed to yield an uncontaminated lignin (20). Studies on this material indicate that white rot fungi can utilize it (119-220) other wood destroying fungi may require a period of training i.e. a long incubation period in contact with the substrate before they can utilize it (49-71).

Although lignosulfonic acid disappears from media in which fungi are growing (1) it seems from other studies that lignosulfonate forms a complex with the mycelium and that the utilization therefore is only apparent (18).

Keratin the insoluble protein of wool hair and feathers has been implicated as a possible substrate for fungus growth since the observations of Ward (229) on *Onygena equina*. Jensen (97) reported growth and ammonia liberation by *Streptomyces* spp. in sand culture with keratin as the sole source of carbon and nitrogen. In a survey of keratin utilization (237) it was found that untreated wool is not attacked by common saprophytic fungi actively keratinolytic forms are all dermatophytes especially species of *Trichophyton* and *Microsporum*. The action of *Microsporum gypseum* on keratin and collagen is analyzed by Stahl et al. (191).

The success of keratin-containing materials as bait for chytrids (101-102) and the ability of *Polychytrium aggregatum* to make visible growth on keratin (2) suggest that these fungi which are not isolated by conventional procedures of soil and fabric microbiology are worth study as possible agents of keratin destruction in nature.

Purified rubber is attacked by both fungi and actinomycetes (217).

as in decomposition of pure hydrocarbons the actinomycetes appear to be the more active

Tannins are utilized by several common saprophytes (47 109 168) presumably after elaboration of an adaptive tannase (Chapter 6) The crude resins of *Guayule* are attacked by several fungi (3)

8 THE CONCENTRATION FACTOR IN CARBON NUTRITION

As the concentration of carbohydrate in a medium is increased the economic coefficient declines although more sugar may be utilized the dry weight does not increase proportionately Carbohydrate is evidently being converted to metabolic products—carbon dioxide organic acids soluble polysaccharides etc There also occurs at supraoptimal glucose concentrations a simple failure to utilize the sugar present (119) in which case the economic coefficient is not affected These two related phenomena are surely to be ascribed to the fact that other constituents of the medium especially nitrogen become limiting at the higher carbohydrate concentrations That is within limits growth is increased by higher carbohydrate provided that adequate nitrogen is supplied (27) The role of zinc and other metals may be more complicated than that of nitrogen supply (66) but the simple fact that more cellular synthesis will increase the demand for all nutrients should not be neglected in considering the effect of metals on growth and metabolism

For many fungi increase in carbohydrate beyond an optimum point results in an absolute as well as a relative decrease in growth The data of Figure 2 exemplify this type of response which is com




Figure 2 The effect of glucose concentration on the growth of *Psalliota bispora* Drawn from tabular data of Treschow (16)

mon in the fungi (160 166 227) Although most of such cases have not been analyzed the factor involved is almost certainly the formation of toxic metabolic products In *Entomophthora* spp the effect of high concentrations of glucose is clearly to increase the acidity of the medium (242) and the lowered pH resulting from organic acid accumulation is probably the most general cause of poor growth at high sugar concentrations

The carbohydrate concentration at which growth depressant effects become evident is of course affected by other constituents of the medium especially buffers and the nitrogen source but there are also specific effects Thus growth of *Ustilago violacea* falls off above 0.2 per cent glucose (180) that of *Penicillium javanicum* only above 21.3 per cent (227) It appears probable although there are very few data that in those fungi which grow well at high sugar concentrations (above 10 per cent) the factor which ultimately limits growth is osmotic pressure (Chapter I)

Probably because so much of the early work on fungi concerned the acid tolerant species of *Aspergillus* and *Penicillium* the media in common use for fungi are too high in available carbohydrate giving a distorted picture of growth and metabolism Ideally the level of carbohydrate to be used should be determined by experiment for each organism perhaps as a rough rule of thumb 2 per cent of available carbohydrate in a liquid medium may be suggested as an upper limit considering the amounts of nitrogen usually recommended in media Westerdijk (232) suggests that agar stock culture media for fungi contain not more than 3 per cent carbohydrate

9 ENZYME INDUCTION ADAPTIVE GROWTH AND THE UTILIZATION OF MIXED CARBON SOURCES

In this chapter we have had occasion to note that adaptive utilization of substrates occurs in the fungi Operationally adaptive growth is usually detected in an experiment in which a relatively large population of spores in a given medium grows only after prolonged incubation The organism is said to adapt itself to the carbon source It must be understood that in this type of experiment it is possible—and in some instances very probable—that the results can be explained by selection of a mutant pre-existing in the inoculum or arising during the long incubation period

In a related type of experiment it is found that a fungus produces much more of a given enzyme if cultivated on the substrate of the enzyme The literature on this phenomenon is very large and has

been reviewed several times in the past 10 years (1 37 110 245) Amylase inulase maltase sucrase lipase pectin polygalacturonase and tannase are among the more frequently investigated enzymes The results with pectin polygalacturonase are typical in that substances chemically related to the substrate may also increase enzyme formation (156) and in that the response to substrate occurs with some fungi and not with others (25) Here again it must be stressed that a genetic mechanism—selection of variant cells or of variant nuclei in a heterocaryon—is as probable in many instances as a physiological mechanism

Enzyme induction has been defined (42) as a relative increase in the rate of synthesis of a specific apoenzyme occurring in response to a specific chemical substance the inducer Earlier attention was focused on the specific substrate as inducer and the term adaptive enzyme was employed to denote an enzyme formed only or in increased amounts in the presence of substrate constitutive enzymes are defined as those which are formed in quantity in a favorable medium lacking substrate or inducer (103) The inducer is most often also the substrate but it may be a chemically related or an unrelated compound In a classical experiment (196) resting cells of *Escherichia coli* form the enzyme formic dehydrogenase if incubated in the presence of formate not otherwise

It has been shown by Landman (111) that the lactase (β galactosidase) of *Neurospora crassa* is induced under conditions which satisfy the requirements above particularly in that lactase is formed by resting cells exposed to an inducer (Figure 3) The data show also that sugars other than lactose act as inducers Resting cells respond somewhat differently from growing cells and the mechanisms may not be identical The time required for lactase induction in *N. crassa* is much longer than that required for induction of the same enzyme in yeast or in *Escherichia coli*

The foregoing demonstration is the only one to date in the fungi It seems reasonable however to postulate in agreement with Monod (132) that at least some of the reports of adaptive growth and of stimulation of enzyme formation by substrate which have been mentioned do in fact involve enzyme induction even though genetic change is not rigorously excluded

Earlier in this chapter particularly in reference to the growth of fungi with fructose and mannose it was suggested that results with mixed carbon sources may be explained in part by the induction of new enzymes Specifically the effect of a small amount of glucose in permitting rapid growth on fructose or mannose sugars which by




Figure 3 The formation of β galactosidase (lactase) by resting mycelium of *Neurospora crassa*. Arrows indicate time of exposure to the inducing sugar. Redrawn from O. E. Landman *Archives of Biochemistry and Biophysics* Vol 52 p 93-100 (1954) by permission of Academic Press Inc.

themselves support growth only after a long lag may be explained by postulating that the glucose allows sufficient growth so that an enzyme essential to fructose or mannose utilization can be synthesized. Naturally some energy must be available for the enzyme synthesis for resting cells endogenous reserves may be adequate but with a spore inoculum an exogenous energy source may be necessary.

Extensive data on the utilization of mixed carbon sources have been reported by Lilly and Barnett (116). Several fungi e.g. *Sordaria fimicola* and *Thielaviopsis basicola* behave in accordance with the foregoing postulate—they grow on a disaccharide melibiose for example only if a utilizable monosaccharide is available. However results with *Gliomastix convoluta* are not entirely explicable on this basis unless we postulate without proof that the sugar which supplies energy for enzyme synthesis must be present in relatively high concentration. These experiments are complicated by the inclusion of asparagine which supplies some carbon in the basal medium.

The results of Steinberg (194) on the utilization of mixed carbon sources by *Aspergillus niger* are not satisfactorily explained by the hypothesis of enzyme induction. Here a mixture of two unutilizable compounds supports good growth. It is clear from these results and from those mentioned in the preceding paragraph that the interaction of different carbon sources requires further study.

In relation to utilization of mixed carbon sources it should be noted that preferential utilization of one compound is common e.g. the utilization of acetate before glucose (151) and the utilization of soluble sugars before more complex carbon compounds (62 109 188)

Fungi in nature live as a rule in a dilute solution with more than one carbon source available or potentially available. These conditions do not very closely resemble a single substrate pure culture. Lignin and cellulose for example may be much more rapidly decomposed by the action of induced enzymes than they are when given to an organism in culture as the sole source of carbon. The ecological importance of the ability of fungi to synthesize new enzymes—if this ability proves to be general—justifies much more study of the problem. It is to be hoped however that such studies will be guided more than they have been in the past by the wealth of experience accumulated in the investigation of induced enzymes in other microorganisms.

10 EFFICIENCY OF GROWTH

Several indices of the over all efficiency of growth in fungi have been proposed. Of these the economic coefficient has been most frequently determined. It may best be represented as

$$\frac{\text{Mycelial dry weight gm}}{\text{Carbohydrate consumed gm}} \times 100$$

Some authors have used the ratio or its reciprocal without multiplication by 100.

As an approximate index of the efficiency of carbon utilization the economic coefficient is satisfactory. More elaborate indices which take account of energy quantities more directly have been devised (202 206 208 210).

The economic coefficient can be seen by inspection to be maximal when respiratory carbon dioxide and soluble metabolic products are minimal in quantity. Since both of these are affected by cultural conditions no one value or even range of values can be set down as characteristic of the fungi as a whole. Fungi grown on dilute media for no longer a time than is necessary to utilize the carbohydrate usually have an economic coefficient in the range 20 to 35 a few higher values have been reported (119 161 205 209). Very low values (77 228 230) reflect probably the production of significant amounts of soluble carbon compounds in high carbohydrate media.

The economic coefficient although unaffected by temperature (209)

is strongly influenced by substrate (205) metal deficiencies (29 158 167 182) and nitrogen nutrition (157 211) The influence of age is variable (28 155) As one would expect increasing the glucose supplied decreases the economic coefficient (119 159 195)

All of the factors just listed as influencing the economic coefficient almost certainly act by changing the amount of soluble metabolic products formed it is a possibility not so far explored that the loss of carbon as CO₂ is affected but this seems doubtful A low economic coefficient should suggest in the first instance a search for soluble carbon compounds in the medium

11 CONCLUSION

The most important single facet of carbon nutrition is the high degree of specificity differences between species and strains are the rule in the fungi Nevertheless a search for some general principle has been and should be pursued One such is to relate carbon nutrition to phylogeny

It has often been pointed out e.g. by Garrett (69) that the major groups of fungi differ with respect to cellulose and lignin utilization The Zygomycetes do not use either of these materials in the ascomycetes and the Fungi Imperfecti cellulose utilization is common the basidiomycetes contain most of the known lignin utilizing fungi in addition to many species able to grow on cellulose This progression toward utilization of more complex materials would be more impressive if the capacity to break down cellulose were not widespread in the Chytridiales and perhaps in the Oomycetes

The phycomycetes usually conceded to be the most primitive major group of fungi have certain nutritional and metabolic characteristics which taken together separate them rather clearly from the higher fungi

1 The fungi which produce large amounts of lactic acid from carbohydrate are all phycomycetes (Chapter 6)

2 The fungi able to grow at low oxygen tensions are usually phycomycetes (Chapter 1) This ability may of course be related to the ability to form lactic acid

3 Requirements for organic sulfur among fungi occurring in nature are known only in the phycomycetes (Chapter 9)

4 The ability to use the lower fatty acids appears to be more common in the phycomycetes than in other groups (p. 81) although the data are still rather limited

It may be that the appropriate theoretical framework for carbon nutrition is more the ecology than the phylogeny of fungi. The characteristics of the phycomycetes outlined above can be understood as related to the aquatic habit of most of the forms so far studied just as easily as they can be interpreted in terms of taxonomy or phylogeny. The carbon nutrition of the mycorrhizal basidiomycetes is another example. The evidence is incomplete and not entirely satisfactory but it appears that this group is restricted in its carbon nutrition to simple sugars starch and the pectic substances (74 91 130 131 170). These fungi stand therefore in contrast to the litter-decomposing and wood rotting basidiomycetes which use cellulose lignin or both. At least one mycorrhizal basidiomycete however does form an adaptive cellulase (142).

The order Mucorales is cited often as a group of sugar fungi unable to utilize cellulose (26). This characterization appears to be sound although relatively few of the Mucorales have been adequately investigated. Ecologically however the sugar fungi include a number of the Fungi Imperfecti especially species of *Aspergillus* and *Penicillium* found in soils high in fresh organic matter and in carbohydrate-containing materials. The line between sugar fungi and cellulose utilizing fungi cuts therefore across taxonomic boundaries since different species of *Aspergillus* for example are found in one or the other group.

In Table 1 are collected some data on the carbon nutrition of different fungi although it cannot be said that the organisms chosen are in any way typical. The organisms of wide substrate range include a large group of soil saprophytes and plant parasites—*Aspergillus oryzae* *Brevilegnia gracilis* *Fusarium oxysporum* f. *nicotianae* *Memnoniella echinata* *Penicillium digitatum* *Phytophthora cactorum* *Psalliota bispora* *Stachybotrys atra* and *Streptomyces coelicolor*. In comparison the aquatic phycomycetes in the list appear more fastidious the contrast between the soil inhabiting *Brevilegnia gracilis* and the closely related aquatic *Saprolegnia delica* is particularly instructive. On the other hand the plant parasite *Chalara quercina* has a much narrower nutritional range than the aquatic *Pythiogeton* sp. The most restricted of all in the group are *Entomophthora coronata* parasitic on insects and the aquatic *Polychytrium aggregatum*.

The fungi parasitic on animals might be expected by analogy to bacterial parasites to be nutritionally specialized. The available data indicate that these fungi are not in fact particularly fastidious with respect to carbon sources (140 200).

One incidental problem may be mentioned here. Throughout this

chapter it has been assumed that carbon sources are utilized by pathways involving hexoses or hexose phosphates as intermediates. However the data of Table I afford several instances in which glucose is inferior to some other carbon compound xylose gluconic acid disaccharides or polysaccharides. None of these cases has been studied. The superiority of xylose or gluconate could mean that the phosphogluconate oxidation pathway (Chapter 7) is more active than others. Or a compound may be superior to glucose for accidental reasons—pH relations accumulation of intermediates contamination with metals or growth factors etc. This second explanation has been suggested earlier for the superiority of particular disaccharides and polysaccharides. It must be admitted however that the question is still open.

Looking at the whole field of carbon nutrition of fungi one is struck with the gaps in our knowledge. Certain groups—the Mucorales and the Peronosporales are examples—have been more or less neglected. Comparative studies of species within a genus particularly species of different habitat might prove most valuable. The biochemical basis of nutritional capabilities is largely a matter of inference at present failure to utilize fructose or mannitol for example probably has a simple enzymatic basis. As just discussed above induced enzyme synthesis in the fungi is still largely a matter for speculation. Some of the data reported in this chapter and more which have been omitted are difficult to accept and important points of this nature should be reinvestigated. Finally the understanding of nutrition in nature with numerous substrates present in low concentration is a longrange goal progress toward which will require radically new techniques of investigation.

BIBLIOGRAPHY

- 1 Adams G. A. and G. A. Ledingham 1949 *Can. J. Research C* 90 101-104
- 2 Ajello L. 1918 *Am. J. Botany* 30 130-140
- 3 Allen P. J., J. Naghski and S. R. Hoover 1911 *J. Bacteriol.* 47 509-568
- 4 Ammann A. 1941 *Phytopathol. Z.* 18 416-446
- 5 Aschan K. and B. Norkrans 1953 *Physiol. Plantarum* 6 61-585
- 6 Baba S. 1911 *Bull. Agr. Chem. Soc. Japan* 17 109-110
- 7 Ball E. 1953 *Bull. Torrey Bot. Club* 80 409-411
- 8 Barnard D. and F. Challenger 1919 *J. Chem. Soc. (London)* 1919 110-117
- 9 Barner H. D. and E. C. Cantino 1952 *Am. J. Botany* 39 746-751
- 10 Barton R. R. 1953 *Mycologia* 45 533-547
- 11 Bavendamm W. 1927 *Ber. deut. botan. Ges.* 4, 357-367
- 12 Beckman C. H., J. E. Kunz and A. J. Riker 1953 *Phytopathology* 43 441-447

- 13 Benedict R G L A Lindensfelser F H Scodola and D H Trauffer 1951
J Bacteriol 62 487-497
- 14 Benedict R G T G Prudham L A Lindensfelser and R W Jackson 1955
Appl Microbiol 3 1-6
- 15 Bernhauer A and F Stanina 1933 *Biochem Z* 264 109-112
- 16 Bhargava K S 1943 *J Indian Botany Soc* 22 85-99
- 17 Bhargava K S 1945 *Lloydia* 8 60-68
- 18 Blank L M and P J Talley 1941 *Am J Botany* 28 564-69
- 19 Blumer S 1937 *Arch Mikrobiol* 8 458-478
- 20 Brauns F E 1939 *J Am Chem Soc* 61 2190-2127
- 1 Brauns F E 1952 *The Chemistry of Lignin* New York Academic Press
pp 808
- 22 Bretzlaff C W Jr 1954 *Am J Botany* 41 58-67
- 23 Brian P W P J Curtis and H G Hemming 1946 *Trans Brit Mycol Soc* 29 173-187
- 24 Brian P W P J Curtis and H G Hemming 1947 *Proc Roy Soc (London)* B 135 106-132
- 25 Brown W and R K S Wood 1953 *Symposium Soc Gen Microbiol* 3 506-539
- 26 Burges A 1939 *Botera Sér trimestri al* 8 64-81
- 27 Burkholder P R and I McVeigh 1940 *Am J Botany* 27 634-640
- 28 Butkewitsch W 1952 *Biochem Z* 151 338-350
- 29 Butkewitsch W and W G Orlov 1922 *Biochem Z* 135 556-56
- 30 Camp A F 1953 *Ann Missouri Botanical Garden* 10 913-93
- 31 Campbell W G 1932 *Biochim J (London)* 26 1829-1838
- 32 Cantino E C 1949a *Am J Botany* 36 95-111
- 33 Cantino E C 1949b *Am J Botany* 36 747-756
- 34 Cantino E C 1950 *Quart Rev Biol* 25 289-277
- 35 Cartwright K St G and W P K Findlay 1946 *Decay of Timber and its Prevention* London H M Stationery Office pp 294
- 36 Cheldehn V H and T E King 1953 *Ann Rev Microbiol* 7 113-140
- 37 Christensen J J and J M Daly 1951 *Ann Rev Microbiol* 5 57-70
- 38 Chrzastacz T and M Zakomorny 1952 *Biochem Z* 279 64-75
- 39 Cochrane V W 1957 *J Bacteriol* 63 459-471
- 40 Cochrane V W and J E Conn 1947 *J Bacteriol* 54 15-218
- 41 Cochrane V W H D Peck Jr and A Harrison 1953 *J Bacteriol* 66 1-23
- 42 Cohn M J Monod M R Pollock S Spiegelman and R Y Starnes 1953
Nature (London) 170 1096
- 43 Conti H J 1921 *N Y State Agr Exp Sta (Geneva N Y) Bull* 83 1-76
- 44 Corum C J 194 *Botan Gaz* 103 749-750
- 45 Crasemann J M 1954 *Am J Botany* 41 502-510
- 46 Czapek F 1903 *Beitr Chem Physiol Path* 3 47-66
- 47 Davis D P E Waggoner and A E Diamond 1953 *Natu r (London)* 172 99
- 48 Day W C S Gottlieb and M J Pelczar Jr 1953 *Appl Microbiol* 1 8-81
- 49 Day W C M J Pelczar J and S Gottlieb 1949 *Arch Biochem Biophys* 23 360-369
- 50 DeLey J and J Vandamme 1957 *J Gen Microbiol* 10 167-171
- 51 Dietzel E H Behrenbruch and M Eucken 1950 *Arch Mikrobiol* 15 159-184
- 52 Diamond A E and G L Pelzer 1954 *Am J Botany* 41 46-50
- 53 Doudoroff M 1951 In W D McElroy and B Glass (eds) *Phosphorus Metabolism* Vol 1 p 4-48 Baltimore Johns Hopkins Press

- 51 Dox A W and R E Neidig 1912 *Biochem* 7 46 397-409
- 55 Dubos R J 1928a *Ecology* 9 12-21
- 56 Dubos R J 1928b *J Bacteriol* 15 223-234
- 57 Ehrensward G 1955 *Exp Cell Research Suppl* 3 102-109
- 58 Emerson R and E C Cantino 1948 *Am J Botany* 35 157-171
- 59 Englis D T and D J Hanahan 1941 *J Am Chem Soc* 63 51-54
- 60 Erikson D 1941 *J Bacteriol* 41 277-300
- 61 Estienne V E Castagne and E Bertrand 1917 *Bull soc chim biol* 99 620-627
- 62 Eyre J C 1932 *Ann Appl Biol* 19 351-369
- 63 Falck R 1926 *Ber deut botan Ges* 41 652-661
- 64 Falck R and W Haag 1927 *Ber deut chem Ges* 60 225-232
- 65 Fergus C L 1952 *Mycologia* 44 183-189
- 66 Foster J W 1949 *Chemical Activities of Fungi* New York Academic Press pp 648
- 67 Fruton J S and S Simmonds 1953 *General Biochemistry* New York John Wiley and Sons pp 940
- 68 Fuller W H and A G Norman 1945 *J Bacteriol* 50 667-671
- 69 Garrett S D 1951 *New Phytologist* 50 149-165
- 70 Gieski F 1937 *Bull intern acad polon sci Classe sci math nat Ser B* 1 1937 89-105
- 71 Gottlieb S W C Day and M J Pelczar Jr 1950 *Phytopathology* 40 926-935
- 72 Gottlieb S and M J Pelczar Jr 1951 *Bacteriol Rev* 15 55-16
- 73 Greathouse G A D E Klemme and H D Barker 1912 *Ind Eng Chem Anal Ed* 14 611-620
- 74 Harley J L 1950 *Ann Rev Microbiol* 6 367-386
- 75 Harris G C M 1945 *Ann Appl Biol* 32 38-39
- 76 Harrold C E and M Flung 1952 *J Biol Chem* 194 399-406
- 77 Harter L L and J L Weimer 1951 *J Agr Research* 21 211-226
- 78 Haskins R H and W H Weston Jr 1950 *Am J Botany* 37 739-750
- 79 Hawker L F and S D Chaudhuri 1946 *Ann Botany* 10 185-191
- 80 Hawkins L A 1915 *Am J Botany* 2 375-388
- 81 Hawley L F L C Fleck and C A Richards 1928 *Ind Eng Chem* 20 504-507
- 82 Herrick J A 1910 *Ohio J Sci* 40 123-129
- 83 Hestrin S 1948 *Wallerstein Lab Commun* 11 193-207
- 84 Hestrin S 1949 *Wallerstein Lab Commun* 12 45-54
- 85 Heuser E B F Shema W Shockley J W Appling and J F McCoy 1910 *Arch Biochem* 21 343-350
- 86 Hill E G and A R Patton 1947 *Science* 105 481-489
- 87 Hofmann E 1931 *Biochem Z* 243 423-428
- 88 Holden M 1950 *Biochem J (London)* 47 426-431
- 89 Horowitz Wassowa L M and M J Linschitz 1935 *Zentr Bakteriell Parasitenk Abt II* 92 424-435
- 90 Horr W H 1936 *Plant Physiol* 11 81-93
- 91 How J E 1940 *Ann Botany* 4 155-150
- 92 Hubbard C V and H H Thornberry 1950 *Trans Illinois State Acad Sci* 43 61-74
- 93 Hungate R E 1916 *J Bacteriol* 51 51-55
- 94 Ingraham J L and R Emerson 1954 *Am J Botany* 41 146-152
- 95 Jarvis F G and M J Johnson 1947 *J Am Chem Soc* 69 3010-3017
- 96 Jeffreys E G P W Brian H C Hemming and D Lowe 1953 *J Gen Microbiol* 9 514-541
- 97 Jensen H L 1950 *J Agr Sci* 20 390-398

- 99 Jensen H L 1931 *Proc Linnean Soc N S Wales* 56 315-310
- 99 Jermy M A 1953 *Australian J Biol Sci* 6 48-69
- 100 Johnson G T and A C Jones 1911 *Mycologia* 33 421-433
- 101 Karling J S 1946 *Am J Botany* 33 751-754
- 102 Karling J S 1947 *Am J Botany* 34 27-32
- 103 Karlström H 1938 *Ergeb Enzymforsch* 7 350-316
- 104 Khambata S R and J V Bhat 1953 *J Bacteriol* 66 505-507
- 105 Kitz D A and W H Peterson 1953 *J Biol Chem* 203 661-668
- 106 Klein H P and M Doudoroff 1950 *J Bacteriol* 9 739-750
- 107 Klemme D E 1947 *J Bacteriol* 43 171-180
- 108 Klemme D E G A Greathouse K Bollenbacher and S Pope 1945 *U S Dept Agr Circ No 131* 1-11
- 109 Knudson L 1913 *J Biol Chem* 14 159-181 185-202
- 110 Kynl R 1914 *Jahrb wiss Botan* 53 465-501
- 111 Landman O E 1954 *Arch Biochem Biophys* 52 93-109
- 112 Landman O E and D M Bonner 1957 *Arch Biochem Biophys* 41 253-265
- 113 Leben C and C W Keitt 1948 *Am J Botany* 35 331-343
- 114 Lein J D C Appleby and P S Lein 1951 *Arch Biochem Biophys* 31 72-80
- 115 Leonian L H and V G Lilly 1940 *Am J Botany* 27 19-26
- 116 Lilly V G and H L Barnett 1953 *West Va Univ Agr Exp Sta Bull* 367T 5-58
- 116a Lilly V G and H L Barnett 1956 *Am J Botany* 43 69-714
- 117 Lindeberg G 1944 *Symbolae Botan Upsalenses* 8(2) 1-183
- 118 Lindberg G 1946 *Arkiv Botan* 33A (10) 1-16
- 119 Lockwood L B and G E N Nelson 1946a *Arch Biochem* 10 365-374
- 120 Lockwood L B and G E N Nelson 1946b *J Bacteriol* 52 581-586
- 121 Lockwood L B G E Ward O E May H T Herrick, and H T O'Neill 1951 *Zentr Bakteriol Parasitenk Abt II* 90 411-455
- 122 Lwoff A H Ionescu and A Gutmann 1949 *Compt rend (Paris)* 278 34-344
- 123 McBeth I G and F M Scales 1913 *U S Dept Agr Bur Plant Ind Bull* 266 1-57
- 124 Machlis L 1953 *Am J Botany* 40 460-464
- 125 McKee W E 1950 *Science* 123 509
- 126 Marsh P B K Bollenbacher M L Butler and K B Raper 1949 *Textile Research J* 19 465-481
- 127 Martin J P 1945 *J Bacteriol* 50 349-360
- 128 Mehrotra B S 1949 *J Indian Botan Soc* 8 108-124
- 129 Mehrotra B S 1951 *Lloydia* 14 122-128
- 130 Meln E 1953 *Untersuchungen über die Bedeutung der Baum Mykorrhiza* Jera Gustav Fischer pp 152
- 131 Meln E 1948 *Trans Brit Mycol Soc* 50 97-99
- 132 Monod J 1947 *Growth* 11 225-283
- 133 Moore W F 1919 *J Gen Microbiol* 3 143-147
- 134 Morton A G and A Macmillan 1954 *J Exp Botany* 5 232-252
- 135 Moyer A J and R D Coghill 1946 *J Bacteriol* 51 57-78
- 136 Mudge C S 1917 *J Bacteriol* 2 403-415
- 137 Müller H 1950 *Arch Mikrobiol* 15 137-148
- 138 Newth F H 1951 *Advances in Carbohydrate Chem* 6 83-106
- 139 Nickerson W J and R R Mohan 1953 *Intern Congr Microbiol. 6th Congr Rome* 5 137-146
- 140 Nickerson W J and J W Williams 1947 In W J Nickerson (ed) *Biology of Pathogenic Fungi* p 130-156 Waltham Mass The Chronica Botanica Co

- 141 Nord F F and E. W Engel 1938 *Biochem Z* 296 153-160
- 142 Norkrans B 1930 *Symbolae Botan Upsalienses* 11 (1) 5-126
- 143 Norman A G 1931 *Ann Appl Biol* 18 241-259
- 144 Obaton F 1932 *Compt rend (Paris)* 194 302-304
- 145 O'Connor R C 1940 *Biochem J (London)* 34 1008-1011
- 146 Oddoux L 1952 *Compt rend (Paris)* 284 654-657
- 147 Otto H 1916 *Beitr Allgem Botan* 1 190-260
- 148 Pasteur I 1860 *Compt rend (Paris)* 51 298-299
- 149 Pelczar M J Jr S Gottlieb and W C Day 1950 *Arch Biochem* 25 449-451
- 150 Perlman D 1948a *Am J Botany* 35 36-41
- 151 Perlman D 1948b *Am J Botany* 35 360-363
- 152 Perlman D 1950 *Bull Torrey Botan Club* 77 103-109
- 153 Perlman D D A Kita and W H Peterson 1946 *Arch Biochem* 11 173-129
- 154 Peterson W H 1946-1947 *Harvey Lectures Ser* 42 216-309
- 155 Peterson W H E B Fred and E G Schmidt 1922 *J Biol Chem* 54 19-34
- 156 Phaff H J 1947 *Arch Biochem* 13 67-81
- 157 Pinck L A and F E Allison 1914 *Soil Sci* 57 155-161
- 158 Porges N 1932 *Potan Gaz* 94 197-205
- 159 Prill E A P R Wenck and W H Peterson 1950 *Biochem J (London)* 29 21-33
- 160 Quantz L 1943 *Jahrb wiss Botan* 91 120-163
- 161 Raaf H 1941 *Arch Microbiol* 12 131-189
- 162 Reese E T 1947 *J Bacteriol* 53 389-400
- 162a Reese E T H Cravetz and G R Mandels 1955 *Farlowia* 4 409-421
- 163 Reese E T and M H Downing 1951 *Mycologia* 43 16-28
- 164 Reese E T and H S Levinson 1952 *Physiol Plantarum* 5 345-366
- 165 Reischer H 1951 *Mycologia* 43 319-328
- 166 Reitsma J 1932 *Phytopathol Z* 4 461-462
- 167 Rippel A and G Behr 1934 *Arch Mikrobiol* 5 561-577
- 168 Rippel A and J Keseling 1930 *Arch Mikrobiol* 1 60-77
- 169 Pobak H 1942 *Med Vestl Forst Forskstation* 7(3) 1-248
- 170 Romell L G 1938 *Scensk Botan Tidskr* 32 89-99
- 171 Rothstein A 1954 *Symposia Soc Exp Biol* 8 165-201
- 172 Ryan F J G W Beadle and E L Tatum 1943 *Am J Botany* 30 784-799
- 173 Saksena R K and S K Bose 1944 *J Indian Botan Soc* 23 108-112
- 174 Satina S and A F Blakeslee 1928 *Proc Natl Acad Sci U S A* 14 279-283
- 175 Schade A L 1940 *Am J Botany* 27 376-384
- 176 Schatz A H D Isenberg A A Angnst and V Schatz 1951 *J Bacteriol* 63 1-4
- 177 Schatz A K Savard and I J Pintner 1949 *J Bacteriol* 79 117-12
- 178 Scheffer T C 1936 *U S Dept Agr Tech Bull No* 52 1-45
- 179 Schmidt E G W H Peterson and E B Fred 1923 *Soil Sci* 15 479-489
- 180 Schopfer W H and S Blumer 1938 *Arch Mikrobiol* 9 305-367
- 181 Schopfer W H and M Guisliand 1945 *Z Vita ninforsch* 16 181-206
- 182 Schulz G 1937 *Panta* 2, 196-218
- 183 Schutter J and P W Wilson 1950 *J Gen Microbiol* 1 446-451
- 184 Shirakawa H S 1950 *Am J Botany* 42 379-381
- 185 Sistrom D E and I Machlis 1955 *J Bacteriol* 70 0-
- 186 Su R C H 1951 *Microbial Decomposition of Cellulose* New York Reinhold Publ Corp pp 331

18 Siu R. C. H. R. T. Darby P. R. Burkholder and E. S. Barghoorn 1949 *Textile Research J* 19 481-488

188 Siu R. C. H. and J. W. Sinden 1951 *Am J Botany* 38 281-290

189 Skinner C. E. and F. Dravis 1937 *Ecology* 18 391-394

190 Stadlman T. C. R. H. Vaughn and G. L. Marsh 1915 *J Bacteriol* 50 601-700

191 Stahl W. H. B. McQue G. R. Mandels and R. C. H. Siu 1950 *Textile Research J* 20 510-579

197 Stahl W. H. and H. Pessen 1953 *Appl Microbiol* 1 30-35

193 Stamer R. Y. 1942 *Bacteriol Revs* 6 143-196

194 Steinberg R. A. 1939 *J Agr Research* 59 749-764

19 Steinberg R. A. 1942 *J Agr Research* 64 615-633

196 Stephenson M. and L. H. Suckland 1933 *Biochem J (London)* 27 153-155

197 Stern A. M. Z. J. Ordal and H. O. Halvorson 1954 *J Bacteriol* 68 24-27

198 de Stevens G. and F. F. Nord 1952 *J Am Chem Soc* 74 336-337

199 Stevens N. E. and R. A. Chapman 1942 *Phytopathology* 32 181

200 Stockdale P. 1953 *Biol Revs Cambidge Phil Soc* 28 81-101

201 Tamiya H. 1932 *Acta Phytochim (Japan)* 6 1-129

202 Tamiya H. 1942 *Advances in Enzymol* 2 183-238

203 Tamiya H. and S. Usami 1940 *Acta Phytochim (Japan)* 11 261-293

204 Tatum E. L. R. W. Barratt and V. M. Cutter Jr 1949 *Science* 109 509-511

205 Taussan W. O. 1958 *Biochem Z* 193 85-93

206 Taussan W. O. 1955 *Mikrobiologiya* 4 166-175

207 Tempel E. 1931 *Arch Mikrobiol* 2 40-71

208 Terroine E. F. and R. Bonnet 1930 *Bull soc chim biol* 12 10-19

209 Terroine E. F. and R. Wurmser 1922a *Bull soc chim biol* 4 519-567

210 Terroine E. F. and R. Wurmser 1922b *Compt rend (Paris)* 174 1435-1437

211 Terroine E. F. and R. Wurmser 1922c *Compt rend (Paris)* 175 228-230

212 Thaler H. and W. Eisenlohr 1941 *Biochem Z* 308 88-102

213 Thale H. and G. Geist 1939 *Biochem Z* 302 121-136

214 Thom C. H. Humfeld and H. P. Holmes 1933 *Am Dyestuff Repts* 23 581-586

215 Thorne C. B. and W. H. Peterson 1918 *J Biol Chem* 176 413-419

216 Treschow C. 1944 *Dansk botan Ark v* 11(6) 1-180

217 Turfitt G. E. 1944 *J Bacteriol* 47 487-493

218 Turfitt G. E. 1947 *J Bacteriol* 54 557-567

219 Umbreit W. W. 1939 *J Bacteriol* 38 73-89

220 Van Vleet W. F. 1954 *Biochim et Biophys Acta* 15 211-16

221 Waksman S. A. 1919 *Soil Sci* 8 71-215

222 Waksman S. A. 1931 *Arch Mikrobiol* 135-151

223 Waksman S. A. 1937 *Principles of Soil Microbiology* 2nd ed Baltimore Williams and Wilkins Co pp 894

224 Waksman S. A. 1940 *Botan Rev* 6 637-663

225 Waksman S. A. and R. A. Diehm 1931 *Soil Sci* 37 73-95 97-117

226 Waksman S. A. and I. J. Hutchings 1936 *Soil Sci* 47 119-130

227 Ward G. E. L. B. Lockwood O. E. May and H. T. Herrick 1935 *Ind Eng Chem* 27 318-322

228 Ward G. E. L. B. Lockwood O. E. May and H. T. Herrick 1936 *J Am Chem Soc* 58 1786-1788

229 Ward H. M. 1899 *Phil Trans Roy Soc London S B* 191 763-791

230 Wasmuth G. M. 195 *Biochem Z* 28 226-234

231 Webley D. M. and P. C. DeKock 1950 *Biochem J (London)* 51 371-375

- 141 Nord F F and E W Engel 1938 *Biochem J* 296 153-170
- 142 Norkrans B 1930 *Symbolae Botan Upsalienses* 11 (1) 5-106
- 143 Norman A G 1931 *Ann Appl Biol* 18 241-259
- 144 Obaton F 1932 *Compt rend (Paris)* 191 302-301
- 145 O Connor R C 1910 *Biochem J (London)* 34 1008-1011
- 146 Oddoux L 1932 *Compt rend (Paris)* 231 634-637
- 147 Otto H 1916 *Beitr Allgem Botan* 1 190-200
- 148 Pasteur L 1860 *Compt rend (Paris)* 51 298-299
- 149 Pelczar M J Jr S Gottlieb and W C Day 1930 *Arch Biochem* 23 449-451
- 150 Perlman D 1919a *Am J Botany* 53 36-41
- 151 Perlman D 1918b *Am J Botany* 35 360-363
- 152 Perlman D 1930 *Bull Torrey Botan Club* 77 103-109
- 153 Perlman D D A Kita and W J Peterson 1910 *Arch Biochem* 11 123-129
- 154 Peterson W H 1916-1917 *Harvey Lectures Ser* 42 276-300
- 155 Peterson W H L B Fred and E G Schmidt 1922 *J Biol Chem* 44 19-34
- 156 Phaff H J 1917 *Arch Biochem* 13 67-81
- 157 Pinck L A and F E Allison 1914 *Soil Sci* 5 155-161
- 158 Porces N 1930 *Botan Gaz* 91 197-203
- 159 Prill E A P R Wenck and W H Peterson 1933 *Biochem J (London)* 29 21-33
- 160 Quantz L 1913 *Jahrb wiss Botan* 91 120-164
- 161 Raaf H 1911 *Arch Microbiol* 12 131-182
- 162 Reese F T 1917 *J Bacteriol* 53 389-400
- 162a Reese E T H Cravetz and C R Mandels 1935 *Farlowia* 4 409-411
- 163 Reese E T and M H Downing 1931 *Mycologia* 43 16-28
- 164 Reese E T and H S Levinson 1932 *Physiol Plantarum* 5 345-360
- 165 Reischer H 1931 *Mycologia* 43 319-328
- 166 Reitsma J 1932 *Phytopathol* 24 461-502
- 167 Rippel A and G Behr 1934 *Arch Mikrobiol* 5 561-577
- 168 Rippel A and J Kesseling 1930 *Arch Mikrobiol* 1 60-77
- 169 Rohak H 1940 *Med Vestl Forst Forskstation* 7(3) 1-248
- 170 Romell L C 1938 *Svensk Botan Tidskr* 32 89-99
- 171 Rothstein A 1934 *Symposia Soc Exp Biol* 8 163-201
- 172 Ryan F J C W Beadle and E L Tatum 1913 *Am J Botany* 30 781-799
- 173 Saksena R K and S K Bose 1944 *J Indian Botan Soc* 23 103-110
- 174 Satina S and A F Blakeslee 1928 *Proc Natl Acad Sci U S A* 14 229-233
- 175 Schade A L 1910 *Am J Botany* 27 376-384
- 176 Schatz A H D Isenberg A A Angrist and V Schatz 1934 *J Bacteriol* 68 1-4
- 177 Schatz A K Savard and J J Pintner 1919 *J Bacteriol* 18 117-123
- 178 Scheffer T C 1936 *U S Dept Agr Tech Bull No* 527 1-4
- 179 Schmidt E G W H Peterson and F B Fred 1923 *Soil Sci* 15 479-493
- 180 Schopfer W H and S Blumer 1938 *Arch Mikrobiol* 9 303-367
- 181 Schopfer W H and M Cuilloud 1915 *Z Naturforsch* 16 181-200
- 182 Schulz G 1937 *Planta* 27 196-218
- 183 Schutter J and P W Wilson 1935 *J Gen Microbiol* 10 446-454
- 184 Shurakawa H S 1935 *Am J Botany* 40 379-384
- 185 Siström D E and L Machlis 1935 *J Bacteriol* 70 0-3
- 186 Siu R G H 1951 *Microbial Decomposition of Cellulose* New York Reinhold Publ Corp pp 531

4. Carbon Metabolism

Part I

Introduction

The sheer bulk of the reported investigative work on carbon metabolism dictates some subdivision of the topic for the sake of convenience. In this and the two succeeding chapters we are concerned with the principal transformations of carbon compounds brought about by fungi, excluding those primarily important in the metabolism of nitrogenous compounds (Chapter 8) and a few which are better considered in conjunction with respiration (Chapter 7).

Broadly considered, the physiological problems of carbon metabolism are two: the conditions under which a compound is utilized and the conditions under which a compound is manufactured by the cell. These lead to two corresponding biochemical problems: the pathway of breakdown and the pathway of biosynthesis. The natural occurrence of compounds and processes is also of interest. For any class of compounds the information is usually incomplete; thus the physiology of acid formation is better known than are the biosynthetic pathways involved, and very little solid information on the breakdown of the acids is available. Many interesting compounds are known so far largely as curiosities, since neither their physiological significance nor their biosynthesis has been explored. It seems warranted, however, to include at least some examples of these in a discussion of fungus physiology.

Because so much work has been done on the metabolites which accumulate in large amounts, it is salutary to recall that probably most fungi in culture form cell material and carbon dioxide exclusively. Thus of 40 basidiomycetes investigated by Whitaker (21)

- 232 Westerdijk J 1917 *Antonie van Leeuwenhoek J Microbiol Serol* 12 223-231
- 233 Whiffen A J 1911 *J Flisha Mitchell Sci Soc* 57 321-330
- 234 Whiffen A J 1915 *J Flisha Mitchell Sci Soc* 61 114-123
- 235 Whistler R L and C L Smart 1955 *Polysaccharide Chemistry* New York Academic Press pp 493
- 236 White W L R T Darby G M Stechert and K. Sanderson 1918 *Mycologia* 10 54-81
- 237 White W L G R Mandels and R G H Siu 1950 *Mycologia* 42 199-223
- 238 White W L R G H Siu and E T Reese 1918 *Bull Torrey Botan Club* 75 601-632
- 239 Wigglesworth V B 1918 *Biol Revs Cambridge Phil Soc* 23 408-451
- 240 Willaman J J 1920 *Botan Gaz* 40 221-229
- 241 Wilson W E 1912 *Phytopathology* 32 130-140
- 242 Wolf F T 1951 *Bull Torrey Botan Club* 78 211-210
- 243 Wolf F T 1955 *Bull Torrey Botan Club* 82 313-314
- 244 Wooster R C and V H Cheldelin 1915 *Arch Biochem* 8 311-320
- 245 Young V H 1918 *Plant World* 21 75-81 114-133
- 246 ZoBell C E 1946 *Bacteriol Revs* 10 1-50
- 247 ZoBell C E 1950 *Advances in Enzymol* 10 445-486

It is not always appreciated that a carbon balance implies an oxidation reduction balance and that some error must have been made if the oxidized products cannot be accounted for by the reduced products

The use of resting cells in metabolism studies is another valuable technique but its limitations are not always recognized. The essential feature of the method is that cells are grown in a complete medium washed and put in contact with usually a single compound. This compound may be a putative precursor of a fungal metabolite which is of interest or one may be concerned with the dissimilation of the compound itself. In a short space it is impossible to describe all the pitfalls of this method but some of the more important may be summarized

1 The previous history of the cells may influence the results obtained at least quantitatively (1, 14, 19)

2 The thick inhomogeneous mat of a surface culture is probably always less desirable than the more nearly homogeneous mycelium from a shaken or aerated culture

3 The purpose of the method is defeated if a source of nitrogen is included generally the cells contain and liberate some nitrogenous compounds

4 The endogenous metabolism must be taken into account or even if possible manipulated e.g. by starvation. Thus it is possible that an added compound may simply protect an endogenously formed metabolite from utilization (7) and appear therefore to be a precursor of it

5 Probably the technique is of more value in the study of the dissimilation of a compound than it is in the study of synthetic pathways. A compound e.g. acetate has so many metabolic channels open to it that its effect on a synthesis is difficult to interpret. Similar problems arise in studies of dissimilation but in general the interpretation is clearer provided the products account for the bulk of the compound disappearing. It has been observed (16) that the weight of resting cells may increase during incubation with sucrose presumably this is a manifestation of oxidative assimilation (Chapter 7)

The use of the resting cell technique is discussed by Foster (11) who offers the particularly valuable suggestion that in precursor studies a known low level of carbohydrate—just sufficient to allow minimal synthesis of the product—should be supplied

The details of particular enzymatic methods are beyond our scope here a compendium of such methods has recently appeared (9). Two common problems which arise repeatedly in the fungi may however

only 3 produced significant amounts of soluble carbon compounds. The classical studies of Birkinshaw, Raistrick and others were carried out on common saprophytes growing in a medium relatively high in carbohydrates; even under these conditions numerous species of *Penicillium*, *Aspergillus* and other genera were found to form only carbon dioxide and cell substance. It is safe also to assume that fungi growing in soil do not accumulate the same products which they do in pure culture.

Carbon balance studies provide essential information as to the type of metabolism involved. The methods of Birkinshaw and Raistrick (6) are still valuable and others have been developed more recently (18). Table 1 illustrates a typical carbon and oxidation-reduction balance. Determinations of carbon balances include many on members of the Fungi Imperfecti (2, 4, 5, 12, 15, 20) and a few on the lower phycomyces (8, 13) and on the higher fungi (3, 18, 21).

Table 1. Carbon and Oxidation-Reduction Balances in *Aspergillus Niger* (18)

	mM	Carbon mM	Oxidation Reduction Value per Mole*	Over all Oxidation Reduction Value
Substrate utilized				
Glucose	38.0	228.0	0.0	0.0
Ammonium chloride	1.96		0.0	0.0
Oxygen	109.56		2.0	+18.6
Total		228.0		218.6
Products formed				
Carbon dioxide	88.20	88.20	2.0	176.4
Citric acid	16.13	96.78	3.0	48.4
Extracellular polysaccharide (as glucose)	2.51	15.06	0.0	0.0
Cell				
Carbohydrates (as glucose)	2.93	17.58	0.0	0.0
Chitin	0.114	3.65	-6.0	-0.68
Protein nitrogen	0.765	5.92	-1.78†	-2.72
Lipids	Trace			
Total		227.2		221.4

Carbon recovery = $227.2/228.0 = 99.7\%$

Oxidation-reduction balance = $221.4/218.6 = 1.01$

* See original paper for calculation

† Per atom nitrogen

growth or disappearance of substrate—rather than at the same chronological age

The physiologist often perhaps inevitably overlooks problems of taxonomy and morphology. Particularly in rather difficult genera like *Penicillium* and *Aspergillus* the careful description of cultures with at least some indication that the species has been correctly named is of definite value to the next student of the problem. The question of morphology arises in a different context in the fungi in that conditions of growth or special treatments employed in preparing cells may drastically alter the ratio of different kinds of cells e.g. the ratio of spores to mycelium or even more important the ratio of living to dead cells. It should not be necessary to labor this point.

BIBLIOGRAPHY

- 1 Bernhauer K. 1928 *Biochem Z* 197 281-308
- 2 Birkinshaw J H J H V Charles A C Hetherington and H Raistrick 1931a *Phil Trans Roy Soc London Ser B* 220 55-9
- 3 Birkinshaw J H J H V Charles A C Hetherington and H Raistrick 1931b *Phil Trans Roy Soc London Ser B* 220 99-125
- 4 Birkinshaw J H J H V Charles H Raistrick and J A R Stoyle 1931a *Phil Trans Roy Soc London Ser B* 220 7-54
- 5 Birkinshaw J H J H V Charles H Raistrick and J A R Stoyle 1931b *Phil Trans Roy Soc London Ser B* 220 93-98
- 6 Birkinshaw J H and H Raistrick 1931 *Phil Trans Roy Soc London Ser B* 220 11-26
- 7 Butkevitch W (Butkevich V S) E V Menzhnikova and E I Trofimova 1934 *Mikrobiologiya* 3 319-342
- 8 Cantino E C 1949 *Am J Botany* 36 95-112
- 9 Colowick S P and N O Kaplan (eds) 1950 *Methods Enzymology* Vol 1 New York Academic Press pp 835
- 10 Etheridge D E 1955 *Can J Botany* 33 416-428
- 11 Foster J W 1949 *Chemical Activities of Fungi* New York Academic Press pp 648
- 12 Hockenhull D J D 1950 *J Exp Botany* 1 194-200
- 13 Ingraham J L and R Emerson 1954 *Am J Botany* 41 146-152
- 14 Kluyver A J and L H C Perquin 1933 *Biochem Z* 266 82-95
- 15 Lockwood L B J J Stubbs and C E Senseman 1938 *Zentr Bakteriell Parasitenk Abt II* 98 167-171
- 16 Löffler S and G M Toupinina 1938 *Compt rend acad sci U.R.S.S* 21 301-311
- 17 Pan S C L W Nicholson and P Kolachov 1953 *Arch Biochem Biophys* 42 421-434
- 18 Shu P and J A Thorn 1959 *Can J Botany* 30 252-6
- 19 Thies W 1930 *Zentr Bakteriell Parasitenk Abt II* 82 31-347
- 20 Wells P A A J Moyer and O E May 1936 *J Am Chem Soc* 58 555-58
- 21 Whittaker D R 1951 *Can J Botany* 29 159-17

be mentioned. First it is not always taken into account that the typical preparation contains many enzymes in addition to the one under study. Thus the appearance of isomaltose during the hydrolysis of starch by amylase was once taken as evidence for the structure of the starch molecule. It is now known that it merely denotes the presence of a contaminating transglycosidase (17).

A second common difficulty in enzymatic work is the tendency to read too much into experiments with intact cells. Thus the appearance of unsaturated fatty acids in the medium during the metabolism of saturated fats by intact cells is certainly *prima facie* evidence for the existence inside the cells of a fatty acid dehydrogenase. It is not however a demonstration of the enzyme and should not be described as such. *Ex hypothesi* an intracellular enzyme can be demonstrated unequivocally only in a cell free preparation although properly designed experiments with whole cells may establish a high probability that the enzyme is present.

The variability of fungi is a real problem which arises in connection with virtually all phases of metabolism. However the amount of variation should not be exaggerated and should not be used as a crutch for a failing hypothesis. Complete loss of metabolic capacities is of course well known to be inducible by mutagenic agents and presumably therefore may occur as a result of spontaneous mutation. Most such mutants however are nonviable except under carefully adjusted conditions of culture. It appears that quantitative changes are more common than qualitative. An organism may change in the amount of a given substance formed but still form some. Exceptions will occur and must be taken into account but the progress of a cumulative science requires that the material foundation be assumed constant unless the evidence clearly proves that it is not.

Variability may result from changes in the growth rate such as those described among geographical isolates and cultural variants of *Fomes annosus* (10) rather than from a loss of metabolic capacities. If two strains have the same metabolism but different growth rates analysis at one arbitrary time for a metabolite may give an exaggerated picture of the differences between them. Thus if a given compound is first formed and then utilized after exhaustion of the principal carbon source a rapidly growing strain may complete the process of reutilization at a time when a more slowly growing strain is still in the phase of rapid formation of the compound. It would appear from analysis at this single time that the rapidly growing strain was deficient in the synthesis. To circumvent this difficulty it is necessary to study cultures at the same physiological age—judged by the amount of

molecule but there is no evidence directly on this point in the fungi and actinomycetes


Free sugars appear in the mycelium (130-227) and in the culture medium (84-85) but their significance is not known

The phosphate esters of the monosaccharides are hydrolyzed by phosphatases (phosphomonoesterases). The phosphatases active on glycerophosphoric acid may be presumed to act also on hexose phosphates. No direct study of these reactions in fungi has been made but there is no doubt that at least the common saprophytic fungi and actinomycetes can hydrolyze hexose and pentose phosphates. Commercial enzyme preparations made from fungi contain acid phosphatases (179) and have been used in many laboratories for the dephosphorylation of sugar phosphates. Phosphatases are considered in more detail in Chapter 9.

2 THE METABOLISM OF OLIGOSACCHARIDES

The literature on the enzymatic breakdown of disaccharides and trisaccharides by fungi is extensive but as in so much of fungus physiology relatively few types have been studied. Most of our information comes from work on the Fungi Imperfecti, the wood-destroying basidiomycetes and the Saprolegniaceae. Many other groups of fungi have not been investigated and must therefore be excluded from any generalization.

The most significant recent advance in our approach to the breakdown of oligosaccharides is the concept of transglycosidation. Until the formulation of this concept the breakdown of sucrose, for example, was visualized as a simple hydrolysis



Since sucrose is made up of glucose and fructose units linked by a glycosidic bond, this hydrolytic reaction may be schematized



It now appears that at least some and possibly all of the so-called hydrolytic enzymes are in fact group transferases, capable of transferring a sugar residue to any suitable acceptor. The acceptor may be water, in which case Equation 2 describes the reaction, but other sugars or alcohols can also serve as acceptors. In a particular case, the action of yeast invertase, the fructose residue of sucrose is believed to be transferred as follows (72)

5. Carbon Metabolism

Part II

The Metabolism of Carbohydrates and Related Compounds

The principal topic of this chapter on carbon metabolism is the metabolism of oligosaccharides and polysaccharides anticipated to some degree by the discussion of their role in carbon nutrition (Chapter 3). The related compounds included are the amino sugars and the polyhydric alcohols. Discussion of the monosaccharides is limited to a few synthetic reactions.

1 THE METABOLISM OF MONOSACCHARIDES

The breakdown of monosaccharides by fungi has been considered separately as respiration in Chapter 7. A second aspect of monosaccharide metabolism is the synthetic one in which the sugar or a derivative of it is incorporated into larger molecules. As an example glucose in the glucolipid ustilic acid is found by isotope studies to be in part incorporated as such and in part resynthesized from triose before incorporation (30). In this case and in the formation of glucose polysaccharides no modification of the basic glucose molecule need be assumed. However the organism may synthesize a complex molecule containing sugar moieties which were not provided in the culture medium. Examples include galactose in polysaccharides (Chapter 2) and amino sugars in certain antibiotics (p. 122). In these processes it is reasonable to assume that the new sugar or a derivative of it (e.g. a phosphate ester) is made and then incorporated into the larger

Table 1 Transglycosidase Activity in Fungi

Enzyme Activity	Substrate	Fungi
Transfructosidase	Sucrose	<i>Aspergillus</i> spp (9 10 16 66 169 170)
	Raffinose	<i>Aspergillus</i> spp <i>Penicillium spinulosum</i> (15)
Transglucosidase	Maltose	<i>Aspergillus</i> spp (47a 169 173 174) <i>Pennis- cillium chrysogenum</i> (202)
	Isomaltose	<i>Aspergillus oryzae</i> (171 172)
	Cellulose	<i>Aspergillus niger</i> (11 12)
		<i>Chaetomium globosum</i> (48)
		<i>Myrothecium verrucaria</i> (119)
Transgalactosidase	β Nitrophenyl	<i>Aspergillus</i> spp (113 226)
	β glucoside	<i>Penicillium chrysogenum</i> (226)
	Lactose	<i>Aspergillus oryzae</i> (244 245)




acting on sucrose or raffinose (Table 1). A fructose-enzyme complex is postulated as in Equations 3, 4, and 5 above. In *Penicillium spinulosum* there is no evidence for any sucrase other than the transfructosidase but in *Aspergillus oryzae* there is probably a second enzyme acting on sucrose (16). Paper chromatography of the enzymes of *A. oryzae* similarly suggests that more than one sucrase is present (112).

Sucrase acting as it does on β fructofuranosides is instrumental in the metabolism of raffinose and stachyose.

Mandels (143) presents indirect evidence for the non-hydrolytic metabolism of sucrose by spores of *Myrothecium verrucaria* possibly to be credited to a sucrose phosphorylase of the type known in bacteria.

Maltose is used as a source of carbon by almost all fungi (Chapter 3). Correspondingly beginning with the work of Bourquelot (33) we find that the enzyme maltase is found in almost every fungus and actinomycete investigated although quantitative differences between strains are common (127). It has been claimed that *Polyporus betulinus* lacks maltase (138) and the chytrids which are unable to grow on maltose (Chapter 3) may be found to lack the enzyme.

Maltase is an α -glucosidase and is found in bacteria, animals, and higher plants. The earlier separation of fungus from yeast maltase as a glucomaltase is probably not valid but the two maltases are not identical (89). As with sucrase recent investigations have established in fungi the existence of a transglycosidase acting on maltose to form a series of transient oligosaccharides (Table 1). The existence of a specific hydrolase for maltose may be doubted but it is not yet known whether the transglucosidase of *Aspergillus niger* and *A. oryzae* the only fungi so far studied is of general occurrence as a single



The fructose enzyme complex is of course hypothetical but it is necessary to explain the observed reactions. Because of the irreversibility of the fifth reaction and the great excess of water present in a long term experiment virtually all of the fructose residues are transferred to water and the result is described grossly by Equation 2. The technique of paper chromatography applied in short term experiments has been essential to the elucidation of this type of group transfer. The characteristic of transglycosidase action on disaccharides is the transient appearance of new oligosaccharides.

Disaccharides Most of the work on this topic has been concerned with the breakdown of sugars especially the common disaccharides such as sucrose, maltose, and lactose. Synthesis of disaccharides occurs in the fungi, the best known case being trehalose (p. 108). Sucrose is attacked by an enzyme variously known as invertase, saccharase, glucosaccharase, and sucrase; the name sucrase is preferred. The first indication of its presence in fungi came from the work of Béchamp (18-19). As might be expected from the frequent utilizability of sucrose as a carbon source (Chapter 3), the enzyme is widely distributed in the fungi. Sucrase is somewhat rare in the Mucorales—again in agreement with nutritional data—but does occur in some species (93-203). Sucrase is found in some isolates of *Streptomyces* (101-110) not in any of five species of dermatophytes (229).

Fungus sucrase may be entirely intracellular (148) but is usually found also in the medium at least after the onset of autolysis (58). It occurs in spores (71-141) in *Mycrothecium verrucaria* and *Aspergillus luchuensis*; there is evidence that it or a sucrose phosphorylase is located at or near the spore boundary (142). This boundary location is thought to be characteristic of sucrase in the yeast cell (62-155).

It has long been argued that the predominant sucrase of fungi is a glucosaccharase differing from the fructosaccharase of yeast (121); the history of this controversy is reviewed by Neuberg and Mandl (156). The question has now been reexamined in the light of the finding that both yeast and fungus sucrases are probably transfructosidases. It appears that the two sucrases differ but not in the way formerly thought and that it is not necessary to postulate a glucosaccharase (15-66).

Transfructosidase activity has been demonstrated in several fungi

reserve carbohydrate formed early in metabolism and used up later (43) under some conditions reutilization may not occur during the life of the fungus (164). An adequate picture of the course of trehalose synthesis and breakdown is not yet available. In yeast trehalose is synthesized as its phosphate from glucose and a glucose phosphate with uridine diphosphate glucose as coenzyme (68, 125). Whether the same mechanism is operative in the filamentous fungi remains to be seen.

Melibiose, an α -galactosido-glucose occurring naturally as a constituent of raffinose, is split by an α -galactosidase (melibiase) known in yeast and higher plants and reported from *Aspergillus* spp. (100, 230).

Trisaccharides and Tetrasaccharides Raffinose, composed of galactose, glucose and fructose, is split by enzymes from most fungi which have been tested. This is to be expected in view of the fact that sucrase splits raffinose to melibiose and fructose. The complete hydrolysis of raffinose requires both sucrase and melibiase; melibiase alone yields sucrase and galactose. In all likelihood hydrolysis of raffinose by unspecified fungal enzymes should usually be attributed to sucrase (2, 15).

Melenitose is hydrolyzed at least partially by several fungi (15, 122). Detailed studies of the reaction are lacking; complete hydrolysis requires an enzyme able to attack the disaccharide, turanose.

Gentianose is completely hydrolyzed to glucose and fructose by an enzyme preparation from *Aspergillus niger* (40, 41); the effective preparation has been shown (42) to contain a β -glucosidase acting on the gentiobiose moiety of gentianose.

Stachyose, a tetrasaccharide composed of galactose, glucose and fructose in the ratio 2:1:1, is split by enzymes from *Penicillium spinulosum* and *Aspergillus* spp. completely to its constituent sugars (15, 228). Presumably sucrase and an α -galactosidase are required.

3 THE METABOLISM OF OTHER GLYCOSIDES

The oligosaccharides just considered are one class of glycoside in which the glycosidic bond connects two monosaccharide residues. The same bond can be formed between a sugar and certain other molecules; the non-carbohydrate moiety of the resulting compound is then termed an aglycon. Information on the metabolism of glycosides of this type by fungi is restricted to a few compounds in all of which glucose or mannose is the sugar involved.

The β -glucosides have received more attention than any others in

enzymatic entity Transglucosidases which act on maltose to form polysaccharides are known in bacteria but have not yet been reported from fungi

The disaccharide lactose is split to its constituent hexoses by the enzyme lactase a β galactosidase The enzyme has been reported in several fungi and actinomycetes but is apparently not of such general occurrence as either sucrase or maltase Some of the early work on lactase in commercial enzyme preparations and in basidiomycete sporophores may be open to question because of the possibility of bacterial contamination

The adaptive formation of lactase in *Neurospora crassa* has been proved (Chapter 3) A purified enzyme from *Aspergillus oryzae* acts as a transgalactosidase galactose residues are transferred to alcohols to lactose itself to form a trisaccharide or to water to yield free galactose (Table I) Whether all action of lactase can be credited to this enzyme is not yet known

Cellobiose is formed during the hydrolysis of cellulose Although it is common to refer to the enzyme splitting it as cellobiase there is no reason to think that this enzyme differs in any way from β glucosidases acting on other β glucosides Specific assays with cellobiose as substrate have been made infrequently but it seems that most fungi can split it to its constituent glucose residues (191) Resting and growing cells of *Aspergillus niger* form new oligosaccharides from cellobiose (11 12) and additional evidence for transglucosidase action on cellobiose has been obtained with other fungi (Table I) Cellobiose is synthesized by *Ustilago zae* as a component of ustilagic acid (128)

Trehalose an α glucosido-glucose is not attacked by the usual α glucosidases but rather by a special enzyme trehalase found in animals plants and fungi (89) Early work on trehalase in fungi is reviewed by Czapek (59) this and more recent data indicate the enzyme is common (203 256) The spores of *Aspergillus luchuensis* contain measurable amounts of the enzyme (139) In *Lycogala* sp a myxomycete trehalase of the fruit body disappears during ripening (108)

Trehalose is the only disaccharide to be formed free in appreciable amounts by fungi The discovery of the sugar in ergot (*Claviceps purpurea*) by Wiggers (254) and early reports of its occurrence in other fungi are reviewed by Zellner (261) and Obaton (164) Trehalose is especially common in the basidiomycetes but is found also in many other fungi (31 36) Usually it constitutes 2 to 3 per cent of the mycelial dry matter (108 193 227)

In all probability trehalose in fungi has the role of a translocatable

least in its early phases results in the formation of dextrans reducing sugars appearing only later

2 Beta amylase (exoamylase saccharifying amylase) of resting seeds
Attack on starch liberates maltose immediately

It has been repeatedly found with highly purified or crystalline preparations that the amylase of *Aspergillus oryzae* is of the α -amylase group (49 73 235). So also apparently is the amylase of *Streptomyces* spp (213). The crude amylase of *A. oryzae* contains more than one component as judged by electrophoretic data (82).

The enzyme of *Rhizopus delemar* termed gluc amylase (177 178) can be separated from α -amylase and resembles β amylase in certain respects. There seems however no reason to believe that a typical β amylase occurs in the fungi (127 129 146).

Table 2 summarizes some of the properties of the purified amylase of *Aspergillus oryzae* and its differences from other α -amylases.

Amylase activity defined as the ability of cell free materials to hydrolyze starch or glycogen is virtually universal in the fungi and actinomycetes. *Penicillium digitatum* does not form detectable amylase in culture (103) and in view of its inability to grow on starch (Chapter 3) is presumably deficient for the enzyme. Early studies on fungal amylase are summarized by Thaysen and Galloway (230) and the industrial technology of the enzyme is reviewed by Linglykke et al (123).

Table 2 Properties of Alpha amylases (74)

	Source of Crystalline Alpha amylase				
	Malt	<i>Aspergillus oryzae</i>	<i>Bacillus subtilis</i>	Pig Pancreas	Human Saliva Pancreas
Activity per mg N	2350	2400	3600	4000	6200
Activity per mg protein	315	310	500	630	980
Nitrogen per cent	13.4	12.9	15.8	15.8	15.8
Optimum pH for activity	4.7-5.4	5.5-5.9	5.3-6.8	6.9	6.9
Optimum pH for stability	4.9-9.1	5.5-8.5	4.8-8.5	7.0-8.5	4.8-11
Isoelectric point	5.6	circ 4.2	?	5.3	5.3
Absorption maximum (m μ)	280	280	280	280	280
Saccharogenic/dextrinogenic ratio	9.8	9.8	9.8	9.8	9.6
Activation by Ca	+	-	-	-	-
Activation by Cl	-	-	+	+	+
Energy of activation	7050	10650	13350	13500	13350
			0 to 15		

this general category. The enzyme β glucosidase (formerly termed emulsin) occurs in almost all of the fungi and actinomycetes which have been examined for it (31 38 39 101 147 201 229) although it is not always found in the medium (191) and may not be so nearly universal as for example maltase or amylase. *Fusarium oxysporum* var *lycopersici* forms detectable amounts of the enzyme within the living infected plant (60).

Although it is common practice to speak of a single β glucosidase hydrolyzing all β glucosides—including the β linked disaccharides cellobiose and gentiobiose—it seems more likely that a group of related enzymes is involved (238) or that at least there are distinct components of the enzyme (111, 112). The β glucosidases of *Aspergillus niger* and *A. oryzae* for example act at different rates on different synthetic β -glucosides (151). The enzyme of *A. niger* has been purified about 10 000 fold (160 161).

During the action of β glucosidase on synthetic β glucosides the glucose residue can be transferred to alcohols indicating trans glucosidase activity similar to that with cellobiose as substrate (Table I).

Alpha glucosides in principle at least should be attacked by maltase or α glucosidase. However maltases from different sources have different behavior toward e.g. methyl α glucoside and the maltase of fungi seems to be of minimum activity toward this substrate (89). A crude preparation from *Aspergillus niger* hydrolyzes methyl α -glucoside slowly (65) and in some fungi there is an α glucosidase distinct from maltase (15) but the problem has been too little studied for any definite conclusion to emerge. *Aspergillus niger* cultivated with methyl α galactoside forms the corresponding α galactosidase (182).

An enzyme of *Streptomyces griseus* converts mannosidostreptomycin to streptomycin (175). The enzyme has been found to split synthetic methyl α mannoside (99) and may therefore be termed an α mannosidase. The same action with methyl α mannoside as substrate is known in *Aspergillus niger* (98).

4 THE METABOLISM OF POLYSACCHARIDES

The Breakdown of Starch and Glycogen These two related polysaccharides are hydrolyzed by amylases (diastases) of which there are in nature two general types (154).

1 Alpha amylase (endoamylase dextrinizing amylase liquefying amylase) found in animals and microorganisms. Attack on starch at

Crude cellulase preparations from several fungi are alike in that their optimum pH is in the acid range 4.5 to 5.5 (90, 163, 204); the optimum for the enzyme of *Stachybotrys atra* is however higher pH 6.5-8.0 (230a). Studies on *Tricholoma fumosum* indicate that the enzyme is formed only by cells in contact with cellulose or lichenin and that the enzyme in a culture is predominantly extracellular (163). Cellulose is apparently attacked at several points along the chain of glucose residues inasmuch as in the early stages of attack on cellulose there is a decline in optical density without a corresponding liberation of sugars (162, 163a). Other characteristics of fungal cellulases are summarized by Siu (214).

For the present a general statement on the nature of cellulase is not possible. The classical view (183) postulates at least two enzymes: a cellulase splitting the long polymeric chain to cellobiose residues and a cellobiase (β -glucosidase) splitting the disaccharide to glucose. Although substantial evidence exists for the appearance of cellobiose as an intermediate (119, 159a, 250) the process is probably more complex. Two theories both based on work with *Myrothecium verrucaria* have been proposed recently. The first of these postulates a minimum of three enzymes as required for cellulose hydrolysis (83, 91a, 150, 190, 191) with only one of them characteristic of the cellulolytic organisms and the others occurring also in non-cellulolytic fungi. In sharp distinction to this view other studies on the same organism (251, 253) have resulted in the isolation of an apparently single enzyme purified to electrophoretic homogeneity with a molecular weight of 63,000. This single enzyme hydrolyzes cellulose to glucose without the obligatory participation of cellobiose as an intermediate although cellobiose may be formed during the process. The enzyme acts at different rates on cellobiose, cellotriose and higher oligosaccharides of this group (252) and is inhibited by sulphhydryl reagents (14). It is obvious that further work on the enzyme will be necessary before any general theory can be formulated.

The Breakdown of Hemicelluloses and Pentosans The hemicelluloses are poorly defined plant constituents particularly of wood and gums. The principal components of the hemicelluloses are polysaccharides and polyuronides. The literature reports essentially two types of work: that on hemicellulose fractions of unknown composition and that on more or less purified polysaccharides derived from hemicellulose.

Crude hemicellulose fractions from wood are attacked by enzymes of *Aspergillus oryzae* yielding free sugars and a soluble polysaccharide




Figure 1 Amylase liberation by *Aspergillus oryzae*. Redrawn from Cawther and Lennox (58) by permission of the Australian Journal of Biological Sciences

Amylase is with some exceptions (61) found in the culture medium as well as in the cells. The amount of enzyme is usually increased by inclusion of starch in the growth medium (92-210) and is affected by the nitrogen source (210) and by small amounts of salts (69). Figure 1 illustrates the time course of appearance of extracellular amylase in *Aspergillus oryzae*; it is to be noted that the peak of amylase activity in the culture filtrate is reached only after the onset of autolysis.

It has been reported (26) that the culture filtrate of *Streptomyces microflavus* forms gentiobiose during the hydrolysis of starch.

The Breakdown of Cellulose Since many fungi utilize cellulose as a carbon source (Chapter 3) it is to be expected that an enzyme or enzymes are released from the cell and hydrolyze cellulose to soluble sugars. Cellulose hydrolyzing preparations free of living cells have indubitably been made from several fungi (83-191-214) and distinguished from enzymes acting on lichenin and hemicelluloses (90). However, two important limitations apply to most of the reported work on cellulase: (1) native cellulose is relatively resistant to attack and most assays of cellulase have therefore utilized various types of modified cellulose; (2) most data derive from studies with crude cellulase preparations contaminated with other carbohydrases and with interfering proteins (232). It is difficult therefore to assign more than a descriptive value to much of the work so far reported.

study of the pectic materials. Historically interest in the pectic enzymes was first stimulated by the discovery (13-217) that crude enzyme preparations from plant pathogenic fungi macerate plant tissues by destruction of the middle lamella, the cementing material between cells. First thought to be a cellulase, the tissue macerating enzyme has also been termed cytase. Other early studies are reviewed by Brown (46).

The insoluble pectic materials as they are found in plants are now called protopectin (116). Neither the structure nor the composition of protopectin is known. The corresponding enzyme protopectinase (pectosinase) is defined in principle as the enzyme converting protopectin to water-soluble pectinic acids (255). Operationally the enzyme is defined by its ability to macerate blocks of living plant tissue. It is found in pathogenic and non-pathogenic fungi belonging to all of the major taxonomic groups (Table 3).

Table 3 Occurrence of the Pectic Enzymes in the Fungi*

Enzyme	Occurrence
Protopectinase	<i>Aspergillus</i> spp (61-222) <i>Botrytis</i> spp (46-237) <i>Byssoschlamys fulva</i> (17-105) <i>Chaetomium globosum</i> (222) <i>Daedalea flauida</i> (32) <i>Fomes fraxineus</i> (152) <i>Fusarium</i> spp (51-149-211) <i>Gloeosporium fructigenum</i> (<i>Glomerella cinulata</i>) (149) <i>Lenites sepiaria</i> (263) <i>Macrophomina phaseoli</i> (185) <i>Metarrhizium</i> sp (222) <i>Monilia fructigena</i> (237) <i>Mucor racemosus</i> (93) <i>Phytophthora erythroseptica</i> (149) <i>Polyporus</i> spp <i>Polystictus</i> spp (32) <i>Pythium debaryanum</i> (4-97) <i>Rhizoctonia</i> sp (258) <i>Rhizopus</i> spp (94-95) <i>Stereum purpureum</i> (145) <i>Trametes</i> spp (32)
Pectin methylesterase	<i>Aspergillus</i> spp (81-114) <i>Botrytis cinerea</i> (80) <i>Fusarium</i> spp (87-241) <i>Penicillium</i> spp (114-176) <i>Sclerotinia cinerea</i> (61) <i>Sclerotium bataticola</i> † <i>Thielaviopsis basicola</i> † (231)
Pectin polygalacturonase	<i>Aspergillus</i> spp (6-64-81-114-208) <i>Botrytis cinerea</i> (81-114-257) <i>Byssoschlamys fulva</i> (102-192) <i>Colletotrichum phomoides</i> (185a) <i>Fusarium</i> spp (47-185a-211-241-243) <i>Mucor</i> spp (185a) <i>Neurospora</i> spp (64-197) <i>Penicillium</i> spp (64-176-185a-243) <i>Rhizopus nigricans</i> (185a) <i>Stachybotrys atra</i> <i>Syncephalastrum racemosum</i> <i>Thamnidium elegans</i> <i>Trichoderma</i> spp (64-185a) <i>Trichellium albo-atrum</i> (205)

* Including only the three major enzymes; for other enzyme activities see text.

† Assayed by hydrolysis of half-calcium salt of methyl D-tartaric acid (157).

(166-167) As might be expected the wood destroying basidiomycetes are especially active in the hydrolysis of hemicelluloses (31-189) the gross utilization of hemicelluloses has been mentioned earlier (Chapter 3) The only other fungi which have been studied are some members of the Saprolegniaceae (21-201) and of the genus *Phytophthora* (147) all of which form reducing sugars from the crude hemicellulose of date endosperm

The most abundant of the pentosans are xylans the rarer araban are found in association with pectic substances The limited literature on fungus metabolism of xylans is reviewed by Voss and Butter (240) Enzymes (xylanases) attacking xylan have been found in *Aspergillus oryzae* (90) and *Glomerella cingulata* (96) The xylanase of *Chaetomium globosum* is formed in appreciable amounts only if the organism is grown on xylan and consists of an exocellular component converting xylan to xylobiose and an intracellular enzyme hydrolyzing the xylobiose (217-218)

Arabins—pentosans made up largely of L arabinose—are hydrolyzed by a crude enzyme preparation from *Aspergillus oryzae* (67)

Simpson (212) surveyed the action of 112 isolates of 24 genera on a purified pentosan of unspecified composition An active hydrolysis was brought about by the culture filtrate of about half of the isolates tested the most active filtrates were from cultures of strains of *Aspergillus*, *Fusarium*, *Ficoidium*, and *Trichoderma* Of the 54 active strains only 6 formed the enzyme during growth on glucose in the rest growth on the pentosan—replaceable in some instances by free pentoses—appeared to be necessary for enzyme synthesis In the genus *Streptomyces* 5 of the 19 isolates tested formed an extracellular enzyme active on the pentosan

A purified arabogalactan from *Larix* is split by an enzyme from *Aspergillus niger* (189) Mannans are hydrolyzed by an enzyme of *Chaetomium globosum* which is apparently distinct from the enzyme acting on xylans (217)


The chemistry of the substrates is poorly known and the nature of the enzymes involved in their hydrolysis is uncertain In particular it is not possible from the data available to decide on the possible identity or nonidentity of enzymes which have been studied by different workers It is generally accepted that the enzymes attacking the pentosans and the crude hemicelluloses are distinct from cellulose hydrolyzing enzymes and from lichenase (180) but beyond this point specificity is not known

The Breakdown of Pectic Substances Perhaps nowhere in the field of carbon metabolism is the confusion so great as it is in the

The enzyme occurs in commercial enzyme preparations of pectin polygalacturonase (75) and in the fungi listed in Table 3. The enzyme of *Fusarium oxysporum* var *lycopersici* appears to differ from that of commercial preparations from other fungi (241) and the latter in turn differs from the pectin methylesterase of higher plants (145).

Production of extracellular pectin methylesterase by *Botrytis cinerea* and *Aspergillus niger* occurs if pectin is the growth medium but only to a negligible degree with glucose as the carbon source (3, 80, 81). The extracellular enzyme of *Penicillium chrysogenum* is formed only if pectin, pectic acid, or chemically related substances are in the growth medium (176). Recently the role of pectin methylesterase in the pathogenicity of fungi to plants has received attention.

The third major category of the pectic substances is that of the pectic acids derived from pectinic acids by deesterification. They are colloidal polygalacturonic acids with molecular weights ranging from 25,000 to 100,000. The molecule consists of D-galacturonic acid units and may be visualized as follows:



The enzyme hydrolyzing the pectic acids has been named variously pectinase, pectolase, and polygalacturonase. The most unambiguous name appears to be pectin polygalacturonase (116). It acts on the de-esterified portions of pectinic acids or on the pectic acids resulting from complete deesterification of pectinic acids. It is often contaminated with pectin methylesterase, and the mixture of the two enzymes will, of course, act on pectinic acids both by deesterification and by hydrolysis. The product of pectin polygalacturonase activity on a polygalacturonide is eventually free D-galacturonic acid. The properties of the enzyme have been studied extensively by Matus (144).

Table 3 lists the fungi which have been shown to form the enzyme as assayed on pectinic acid or polygalacturonic acids, i.e., excluding assays of tissue macerating enzyme. Two comments are apposite to the problem of the occurrence of the enzyme in fungi. First, it is probable that the fungi producing protopectinase always form pectin polygalacturonase or that the two enzymes are identical. If either of these hypotheses is true, then the sources of protopectinase given in Table 3 must be included, adding numerous phycomycetes and

The indefinite nature of the substrate the crudity of the enzyme assay and the fact that other enzymes may solubilize the pectic materials of plants (116) are compelling reasons to doubt the real existence of protopectinase. This doubt is reinforced by the finding (102) that a partially purified pectin polygalacturonase from *Byssolhamys fulva* causes disintegration of tobacco tissue. Although it is possible of course that the preparation was contaminated with a protopectinase it seems more probable that pectin polygalacturonase has the ability usually ascribed to protopectinase to macerate tissue. It is significant in this connection that all protopectinase preparations examined contain pectin polygalacturonase (115).

The present nomenclature of the pectic enzymes was first clearly formulated in 1927 (61) and fully worked out only in 1944 (117). Consequently the earlier work on these enzymes employs different and often highly individual nomenclature. In particular the pectinase of the older literature is usually synonymous with what is now called protopectinase i.e. the tissue macerating enzyme. Pectinase as now understood is a synonym for pectin polygalacturonase. The nomenclature used here is that of Kertesz (116).

The pectinic acids are colloidal macromolecules composed of α -D anhydrogalacturonic acid units linked together through 1,4- α -glycosidic bonds. The carboxyl groups of some of the galacturonic acid residues are esterified with methyl alcohol. Certain pectinic acids of specific properties are designated pectins. Pectinic acids are obtained from the crude plant pectic materials by partial hydrolysis.


Excluding protopectinase from consideration the known and postulated pectic enzymes of fungi are of three types depending on their action on pectinic acids. They are

1 Pectin methylesterase catalyzing the hydrolysis of the methyl ester groups of pectinic acids and converting pectinic acids to pectic acids (polygalacturonic acids)

2 Pectin polygalacturonase hydrolyzing the pectic acids to free D-galacturonic acid

3 Pectin-depolymerase a less well known enzyme thought to cause a partial hydrolysis of pectic acids to smaller units without the appearance of free galacturonic acid

Pectin methylesterase (pectase pectinesterase pectinmethoxylase) was originally defined (61) as an enzyme converting soluble pectin (pectinic acid) into insoluble calcium pectate. Its action may be schematized as



enzymes as in that of so many other enzymes is to determine the action of single enzymes on substrates of definite and known composition. Once such information is available there will be some hope of successfully analyzing the more complex situation in nature.

The Breakdown of Other Polysaccharides It is generally considered that a special enzyme, inulase, catalyzes the breakdown of the plant fructosan, inulin. The enzyme was discovered in *Aspergillus niger* by Bourquelot (37) and has since been found in many other fungi (25, 59, 79). It has often been reported that cultivation on inulin as carbon source increases the amount of enzyme (20, 184, 261) but some inulase is always formed whatever the carbon source. No systematic study of the enzyme or its occurrence has been made; some properties of fungal inulase have been determined by Pigman (179) and it is possible that two different enzymes are responsible for the activity (153). On the other hand, there is some evidence that inulase is a transfructosidase active also on sucrose (150a).

A bacterial dextran, composed of anhydroglucose units linked by 1,6- α -glucosidic bonds, is hydrolyzed by a dextranase present in the mycelium and culture fluid of several fungi, e.g. *Penicillium funiculosum*, *Lenticiellium coccorum*, and *Spicaria violacea* (106, 234). The enzyme is formed only if dextran is present in the medium.

Limit dextrin, the polysaccharide fraction remaining after starch hydrolysis by α -amylase, is hydrolyzed by a limit dextrinase from *Aspergillus oryzae*; the enzyme has been crystallized (235).

A commercial enzyme preparation, probably made from an *Aspergillus* sp., hydrolyzes a galactogen of snail eggs, yielding D- and L-galactose (248).

Lichenin, the polysaccharide of Irish moss (*Chondrus crispus*), resembles cellulose except for its high proportion of 1,3-glycosidic bonds. It is hydrolyzed by an enzyme found in animal and plant material (180) and in *Aspergillus oryzae* (200) and *Merulius lacrymans* (181). Lichenase is probably distinct from cellulase and xylanase (78, 90).

The mucopolysaccharides are complex polysaccharides yielding amino sugars on hydrolysis (16); examples include hyaluronic acid, chitin, agar, and unidentified cell wall materials. Among the enzymes attacking these materials, the bacterial hyaluronidase and lysozyme are fairly well known, but the mucopolysaccharases of fungi are not. Chitinases are known among actinomycetes and both higher and lower fungi (49a, 233); the enzyme or enzymes responsible for chitin hydrolysis in *Streptomyces* sp. are adaptive (194). The lytic enzymes of

basidiomycetes to a list which is predominantly composed of ascomycetes and imperfect fungi. Second the fact that many fungi form pectin polygalacturonase only if pectinic acids or related materials are present in the medium means that a complete survey of representative fungi should use such a medium in addition. In any such survey the pH of the medium should be kept below 6.0 by reason of the rapid inactivation of the enzyme at higher pH values (116). From the data available and in the light of these considerations it seems likely that most fungi are able at least under favorable conditions to elaborate pectin polygalacturonase.

The pectin polygalacturonase of fungi differs in its properties from the corresponding enzymes of yeast and higher plants (118-133). The enzyme is highly specific to pectic acids failing to hydrolyze other galacturonides or other uronides (131). Separation from pectin methyl-esterase has been accomplished (131-136). It is not yet possible to assert that pectin polygalacturonase is a single enzymatic entity. One view (114) is that it is a single enzyme acting by random scission of the pectic acid molecule to form shorter galacturonic acid chains and eventually free galacturonic acid. On the other hand it has recently been suggested that three different enzymes are involved in *Aspergillus foetidus* (6-63).

The pectin polygalacturonase of *Penicillium chrysogenum* and of *Fusarium* spp. appears to be elaborated adaptively in media with pectin as carbon source but not in glucose media (47-176) but the same enzyme in other fungi is formed in glucose media (64-80-81). All these experiments were carried out by growing cells from a spore inoculum in media with or without the presumed inducing substrate hence a possible effect of mutation or selection was not rigorously excluded.

Other enzymes depolymerases acting on pectins have been described in plant parasitic and saprophytic fungi (17-19a-88-195-209). These enzymes share in common the characteristic that only certain of the glycosidic linkages of the substrate are attacked resulting in the liberation of short chain polyuronides. The existence of such enzymes in fungi has been challenged (192-197) but the evidence especially that from work on *Neurospora crassa* (195) is strong.

Assay methods for the pectic enzymes are reviewed by Kertesz (116). Cup plate assays have been devised (53) and may be particularly useful in survey studies. There is some doubt (116) that assay of pectin methyl-esterase by hydrolysis of the half calcium salt of methyl p-tartronic acid (157) is valid.

It seems that the immediate problem in the study of the pectic

phosphorylase (22f). Presumably other environmental factors are influential on a priori grounds it is to be expected that the concentration of available carbohydrate and the carbon nitrogen ratio of the medium are important. Oxygen is required for the synthesis of polysaccharide by *Aspergillus niger* (206).


The function of fungal polysaccharides is almost certainly as a reserve food and the data of Ergle (70) on glycogen utilization during the germination of sclerotia bear out this contention. This view of polysaccharides as one form in which fungi can store energy was first clearly stated by Chrząszcz and Tarkow (53) in opposition to an earlier view of polysaccharide formation as a symptom of a deranged metabolism. It should be noted that under some cultural conditions the polysaccharide formed may never in fact be utilized, cell death and autolysis intervening before the carbohydrate supply becomes limiting and before the reserves are attacked.

Several fungi pathogenic to man form polysaccharides often associated with a capsule and often immunologically active. Those of *Coccidioides immitis* and *Cryptococcus (Torulopsis)* spp. are listed in Chapter 2; others are reviewed by Nickerson (158).

D THE METABOLISM OF AMINO SUGARS

The only amino sugar which has been studied extensively is D-glucosamine (2-amino-D-glucose); interest in which arises primarily from the fact that its N-acetyl derivative is the basic unit of chitin, the cell wall material of most fungi (Chapter 2).

Conceivably the conversion of glucose to glucosamine could proceed with glucosone $\text{CH(OH)-(CHOH)}_3\text{-CO-CHO}$ as intermediate and glucosone is formed by *Aspergillus parasiticus* at least under some conditions (29). However in *Neurospora crassa* extracts glucosamine is formed by transamination from glutamine (126) as follows:



It appears from other work that the hexose phosphate involved directly in the reaction is fructose-6-phosphate (26).

Glucosamine appears in chitin and in animal mucopolysaccharides as the N-acetyl derivative. Introduction of the acetyl group involves acetyl-coenzyme A in both animals and *Neurospora crassa* (52, 126). In *N. crassa* it is not certain whether free glucosamine or its phosphate is the compound acetylated.


Amino sugars are frequently found as constituents of antibiotics

actinomycetes e.g. the enzyme system of *Streptomyces albus* (134 135) act on cell wall polysaccharides and can therefore tentatively be classed with the mucopolysaccharases. Enzymes of this type have been reported frequently from actinomycetes (114a 249) and must be present in those fungi in which the chitin breaks down during autolysis (216).


Certain marine actinomycetes liquefy agar (107 219). The enzyme responsible has been called gelase; it is evidently either rare or absent in terrestrial fungi and actinomycetes since agar is used so generally in culture media that gelase-producing organisms would have long since been detected if they were common.

The Synthesis of Polysaccharides The known fungal polysaccharides are reviewed elsewhere (Chapter 2); it will be recalled that they include chitin, polyhexoses and more complex polysaccharides. The most common hexose component is glucose; the next most common galactose, since all of the galactans are formed in media with glucose or sucrose as carbon source; it seems probable that a galactose-4-epimerase is present as in yeast, catalyzing the conversion of glucose to galactose (124).

The metabolic origin of the polysaccharides of fungi has been little investigated but may be expected to follow the pathways known in other organisms. The two best known mechanisms of polysaccharide synthesis in bacteria are in outline



Disaccharide Poly- Mono- Hexose




The first of these is a transglycosidation of which the action of levansucrase is an example; the second is catalyzed by a phosphorylase and is exemplified by glycogen formation in the animal. Levan synthesis in *Aspergillus sydowii* spores may be a transglycosidation; it can be effectuated by a broken cell preparation and the levan is formed during growth on sucrose but not during growth on monosaccharides (120). The polyglucose of *Torulopsis rotundata* on the other hand is probably synthesized by a phosphorylase (139). Evidence on polysaccharide biogenesis in the fungi is so far limited to these two cases.

The polysaccharides of some fungi are formed more abundantly in acid than in neutral media (27 140 207) but the reverse is true of sclerotiose formation (1). In yeast inhibition of polysaccharide formation by alkalinity is believed to result from an inhibition of

melles (24). Intact cells of *Fusarium lini* oxidize erythritol to erythrose (86) and it is likely that similar reactions occur in those fungi which themselves accumulate sugar alcohols.

Glycerol The formation of glycerol by fungi is mentioned in Chapter 7 as an indication of the functioning of the Embden Meyerhof respiratory pathway. The available data (86, 187, 188) are consistent with the metabolism of glycerol via this pathway, i.e.



No enzymatic studies have however been made and the role of phosphorylated intermediates is only conjectural.

Glycerol phosphates are hydrolyzed by phosphomonoesterases (phosphatases) which also attack other organic phosphates (196). Hydrolysis of organic phosphates at acid pH is brought about by mycelial extracts of *Actinomyces israeli* (104) and of fungi (55, 58, 198, 199). In *Trichophyton rubrum* cytochemical tests suggest the presence of another phosphomonoesterase active at alkaline pH values (159) and enzymes of this type (alkaline phosphatases) have been reported in other fungi (50, 236). These and other phosphatases are considered elsewhere (Chapter 9).

Baba (7, 8) isolated an unusual metabolite monoallyl phosphoric acid $\text{CH}_2=\text{CH}-\text{CH}_2-\text{OPO}_3\text{H}$ from cultures of *Aspergillus niger* metabolizing glycerol and showed in addition the presence of an enzyme, a phosphatase, able to dephosphorylate the compound.

Glycols Exogenously supplied glycols are oxidized by some fungi, e.g. *Fusarium lini* converts ethylene glycol to glycolaldehyde (86) and *Aspergillus niger* dehydrogenates 1,2-butanediol to hydroxymethylethyl ketone (246). The dehydrogenation of ethylene glycol may be effected by alcohol dehydrogenase known to catalyze this reaction in other systems.

Fusarium lini also oxidizes 2,3-butylene glycol to acetoin ($\text{CH}_3-\text{CO}-\text{CHOH}-\text{CH}_3$) (86). Acetoin occurs as a metabolite of sugar or pyruvic acid in fungi and actinomycetes (109, 221, 225) but its importance and the pathway of formation in the fungi are not known. In yeast but not in bacteria it seems that ketols of this type are formed with the participation of acetylcoenzyme A and the free aldehyde (215).


including cordycepin of *Cordyceps militaris* (20) and many of the actinomycete antibiotics (22-56). Whether they are related to toxicity is uncertain but the possibility is at least suggested by the number of instances uncovered.

6 THE METABOLISM OF POLYHYDRIC ALCOHOLS

The compounds of this class that have been studied in connection with the carbon metabolism of fungi include the sugar alcohols glycerol and a few glycols. Mannitol and glycerol are the most important.

Sugar Alcohols Mannitol was first discovered in fungi by Bracon not (44-45) and subsequent work during the nineteenth century revealed its general occurrence in the sporophores of basidiomycetes and ascomycetes (36-59, 264). Other analyses of cellular material show that mannitol occurs in mycelium e.g. of *Aspergillus elegans* (91) and *Rhizopus japonicus* (130) in sclerotia (70) and in spores (223, 242, 265). Mannitol may also appear in the medium and very high yields—from 20 to 50 per cent of the glucose consumed—are found in the culture medium of *Aspergillus* sp. (23) and *Byssosclamyces fulva* (186).

Mannitol appears to function physiologically as a reserve food (161, 165); its general utilizability as a carbon source (Chapter 3) is consistent with this function. It is formed in culture from pentoses, hexoses and glycerol (57, 260). The biosynthesis has not been studied but it is a reasonable hypothesis that mannitol is formed by the reduction of fructose or a fructose phosphate. Extracts of *Stachybotrys atra* carry out the reversible reaction (54)



Provided that a hydrogen donor is available this reaction could synthesize mannitol. Consistent with a reductive origin of mannitol, mannitol formation is increased by restriction of the oxygen supply (23).

D-Sorbitol has been reported formed by *Boletus bovinus* (132) and preformed mats of several basidiomycetes dehydrogenate sorbitol to L-sorbose (239). Other sugar alcohols which are formed by fungi include D-xylitol (35), D-arabitol (77, 182a, 259), D-erythritol (168, 220, 262, 265). Some of the sugar alcohols account for an appreciable fraction of the carbon metabolized; thus D-threitol (D-erythritol) constitutes 13 per cent of the dry weight of the mycelium of *Armillaria*

- 40 Bourquelot E 1898 *Compt rend (Paris)* 16 101-1047
- 41 Bourquelot E and H Hérissé 1901 *Compt rend Sci (Paris)* 13^o 571-574
- 42 Bourquelot E and H Hérissé 190^o *Compt rend (Paris)* 135 390-401
- 43 Bourquelot E and H Hérissé 190^o *Bull soc mycol France* 21 50-57
- 44 Braconnot H 1811a *Ann chim* 19 65-304
- 45 Braconnot H 1811b *Ann chim* 80 272-9
- 46 Brown W 1915 *Ann Botany* 29 313-318
- 47 Brown W and R S K Wood 1933 *Synposiura Soc Gen Microbiol* 3 36-339
- 47a Burger M and K Beran 1936 *Czechoslov Mikrobiol* 1 6-31
- 48 Buxton H W and A Jabbar 1934 *Biochim et Biophys Acta* 15 541-548
- 49 Caldwell M L R M Chester A H Doebbeling and G Volz 1941 *J Biol Chem* 161 361-363
- 49a Cantino E C J Lovett and F A Horestein 1937 *Am J Botany* 41 498-500
- 50 Chattaway F W C C Thompson and A J E Barlow 1934 *Biochim et Biophys Acta* 14 583-581
- 51 Chona B L 1931 *Ann Botany* 46 1033-1040
- 52 Chou T C and M Soodak *J Biol Chem* 196 103-109
- 53 Chrzyszcz T and D Tiukow 1939 *Biochem Z* 97 39-50
- 54 Cochrane V W and M Albrecht Unpublished data
- 55 Contardi A and A Ercoli 1933 *Biochem Z* 61 75-300
- 56 Corbaz R et al 1935 *Helv Chim Acta* 38 935-942
- 57 Coyne F I and H Raistrick 1931 *Biochem J (London)* 25 1513-151
- 58 Cwither W G and F G Lennox 1933 *Australian J Biol Sci* 6 410-407
- 59 Crapek F 1931 *Biochemie der Pflanzen* 3rd ed Vol 1 Jena Gustav Fischer pp 878
- 60 Davis D P E Waggoner and A E Dimond 1933 *Nature (London)* 132 9-9
- 61 Davison F R and J J Willaman 1937 *Botan Gaz* 83 359-361
- 62 Dennis D J A Rothstein and R Meier 1934 *Arch Biochem Biophys* 48 55-60
- 63 Dingle J W W Reid and G L Solomons 1933 *J Sci Food Agr* 4 149-1
- 64 Dingle J and C L Solomons 1930 *J Appl Chem (London)* 9 395-399
- 65 Dox A W and R E Neidig 1910 *Biochem Z* 46 397-400
- 66 Edelman J 1934 *Biochem J (London)* 57 2-33
- 67 Ehrlich F and A Kosmahly 1939 *Biochem Z* 210 16-239
- 68 Eland M and K Myrback 1949 *Arch Biochem* 21 949-950
- 69 Erb N M R T Wisthoff and W L Jacobs 1918 *J Bacteriol* 55 813-810
- 70 Ergle D R 1948 *Phytopathology* 38 14-151
- 71 Euler H on 1931 *Fermentforschung* 4 240-7
- 72 Fischer E H L Kohler and J Fellg 1931 *Helv Chim Acta* 34 113-1138
- 73 Fische I H and R de Montmilion 1931a *Helv Chim Acta* 34 1981-1991
- 74 Fischer F H and R de Montmilion 1931b *Helv Chim Acta* 31 1991-1999
- 75 Fitch V B and R B Dushman 1941 *J Am Chem Soc* 67 1155-1157
- 76 Fishman W H 1931 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 109-119 New York Academic Press
- 77 Frejaud M 1939 *Compt rend (Paris)* 208 1113-1114
- 78 Freilinger K and T Ploetz 1939 *Z physiol Chem Hoppe Seyler* 29 19-20
- 79 Carren K H 1938 *Phytopathology* 28 839-841
- 80 C Humann F and E B hnt 191a *Helv Chim Acta* 30 1-38
- 81 C Humann F and E B hnt 1917b *Helv Chim Acta* 30 159-1593

BIBLIOGRAPHY

- 1 Albericci V J T P Curtin and D Reilly 1913 *Biochem J (London)* 37 243-246
- 2 Amelung H 1930 *Z physiol Chem Hoppe Seyler's* 187 171-172
- 3 Asai T and H Saito 1932 *J Agr Chem Soc Japan* 26 381-387
- 4 Ashour W E 1931 *Trans Brit Mycol Soc* 37 343-359
- 5 Auhagen L and S Grzycki 1933 *Biochem Z* 265 217-222
- 6 Ayres A et al 1939 *Nature (London)* 140 834-836
- 7 Baba S 1912 *J Agr Chem Soc Japan* 18 818 (*Chem Abstr* 45 302, 1911)
- 8 Baba S 1913 *Proc Imp Acad (Tokyo)* 19 10-76
- 9 Bacon J S D and D J Bell 1933 *J Chem Soc (London)* 1933 958-2,30
- 10 Barker S A E J Bourne and T R Carrington 1934 *J Chem Soc (London)* 1934 2125-2129
- 11 Barker S A E J Bourne C C Hewitt and M Stacey 1935 *J Chem Soc (London)* 1935 3731-3740
- 12 Barker S A E J Bourne and M Stacey 1935 *Chem & Ind* 1935 1 87
- 13 de Bary A 1886 *Botan Ztg* 44 393-401
- 14 Basu S N and D R Whitaker 1933 *Arch Biochem Biophys* 42 12-4
- 15 Bealing I J 1933 *Biochem J (London)* 53 93-101
- 16 Bealing I J and J S D Bacon 1933 *Biochem J (London)* 53 277-282
- 17 Beaven G H and F Brown 1949 *Biochem J (London)* 45 221-224
- 18 Bechamp A 1858 *Compt rend (Paris)* 46 44-47
- 19 Béchamp A 1864 *Compt rend (Paris)* 59 496-500
- 19a Beckman C H 1936 *Phytopathology* 46 603-609
- 20 Bentley H R K G Cunningham and F S Spring 1931 *J Chem Soc (London)* 1931 2301-2302
- 21 Bhargava K S 1913 *J Indian Botan Soc* 29 83-99
- 22 Binkley S B 1955 *Ann Rev Biochem* 24 596-606
- 23 Birkinshaw J H J H V Charles A C Hetherington and H Raristrick 1931 *Phil Trans Roy Soc London Ser B* 220 153-171
- 24 Birkinshaw J H C E Stickings and P Tessier 1948 *Biochem J (London)* 42 329-337
- 25 Blank L M and P J Tallev 1941 *Am J Botany* 28 561-569
- 26 Blumenthal H J S T Horowitz A Hemerline and S Roseman 1955 *Bacteriol Proc (Soc Am Bacteriologists)* 1955 137
- 27 Boas F 1917 *Biochem Z* 78 308-312
- 28 Bois E and J Savary 1915 *Can J Research B* 23 208-213
- 29 Bond C R E C Knight and T K Walker 1937 *Biochem J (London)* 31 1033-1040
- 30 Boothroyd B J A Thorn and R H Haskins 1935 *Can J Biochem and Physiol* 33 289-296
- 31 Bose S R 1939 *Ergeb En ymforsch* 8 266-276
- 32 Bose S R and S N Sarkar 1937 *Proc Roy Soc (London) B* 123 193-213
- 33 Bourquelot E 1883 *Compt rend (Paris)* 97 1329-1324
- 34 Bourquelot E 1889 *Bull soc mycol France* 5 132-163
- 35 Bourquelot E 1890 *Bull soc mycol France* 6 vii-viii
- 36 Bourquelot E 1893a *Bull soc mycol France* 9 51-66
- 37 Bourquelot E 1893b *Compt rend (Paris)* 116 1143-1145
- 38 Bourquelot E 1893c *Compt rend (Paris)* 117 383-386
- 39 Bourquelot F 1891 *Bull soc mycol France* 10 49-51

- K. Myrback (eds) *The Enzymes* Vol 2 () p 1180-1338 New York Academic Press
- 14 Leloir L F 1953 *Advances in Enzymol* 14 193-218
- 15 Leloir L F and E Cabib 1953 *J Am Chem Soc* 75 544-546
- 16 Leloir L F and C E Cardini 1953 *Biochim et Biophys Acta* 12 1-29
- 17 LeMense F H J Corman J M Van Lanen and A F Langlykke 1947 *J Bacteriol* 54 149-159
- 18 Lemieux R U J A Thorn and H F Bauer 1953 *Can J Chem* 31 1054-1059
- 19 Leopold H and M P Starbanow 1943 *Biochem Z* 314 232-249
- 20 Lim H 193 *J Fac Ig Hokkaido In p Univ (Sapporo Japan)* 37 165-99
- 21 Lineweaver H R Jang and E F Jansen 1949 *Arch Biochem* 20 137-159
- 22 Lippmann E von 1919 *Per deut chem Ges* 4 3431-3434
- 23 Luh B S and H J Iaff 191 *Arch Biochem Biophys* 33 12-27
- 24 McCarty M 1953 *J Exp Med* 96 453-458
- 25 McCarty M 1953 *J Exp Med* 96 69-80
- 26 McCulloch R J and I I Kertesz 191 *J Biol Chem* 160 149-154
- 27 McCulloch R J and I I Kertesz 1947 *Arch Biochem* 13 217-29
- 28 Macdonald J A 1931 *Ann Appl Biol* 21 989-310
- 29 Mager J 1947 *Biochem J (London)* 41 603-609
- 30 Mager J and M Aschner 1947 *J Bacteriol* 53 83-9
- 31 Mandels G R 1951 *Am J Botany* 38 13-29
- 32 Mandels G R 1953 *Exp Cell Res* 5 49-
- 33 Mandels G R 1951 *Plant Physiol* 29 18-26
- 34 Matus J 1919 *Per deut chem Ges* 58 319 80
- 35 Mayo J K 1953 *New Phytologist* 52 162-164
- 36 Meeuse B J D 1953 *J Exp Botany* 3 52-8
- 37 Mehrotra B S 1919 *J Indian Bot Soc* 8 108-111
- 38 Mehrotra B S 1953 *Floyd* 15 18-18
- 39 Menon K P V 1931 *Ann Botany* 48 181-210
- 40 Miller G L and R Blum 1956 *J Biol Chem* 218 131-137
- 41 Mwa T C T Cheng M Fujisaki and A Toishi 1957 *Acta Phytotom (Japa)* 10 155-170
- 42 Montgomery H B S 1936 *Ann Appl Biol* 3 46-48
- 43 Murakami S 1950 *Misc Repts Pres ch I t \ t Resou ces (Tokyo)* No 17-18 87-9 (*Chem Abstr* 47 657 1953)
- 44 Myrback K and G Neumuller 1950 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 653-71 New York Academic Press
- 45 Myrback K and E Vasseur 1943 *Z phys i Chem Hoppe Seylers* 277 171-180
- 46 Nakatsu S 1956 *J Biochem (Tokyo)* 43 119-127
- 47 Neuberg C and I Mandl 1950 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 527-550 New York Academic Press
- 48 Neuberg C and C Ostendorf 1930 *Biochem Z* 29 464-466
- 49 Nickerson W J 1947 In W J Nickerson (ed) *Biology of the Pathogenic Fungi* p 157-166 Waltham Mass Chronica Botanica Co
- 50 Nickerson W J 1951 *Trans A S Acad Sci II* 13 140-145
- 51 Nisawa K 1953 *J Biochem (Tokyo)* 4 82-83
- 52 Niwa K 1950 *J Biochem (Tokyo)* 3 301-308
- 53 Niwa K 1951 *J Biochem (Tokyo)* 38 109-114
- 54 Norkrans B 1950a *Physiol Plantarum* 3 12-87
- 55 Norkrans B 1950b *Symbol e Botan Ups lense* 11 (1) 5-16
- 56 Norkrans B and B G Rånby 1953 *Physiol Plantarum* 9 198-211

- 82 Cille pie J M and F F Woods 1953 *Australian J Biol Sci* 6 41-46^o
- 83 Culligan W and E T Reese 1953 *Can J Microbiol* 1 90-107
- 84 Godin P 1953a *Biochim et Biophys Acta* 11 114-118
- 85 Godin P 1953b *Biochim et Biophys Acta* 11 119-121
- 86 Goepfert G J and F F Nord 1942 *Arch Biochem* 1 281-331
- 87 Gothoskar S S R P Scheffer J C Walker and M A Stahlmann 1953 *Phytopathology* 43 553-556
- 88 Gothoskar S S R P Scheffer J C Walker and M A Stahlmann 1953 *Phytopathology* 43 381-387
- 89 Gottschalk A 1950 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 551-580 New York Academic Press
- 90 Grassmann W R Stadler and R Bender 1933 *Ann Chem Liebigs* 300 20-40
- 91 Grégoire P 1949 *Bull soc chim biol* 31 801-803
- 92a Halliwell C 1957 *J Gen Microbiol* 17 166-183
- 92b Harter L L 1951 *J Agr Research* 20 161-186
- 93 Harter L L 1925 *J Agr Res arch* 30 951-969
- 94 Harter L L and J L Weimer 1921a *J Agr Research* 21 609-673
- 95 Harter L L and J L Weimer 1921b *J Agr Research* 22 311-317
- 96 Hawkins L A 1915 *Am J Botany* 2 313 358
- 97 Hawkins L A and R B Harvey 1919 *J Agr Research* 18 275-297
- 98 Hérissey H 1951 *Compt rend (Paris)* 197 166-168
- 99 Hockenfull D J D G C Ashton K H Lantes and B K Whitehead 1954 *Biochem J (London)* 57 93-98
- 100 Hofmann F 1934 *Biochem Z* 23 194-204
- 101 Hofmann L and E Jaisko 1950 *Biochem Z* 23 269-272
- 102 Holden M 1950a *Biochem J (London)* 41 115-120
- 103 Holden M 1950b *Biochem J (London)* 47 426-431
- 104 Howell A Jr and R J Fitzgerald 1953 *J Bacteriol* 66 437-441
- 105 Hull R 1939 *Ann Appl Biol* 26 800-822
- 106 Hultin L and L Nordstrom 1949 *Acta Chem Scand* 3 1405-1417
- 107 Humm H J and K S Shepard 1946 *Duke Univ Marine Sta Bull* 3 76-80
- 108 Inanoff N V 1952 *Biochem Z* 167 455-458
- 109 Jagannathan V and K Singh 1953 *Enzymologia* 16 150-160
- 110 Jensen H L 1930 *Soil Sci* 30 59-77
- 111 Jermyn M A 1952 *Australian J Sci Research Ser B* 5 433-443
- 112 Jermyn M A 1953 *Australian J Biol Sci* 6 17-9
- 113 Jermyn M A and R Thomas 1953 *Australian J Biol Sci* 6 70-76
- 114 Jermyn M A and R C Tomkins 1950 *Biochem J (London)* 41 437-440
- 114a Jeuniaux C 1953 *Compt rend soc Biol* 149 1301-1308
- 115 Kertész Z I 1936 *Frgel Enzymforsch* 3 233-238
- 116 Kertész Z I 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 745-768 New York Academic Press
- 117 Kertész Z I et al 1944 *Chem Eng News* 22 10-10a
- 118 Kertész Z I and R J McCollough 1950 *Advances in Carbohydrate Chem* 5 9-107
- 119 Kooiman P I A Roelofsen and S Sweetser 1954 *Enzymologia* 16 237-246
- 120 Kopeloff N L Kopeloff and C J Welcome 1950 *J Biol Chem* 45 171-187
- 121 Kuhn R 1953 *Physiol Chem Hoppe Seyler* 129 51-53
- 122 Kuhn R and C F von Grundherr 1956 *Ber deut chem Ges B* 3 165-176
- 123 Langlykke A F C V Smythe and D Perlman 1957 In J B Sumner and

- 06 Schmidt C. F. Jr 1935 *J Biol Chem* 110 511-50
- 07 Schmidt D 1935 *Biochem Z* 158 273-275
- 08 Schubert E 1939 *Biochem Z* 323 18-33
- 09 Schubert E 1934 *Helv Chim Acta* 17 691-700
- 210 Shu P and A. C. Blackwood 1931 *Can J Botany* 9 113-114
- 11 Sideris C. P. 1934 *Phytopathology* 14 481-489
- 12 Simpson F. J. 1935 *Can J Microbiol* 1 131-139
- 213 Simpson F. J. and E. McCoy 1935 *Appl Microbiol* 1 278-286
- 214 Sir R. G. H. 1931 *Microbial Decomposition of Cellulose* New York Reinhold Publ. Corp. pp. 531
- 15 Smith P. F. and D. Hendlin 1935 *J Bacteriol* 63 440-445
- 216 Smithies W. R. 1935 *Biochem J (London)* 53 316-320
- 17 Sørensen H. 1939 *Physiol Plantarum* 5 183-193
- 218 Sørensen H. 1935 *Nature (London)* 137 303-306
- 219 Stanier R. Y. 1917 *J Bacteriol* 41 555-560
- 220 Stodola F. H. 1946 *J Biol Chem* 166 79
- 221 Strauss B. S. 1935 *Arch Biochem Biophys* 55 77-94
- 222 Stuart N. W. and S. L. Emsweller 1913 *Science* 38 569-570
- 223 Sumi M. 1938 *Biochem Z* 193 161-174
- 224 Sussman M. S. Spiegelman and J. M. Renner 1947 *J Cellular Comp Physiol* 29 149-158
- 225 Takahashi T. and T. Asai 1935 *J Agr Chem Soc Japan* 8 62-63
(*Chem Abstr* 6 560^a 1932)
- 226 Takano K. and T. Miwa 1950 *J Biochem (Tokyo)* 37 435-444
- 227 Takata R. 1939 *J Soc Chem Ind Japan* 32 (Suppl. binding) 243-247
(*Chem Abstr* 4 2206 1930)
- 228 Tanret C. 1903 *Bull soc chim Sér* 29 888-896
- 229 Tate P. 1939 *Parasitology* 31 31-54
- 30 Thaysen A. C. and L. D. Galloway 1930 *The Microbiology of Starch and Sugars* London Oxford Univ. Press pp. 536
- 30a Thomas R. 1936 *Australian J Biol Sc* 9 159-183
- 231 Thornberry H. H. 1938 *Phytopathology* 28 20^a 203
- 32 Tracey M. V. 1933 *Biochem Soc Symposia (Cambridge Engl)* 11 49-61
- 233 Tracey M. V. 1935 *Biochem J (London)* 61 579-586
- 234 Tsuchiya H. M. A. Jeanes H. M. Bricker and C. Wilham 1935 *J Bacteriol* 64 513-519
- 235 Underkofler L. A. and D. K. Roy 1931 *Cereal Chem* 8 18-9
- 236 Varma T. N. R. and K. S. Srinivasan 1934 *Eynologia* 17 116-117
- 37 Vasudeva R. S. 1930 *Ann Botany* 41 469-493
- 238 Vebel S. 1930 In J. B. Sumner and K. Myrback (eds) *The Enzymes* Vol 1 (I) p. 593-670 New York Academic Press
- 39 Vitace J. C. E. S. Pallares and F. F. Nord 1916 *Arch Biochem* 9 439-449
- 40 Von W. and C. Butter 1938 *Ann Chem Liebigs* 554 161-183
- 41 Waggoner P. E. and A. E. Diamond 1935 *Phytopathology* 45 79-87
- 42 Wain R. L. and E. H. Wilkinson 1946 *Ann Appl Biol* 33 401-403
- 43 Waksman S. A. and M. C. Allen 1933 *J Am Chem Soc* 55 3108-3118
- 244 Wallenfels K. 1931 *Naturwiss* 8 306-307
- 4 Wallenfels K. and E. Berni 1937 *Angew Chem* 61 8-29
- 246 Walz A. 1931 *J Am Chem Soc* 53 3726
- 47 Ward H. M. 1888 *Ann Botany* 2 319-33
- 48 Weinland H. and K. Nüchterlein 1934 *Z physiol Chem Hoppe Seyler* 98 48-54
- 49 Welsch M. 1942 *J Bacteriol* 44 571-588
- 50 Whistler R. L. and C. L. Smart 1953 *J Am Chem Soc* 75 1916-1918
- 51 Whitaker D. R. 1953 *Arch Biochem Biophys* 43 23-28

- 161 Obaton F 1929 *Peu gér botan* 41 282-292 363-387 424-440 493-512 679-633
- 162 Obaton F 1932 *Compt rend (Paris)* 194 302-304
- 163 O'Dwyer M H 1939 *Biochem J (London)* 33 713-717
- 164 O'Dwyer M H 1940 *Biochem J (London)* 34 149-152
- 165 Oxford A E and H Raistrick 1935 *Biochem J (London)* 29 1599-1601
- 166 Ian S C L W Nicholson and P Kolachov 1953 *Arch Biochem Biophys* 49 406-420 491-431
- 167 Pazur J H 1952 *J Biol Chem* 199 217-226
- 168 Pazur J H 1954 *Biochim et Biophys Acta* 13 158-159
- 169 Pazur J H 1955 *J Phys Chem* 216 531-538
- 170 Pazur J H and D French 1951 *J Am Chem Soc* 73 3330
- 171 Pazur J H and D French 1952 *J Phys Chem* 196 265-272
- 172 Ierlman D and A F Langlykke 1948 *J Am Chem Soc* 70 3968
- 173 Phaff H J 1941 *Arch Biochem* 13 67-81
- 174 Phillips L L and M I Caldwell 1951a *J Am Chem Soc* 73 3559-3563
- 175 Phillips L L and M I Caldwell 1951b *J Am Chem Soc* 73 3 63-6568
- 176 Pigman W W 1943 *J Research Natl Bur Standards* 46 159-175
- 177 Pigman W W 1951 In J B Sumner and L Myrback (eds) *The Enzymes* Vol 1 (1) p 79-144 New York Academic Press
- 178 Floetz T 1939 *Z physiol Chem Hoppe Seyler's* 61 163-168
- 179 Lottevin H 1903 *Ann inst Pasteur* 17 31-51
- 180a Prentice N and L S Cuendet 1951 *Nature (London)* 174 1151
- 181 Pringsheim H 1912 *Z physiol Chem Hoppe Seyler's* 78 266-291
- 182 Pringsheim H and G Kohn 1924 *Z physiol Chem Hoppe Seyler's* 153 80-96
- 183 Radha K 1953 *Pro Indian Acad Sci* 38B 231-234
- 184a Ragheb H S and F W Fabian 1955 *Food Research* 20 614-625
- 185 Raistrick H and C Smith 1933 *Biochem J (London)* 27 1814-1819
- 186 Ramachandran K and J A Walker 1951 *Arch Biochem Biophys* 31 274-233
- 187 Ramachandran K and J A Walker 1952 *Arch Biochem Biophys* 35 193-203
- 188 Ratajak F J and H S Owens 1942 *Potan Co* 104 390-337
- 189 Reese E T W Gilligan and B Norkran 1952 *Physiol Plantarum* 5 319-390
- 190 Reese E T and H S Levinson 1952 *Physiol Plantarum* 5 343-366
- 191 Reid W W 1952 *Biochem J (London)* 50 287-292
- 192 Reuter C 1912 *Z physiol Chem Hoppe Seyler's* 78 167-145
- 193 Reynolds D M 1954 *J Gen Microbiol* 11 150-159
- 194 Robor F R W Barratt and E L Tatum 1952 *J Biol Chem* 195 433 471
- 195 Roche J 1950 In J B Sumner and L Myrback (eds) *The Enzymes* Vol 1 (1) p 474-510 New York Academic Press
- 196 Roelofsen P A 1953 *Biochim et Biophys Acta* 10 410-413
- 197 Sadashige Y 1950 *Arch Biochem* 28 100-110
- 198 Sadasivan A 1951 *Arch Biochem Biophys* 37 172-185
- 199 Saito T 1906-1907 *J Phys Chem* 2 251-265
- 200 Saksema R K and S K Bose 1944 *J Indian Botanical Soc* 23 108-112
- 201 Saroja K R Venkataraman and I V Giff 1955 *Biochem J (London)* 60 399-403
- 202 Satina S and A F Blakeslee 1928 *Proc Natl Acad Sci U S A* 14 29-35
- 203 Saunders P R R G H S U and R N Cenes 1948 *J Bio Chem* 174 697-703
- 204 Scheffer R P S S Cothoskar C F Pierson and R P Collins 1956 *Physio pathology* 40 63-81

6 Carbon Metabolism

Part III

Organic Acids Lipids and Other Compounds

The final group of carbon metabolites of the fungi has as its most important members the organic acids and the lipids. Brief consideration must also be given to other carbon compounds of various types. Many of the miscellaneous compounds included here have not been studied from a physiological point of view. Nevertheless they are involved in metabolism and therefore in physiology they will not remain indefinitely as biochemical curiosities. Space does not permit an exhaustive listing of the aromatic and other compounds known to be produced by fungi; many more will be found in the reviews cited later.

1 THE METABOLISM OF ALIPHATIC ORGANIC ACIDS

The fungi are perhaps best known in microbiology for their tolerance of acidity—although this is much less general than usually thought—and for the ability of some to accumulate non-volatile organic acids. This section considers the metabolism of the organic acids exclusive of the higher fatty acids, the keto acids of the respiratory pathways, and those acids which are basically ring compounds.

Sugar Acids Acids derived by direct oxidation of either or both terminal carbon atoms of a sugar may be grouped together as the sugar acids. Of these the only one to be formed in quantity is gluconic acid. The reaction was discovered by Mollard (379) in *Aspergillus niger* and subsequent work has shown that at least small amounts of gluconate are formed by a variety of fungi (Table 1). So

- 252 Whitaker D R 1954 *Arch Biochem Biophys* 53 439-449
- 253 Whitaker D R J R Colvin and W H Cook 1954 *Arch Biochem Biophys* 49 257-262
- 254 Wiggers H A L 1852 *Ann Pharm* 1 129-182
- 255 Willaman J J 1920 *Botan Ga* 70 221-223
- 256 Willstaedt H and M Borggard 1916 *Arkiv Kemi Mineral Geol* 23B (3) 1-8
- 257 Winstead N N and J C Walker 1954 *Phytopathology* 44 153 158
- 258 Wolf F A 1914 *Mycol Centr* 4 278-78
- 259 Yabuta T and Y Sumiki 1933 *J Agr Chem Soc Japan* 9 492-497 (*Chem Abstr* 27 4236 1933)
- 260 Yamasaki I and M Simomura 1937 *Biochem Z* 291 310-318
- 261 Young V H 1918 *Plant World* 21 75-87 114-133
- 262 Yuill J L 1948 *Nature (London)* 162 652
- 263 Zeller S M 1916 *Ann Missouri Botan Garden* 3 433-509
- 264 Zellner J 1907 *Chemie der höheren Pilze* Leipzig W Engelmann pp 257
- 265 Zellner J 1910 *Monatsh Chem* 31 617-631

Table 1 (continued)

Acid	Organisms
Citric acid $\begin{array}{c} \text{CH}_2\text{---COOH} \\ \\ \text{HO---C---COOH} \\ \\ \text{CH}_2\text{COOH} \end{array}$	<i>Mucor pyriformis</i> (548) <i>Coniophora cerebella</i> (75) <i>Merulius</i> spp (30) <i>Aspergillus</i> spp (354) <i>Botrytis</i> sp (444a 465) <i>Fusarium graminearum</i> (159) <i>Monilia formosa</i> (458) <i>Penicillium</i> spp (134 136 546 547) <i>Sclerotium delphinii</i> (414)
Fumaric acid $\begin{array}{c} \text{HOOC---CH} \\ \\ \text{CH---COOH} \end{array}$	<i>Circinella</i> sp <i>Cunninghamella</i> sp <i>Mucor</i> sp (207) <i>Rhizopus</i> spp (177 207) <i>Neurospora sitophila</i> (spores) (535) <i>Aspergillus</i> spp (465 496 513 549 550) <i>Caldariomyces fumago</i> (<i>Fumago vagans</i>) (141) <i>Monilia formosa</i> (458) <i>Penicillium griseofulvum</i> (432) <i>Streptomyces coelicolor</i> (145)
Fumaryl DL alanine $\begin{array}{c} \text{HOOC---CH} \\ \\ \text{CH---CO---N---CH} \\ \quad \\ \text{H} \quad \text{COOH} \end{array}$	<i>Penicillium resticulosum</i> (68)
Succinic acid $\text{HOOC---CH}_2\text{---CH}_2\text{---COOH}$	<i>Blastocladiella prinitii</i> (106) <i>Mucor</i> spp (196 502) <i>Rhizopus</i> sp (56) <i>Neurospora sitophila</i> (spores) (535) <i>Fomes annosus</i> <i>Merulius</i> spp (400) <i>Aspergillus terreus</i> (70 433) <i>Botrytis cinerea</i> (224) <i>Clasterosporium</i> sp <i>Fumago vagans</i> (80) <i>Fusarium</i> spp (351) <i>Monilia</i> spp (377 457) <i>Penicillium</i> spp (81 410) <i>Streptomyces coelicolor</i> (145)
Malic acid † $\text{HOOC---CH}_2\text{---CHOH---COOH}$	<i>Rhizopus</i> spp (50 503) <i>Neurospora sitophila</i> (spores) (359) <i>Merulius</i> spp (30) <i>Aspergillus</i> spp (52 70 496 550) <i>Botrytis cinerea</i> (224) <i>Fusarium graminearum</i> (159) <i>Penicillium</i> spp (228 298 465)
Acetic acid CH_3COOH	<i>Mortierella</i> spp (416) <i>Mucor mucedo</i> (415) <i>Neurospora crassa</i> (418) <i>Fomes annosus</i> <i>Merulius</i> spp (402) <i>Polyporus anceps</i> (415) <i>Fusarium</i> spp (225 351) <i>Monilia formosa</i> (457) <i>Sclerotium delphinii</i> (414) <i>Memnoniella echinata</i> (419) <i>Actinomyces</i> spp (291) <i>Streptomyces</i> spp (416 421)

Table 1 The Formation of Organic Acids from Carbohydrates by Fungi and Actinomycetes*

Acid	Organisms
Gluconic acid $\text{HOCH}_2-(\text{CHOH})_4-\text{COOH}$	<i>Mucor plumbeus</i> <i>Rhizopus</i> spp † (465) <i>Ceratostomella ulmi</i> (166) <i>Coniophora cerebella</i> † <i>Merulius lacrymans</i> † <i>Polyporus vaporarius</i> (465) <i>Aspergillus</i> spp (80 192 321 379) <i>Dematium pulchellans</i> (422) <i>Fumigato vagans</i> (80) <i>Fusarium lini</i> (399) <i>Penicillium</i> spp (80 321) <i>Sclerotium delphinii</i> (414)
Mannonic acid $\text{HOCH}_2-(\text{CHOH})_4-\text{COOH}$	<i>Aspergillus niger</i> (321) <i>Penicillium luteo purpurogenum</i> (13)
Galactonic acid $\text{HOCH}_2-(\text{CHOH})_4-\text{COOH}$	<i>Aspergillus niger</i> (321)
Glucuronic acid ‡ $\text{HOOC}-(\text{CHOH})_4-\text{CHO}$	<i>Penicillium</i> spp (298 474) <i>Stereum hirsutum</i> (56) <i>Ustilina vulgaris</i> (567)
Saccharic acid $\text{HOOC}-(\text{CHOH})_4-\text{COOH}$	<i>Aspergillus niger</i> (126)
2 Ketogluconic acid $\text{HOCH}_2-(\text{CHOH})_3-\text{CO}-\text{COOH}$	<i>Penicillium brevicompactum</i> (474)
Xylonic acid ‡ $\text{HOCH}_2-(\text{CHOH})_3-\text{COOH}$	<i>Fusarium lini</i> (265)
Arabonic acid ‡ $\text{HOCH}_2-(\text{CHOH})_3-\text{COOH}$	<i>Fusarium lini</i> (265)
Lactic acid $\text{CH}_3-\text{CHOH}-\text{COOH}$	<i>Allomyces arbuscula</i> (292) <i>Abodachlya</i> sp (23) <i>Blastocladia pringsheimii</i> (106) <i>Chytridium</i> sp <i>Macrochytrium botrydoides</i> (152) <i>Monoblepharella</i> sp (374) <i>Mucor rouxii</i> (132) <i>Mucor</i> spp (502) <i>Pythiogeton</i> sp (107) <i>Rhizidium</i> sp (231) <i>Rhizopus</i> spp (453 541) <i>Sapromyces elongatus</i> (231) <i>Botrytis cinerea</i> (224) <i>Monilia</i> spp (377 457) <i>Actinomyces</i> spp (184 291) <i>Sireptomyces</i> spp (171 424 562)

* Exclusive of the keto acids the higher fatty acids and acidic ring compounds
Organisms are listed by groups phycomycetes ascomycetes basidiomycetes
Fungi Imperfecti actinomycetes in that order

† Identification doubtful

‡ See text

§ In sulfite inhibited culture

Table 1 (continued)

Acid	Organisms
Glutaconic acid ‡ $\text{HOOC}-\text{CH}=\text{CH}-\text{CH}_2-\text{COOH}$	<i>Aspergillus niger</i> (26)
Dimethylpyruvic acid $(\text{CH}_3)_2\text{CH}-\text{CO}-\text{COOH}$	<i>Aspergillus niger</i> § (272 438 439)
Fthylene oxide α β -dicarboxylic acid $\begin{array}{c} \text{O} \\ \diagup \quad \diagdown \\ \text{HOOC}-\text{CH}-\text{CH}-\text{COOH} \end{array}$	<i>Aspergillus fumigatus</i> (67 368) <i>Monilia formosa</i> <i>Penicillium viniferum</i> (458)

far as is now known the accumulation of large amounts of the acid is restricted to certain strains of *Aspergillus* spp and *Penicillium* spp and many fungi yield no gluconate by classical isolation procedures (321 372 465)

Because of the potential industrial importance of the process the conditions of gluconate synthesis by *Aspergillus niger* have been intensively studied. It is found generally that a relatively high pH favors the formation of gluconate over that of citrate (101 221 513). Calcium carbonate is customarily added to the medium to control the pH interference with metabolism by precipitated calcium gluconate can be minimized by the addition of boric acid or borax to the medium (386). As in other acid syntheses but to a more pronounced degree the accumulation of gluconate is favored by aeration (464 554). It is also favored by a relatively high glucose concentration (16 269) and a limiting amount of phosphate (269). Under appropriate conditions almost complete conversion of glucose to gluconate occurs (221).

The biochemical basis of gluconic acid formation is probably the action of enzyme glucose aerodehydrogenase (glucose oxidase) (Chapter 7) although *Dematium pullulans* is reported to form gluconate (422) and to lack the enzyme (212). The first product in the reaction is glucono δ lactone which may be found in the medium with the free acid (80). Gluconate is further utilized by fungi (46 514) but the path of metabolism is not known.

The formation by *Aspergillus niger* of other hexonic acids—mannonic acid from mannose and galactonic acid from galactose—has been reported (Table I) but the process has not been studied further. *Penicillium chrysogenum* forms lactobionic acid from lactose (151a).

Glucuronic acid is reported to be formed in traces by fungi (Table 1) although an unequivocal identification is still lacking. Saccharic acid is found in cultures of *Aspergillus niger* (126 539). The appear

Table 1 (continued)

Acid	Organisms
Ethyl acetate $\text{CH}_3\text{COOCH}_2\text{CH}_3$	<i>Indoconidiophora moniliformis</i> (241) <i>Penicillium digitatum</i> (72)
Oxalic acid $\text{HOOC}-\text{COOH}$	<i>Rhizopus nigricans</i> (545) <i>Neurospora crassa</i> (418) <i>Sclerotinia</i> spp (33 545) <i>Armillaria mellea</i> (261) <i>Coniophora cerebella</i> (75) <i>Corticium centrifugum</i> (277) <i>Fomes annosus</i> (401) <i>Marasmius chordalis</i> (478) <i>Merulius</i> spp (401) <i>Polyporus tumulosus</i> (437) <i>Poria caillantii</i> (40?) <i>Ispergillus</i> spp (67 515 545) <i>Botrytis cinerea</i> (224 545) <i>Dematium pullulans</i> (34) <i>Fusarium</i> spp (159 357) <i>Penicillium</i> spp (134 156 545) <i>Sclerotium delphinii</i> (414) <i>Trichophyton tonsurans</i> (533)
Itaconic acid $\begin{array}{c} \text{COOH} \\ \\ \text{C}=\text{CH}_2 \\ \\ \text{CH}_2 \\ \\ \text{COOH} \end{array}$	<i>Aspergillus itaconicus</i> (314 315) <i>A. terreus</i> (105 349) <i>Ustilago eae</i> (263)
cis Aconitic acid $\begin{array}{c} \text{CH}-\text{COOH} \\ \\ \text{C}-\text{COOH} \\ \\ \text{CH}_2-\text{COOH} \end{array}$	<i>Aspergillus niger</i> (455)
Itatartaric acid $\begin{array}{c} \text{COOH} \\ \\ \text{HO}-\text{C}-\text{CH}_2\text{OH} \\ \\ \text{CH} \\ \\ \text{COOH} \end{array}$	<i>Aspergillus terreus</i> mut (492)
Tartaric acid ‡ $\text{HOOC}-(\text{CHOH})_2-\text{COOH}$	<i>Fusarium</i> spp (357)
Clyoxylic acid $\text{OHC}-\text{COOH}$	<i>Aspergillus niger</i> § (87) <i>Botrytis cinerea</i> (224)
Glutaric acid ‡ $\text{HOOC}-(\text{CH}_2)_3-\text{COOH}$	<i>Aspergillus niger</i> (26)

Citric Acid The ability of fungi to accumulate citric acid was first discovered by Welmer (546-517) in species of *Citromyces* (*Penicillium*). Later (155) it was found that higher yields are obtained from selected strains of the black spored *Aspergilli*, usually referred to as *Aspergillus niger*. Certain of these strains are presumably used for the commercial production of citric acid. Although the acid is probably a metabolic intermediate of aerobic metabolism in all fungi (Chapter 7) it appears in significant amounts only in a few forms (Table 1) primarily species—or more correctly certain strains of species—of *Penicillium* and *Aspergillus*. Most strains of *A. niger* have the potentiality for the accumulation of citric, gluconic and oxalic acids; particular strains under a given set of conditions may form predominantly one acid or a mixture of two or three (48-58). Strain differences contribute to the failure of different laboratories to agree on the details of the citric acid process, but the basic physiological events are presumably the same in all strains. Selection may increase the yield of citrate (169-307) and correspondingly a strain may change its capacity as the result of mutation (218-331). No adequate genetic analysis has been made; probably the typical strain is a heterocaryon with nuclei of different genetic types which segregate in spore formation (28).

The biosynthesis of citrate by *Aspergillus niger* was investigated extensively during the period 1920 to 1940, largely by growth and replacement culture experiments which by themselves do not discriminate among the various possible metabolic pathways. Theories based on this work, all obsolete or correct only by chance, are reviewed by Foster (200) and Walker (538).

As discussed in Chapter 7, it now seems that fungi like other organisms utilize the citric acid cycle in aerobic respiration; the evidence which is admittedly still incomplete comes from enzymatic studies and from isotope experiments with citrate formation in *Aspergillus niger*. At least one other pathway of citrate (or isocitrate) metabolism is known in fungi (405) and may be found in the future to make a significant contribution to citrate synthesis. If we postulate the ubiquity of the citric acid cycle in fungi, accumulation of citrate is best regarded as an unusual modification under genetic and environmental control of a process which occurs in all cells under aerobic conditions.

Figure 1 is a simplified diagram of some of the known and probable relationships of organic acids in *Aspergillus niger* and, with some exceptions, very likely in all fungi. The salient and physiologically significant implications of the scheme may be summarized:

ance of pentonic acids in cultures of *Fusarium lini* (265) has not been confirmed and the method of identification used leaves something to be desired

Lactic Acid The formation of lactic acid in fungi is restricted almost entirely to phycomycetes (Table I) and all fungi which convert a major fraction of the carbohydrate to lactate are in this group. The early history of the problem and the cultural conditions favoring lactic acid accumulation are reviewed by Foster (200). Lockwood et al (353) and Ward et al (541-542).

The optical isomer formed by fungi is usually L-lactic (dextro-rotatory) (536-541) although formation of the D and DL forms has been reported (453-457-501).

As discussed in Chapter 7 the anaerobic formation of lactate by *Rhizopus oryzae* conforms to the postulate that glucose is broken down by the Embden Meyerhof sequence of reactions to yield lactate ethanol and carbon dioxide in equimolar amounts the mechanism by which the equimolar ratio is maintained is not yet known. More lactate is formed aerobically than anaerobically and the biosynthesis of this extra lactate is not understood. If it is assumed that all of the glucose used is converted to pyruvate the aerobically produced lactate must arise from it or from ethanol perhaps with 4 carbon acids as intermediates (109).

Some carbon dioxide enters into the lactate molecule (226). Lactic acid production in high yields is obtained from cultures which are aerated either by shaking (54) or by the rotary drum method (542). It should be noted that Sakaguchi and co-workers (460) could not confirm the stimulation of lactate formation by aerobiosis.

Lactate forming organisms other than *Rhizopus* spp. that is the lower phycomycetes and *Streptomyces* do not so far as is known form extra lactate aerobically. Again in contrast to *Rhizopus* some lower phycomycetes anaerobically convert 85 to 100 per cent of the glucose consumed to lactic acid (106-231-292). Evidently the biochemical basis here differs in important respects from that of *Rhizopus* the difference recalls that between the homofermentative and the heterofermentative lactic acid bacteria.

Lactic acid is not an end product of aerobic metabolism but is consumed after exhaustion of the principal carbon source (106-171). An aerobic lactic dehydrogenase is known in *Penicillium chrysogenum* (111) but the enzymatic basis of lactate utilization in those fungi which accumulate the acid is not known. The oxidation of lactic acid by resting cells of *P. chrysogenum* is suppressed by glucose (278).

particular reactions shown in Figure 1 may not all occur in fungi but there is no doubt that fatty acid and amino acid skeletons are built up from intermediates of carbohydrate oxidation. It is significant that strains of *Aspergillus niger* which do not accumulate citrate have a high growth rate (185) and that highly acidogenic mutants are relatively slow growing (218).

5 Carbon may also be drawn off into gluconic or oxalic acid in *Aspergillus niger* and into still other acids in other fungi.

These biochemical relationships explain many of the known physiological requirements for citric acid accumulation. In the first place the concentration of available carbohydrate must be high (354); usually glucose at 15 to 20 per cent is employed. Under these conditions there is an excess of carbon over that required for energy and for the synthesis of cellular material; this excess will then we think accumulate wherever genetic and environmental factors permit rapid formation but only slow breakdown of a metabolite. Such a metabolite may be directly on a major pathway or on a shunt, diversionary pathway (200).

The requirement for a high carbohydrate concentration has its counterpart in the usual finding that highest yields of citrate are obtained if other major nutrients—nitrogen, phosphorus, magnesium, sulfur—are at levels which limit growth (103, 155, 309, 413). Data which contradict this generalization are however often reported and there are undoubtedly secondary physiological effects of nutrients (472, 473).

Minor elements affect citric acid production but contradictory results have been reported. There is no doubt on the one hand that zinc, iron, manganese and copper are essential to growth and acid formation (518) or on the other that high concentrations of metals may lower yields (420). Between these two extremes it appears from recent work that the best yield of citric acid and other acids is obtained at concentrations of the minor elements which allow some growth but are at the same time limiting to growth (130). Increases of zinc, for example, may result in a greater total yield but the percentage of conversion of glucose to acid is lowered (129). Perhaps the clearest example is the inhibitory effect of iron on citrate accumulation; the inhibition disappears in a medium in which growth is limited by low phosphate (472).

The optimum level of any given nutrient affected as it is by the strain used, the conditions of culture and the other nutrients present is a matter for individual determination. The important general

Oxygen is required for citrate synthesis both in surface culture (169) and in the more recently developed submerged processes (308 472). Too vigorous aeration may it has been suggested (186) sweep out the carbon dioxide rapidly enough to reduce citrate formation. The probably essential role of carbon dioxide mentioned above is supported by experiments showing that increases in the carbon dioxide of the atmosphere increase the yield of citrate (259).

Measurements of the oxidation-reduction potential during the culture cycle suggest that there is an optimum potential for citrate formation above the optimum. *Aspergillus niger* forms gluconic acid rather than citric and below it ethanol is formed (332).

Virtually all studies of citric acid formation indicate that a low culture pH is essential to good yields generally in surface culture an initial pH of about 2.0 is recommended for *Aspergillus niger* 3.0–4.0 for *Penicillium* spp. (169 213). In aerated cultures of *A. niger* a somewhat higher initial pH—about 4.5—appears to be optimal (472). At pH values above the optimum gluconic and oxalic acids tend to replace citric (213 385). The mechanism of this pH effect is unknown; it has a counterpart in the effect of pH on the balance of metabolic products formed by the lactic acid and the coliform bacteria.

Metabolic poisons affect citric acid formation and have been used in attempts to elucidate the biochemistry of acid formation. The results are ambiguous as is to be expected from the multiplicity of reactions which compete with or supply carbon to the citrate pool and from the fact that most poisons have several sites of action. The effect of slightly growth-inhibitory concentrations of the lower aliphatic alcohols on the yield of citric acid is very striking (383 455) and may prove on further investigation to be specific.

Fumaric Acid and Related Acids Four of the 4-carbon dicarboxylic acids—oxalic, citric, malic, fumaric and succinic—may be considered together inasmuch as they are readily interconvertible (see Figure 1). Oxalacetate is unstable and does not accumulate appreciably. Probably all the other acids of this group can be detected by sufficiently sensitive methods in the cells or culture fluid of fungi during aerobic metabolism. For this reason the listing of fungi which produce these acids is somewhat deceptive; it is more a history of the interest of investigators than a survey of acid formation.

Fumaric acid is formed in small amounts by many fungi (Table 1); certain of the Mucorales, especially *Rhizopus* spp., are the only fungi in which it accounts for a major fraction of the glucose consumed. *Aspergillus fumigatus*, as originally isolated, formed fumaric acid in large amounts (549) but the strain on examination ten years




Figure 2 Growth and chemical changes in the course of citric acid synthesis by *Aspergillus niger* in shake culture. Redrawn from P. Shu and M. J. Johnson *Industrial and Engineering Chemistry*, Vol. 40, p. 1201, 1202, July 1948. Copyright 1948 by the American Chemical Society and reprinted by permission of the copyright owner.

principle that growth and acid formation are to a degree competitive should not be obscured by these secondary effects important as they may be for other reasons.

The time course of citric acid formation in submerged aerated culture is illustrated in Figure 2 from which it is evident that there is a phase of growth with little acid production followed by a phase of acid formation and slow growth. Maximum yields obviously require that a balance be struck between growth and acid formation; at some point there is enough cell material to convert the residual glucose with maximum efficiency into product. These considerations apply in principle to all acid syntheses and in fact to all syntheses of non-protoplasmic materials.


Reference to Figure 1 makes it clear that many different carbon compounds entering the metabolic pathway at different points may be expected to provide carbon for citrate synthesis and this proves to be true (354). There are so many possibilities here that experiments in which a carbon compound is added to a preformed mycelium in an attempt to trace biochemical pathways generally fail—almost any compound can be fed into the citric acid pool.

berg Wieland reaction This pathway has been proposed in fungi on the basis of both chemical and isotope data (30 49 102 202 203 450) As noted elsewhere (Chapter 7) the evidence is not conclusive although there is of course some mechanism for the conversion of acetate and other 2 carbon compounds to larger molecules


4 The cleavage of isocitric acid directly to succinic and glyoxylic acids (405) This system is probably identical with one known in bacteria (477) its functions during glucose metabolism and its quantitative importance in fungi are not yet known

Formic and Acetic Acids These two volatile acids are more characteristically bacterial than fungal metabolites and under normal conditions of culture neither accumulates in such large amounts as do the non volatile acids so far considered

Formic acid has not been reported to accumulate during carbohydrate utilization but several fungi form it during the metabolism of lactic and pyruvic acids (133 454) These experiments suggest the well known transacetylase reaction



The entrance of formate or a derivative into metabolism is demonstrated by isotope results (302) and by direct utilization of formate (137 262) It has been suggested that formate may enter through the sequence of reactions



Until active cell free preparations are studied these reactions of formate must remain hypothetical in the fungi

Acetic acid is found in traces in the medium of many fungi and actinomycetes the list in Table I omits non specific determinations of volatile acids (54 73) Larger amounts of acetate are formed by *Streptomyces fradiae* (421) and by *Fusarium* spp (351) As also shown in Table 1 two fungi form ethyl acetate from carbohydrate

In view of the central role of acetate and acetyl coenzyme A in metabolism we may assume their presence even if isolable amounts are not formed this generalization is borne out by isotope dilution studies on *Penicillium chrysogenum* (368) and *Streptomyces coelicolor* (144) In particular organisms acetate may be formed from pentose (225) from ethanol by oxidation (398 415) or possibly from glycine by deamination (562) Direct studies have shown that exogenous acetate is oxidized by *Merulius niveus* (30) *Penicillium chrysogenum* (278) and *Zygorhynchus moelleri* (382)

later proved to have lost the capacity to form the acid from glucose except under special circumstances (513-550). Fumarate is also found in the sporophores of basidiomycetes (57^r-576).

Penicillium resticulosum in glucose medium forms fumaric-alanine in which the amino group of alanine is joined in a peptide linkage with one carboxyl of fumarate (68).

Succinic acid accumulation is rarely so great as that of the other acids considered thus far. Resting cells of *Blastocladia pringsheimii* accumulate succinate equivalent to about 10 per cent of the glucose metabolized (106) but the amounts formed by most organisms listed in Table 1 are much smaller. Malic acid has only rarely (56-455) been reported to be formed in large amounts from glucose although it is produced by a variety of saprophytic fungi (Table 1). Most of the identifications of malic acid it should be noted rest on color tests the use of which is often difficult and uncertain. re-examination by enzymatic methods would be profitable.

In fumaric acid synthesis one of the most striking physiological aspects is the effect of zinc which increases growth and glucose consumption and decreases by as much as 80 per cent the formation of fumarate (208). Fumaric acid formation is thus a symptom of retarded growth in the presence of an abundance of carbohydrate. By analogy with acid formation in *Aspergillus niger* it is likely that any metal deficiency has the same effect (129-130) but the role of other metals is still uncertain and iron above its optimum level seems to have little influence (208).

Carbon dioxide has been shown by isotope studies to enter into the molecule of fumaric acid (204-205) and succinic acid (141). Succinate formation by *Streptomyces coelicolor* and *Blastocladia pringsheimii* is enhanced by carbon dioxide (106-114) but the relation of exogenous carbon dioxide to fumarate formation is uncertain (206-207).

The 4 carbon dicarboxylic acids under discussion probably are formed by more than one biochemical mechanism. Four such mechanisms have been suggested namely

1. Oxidative formation in the citric acid cycle. Bulk formation by this mechanism has not been proved.

2. The reductive carboxylation of pyruvic acid either to oxalacetate or directly by the agency of the malic enzyme to malic acid. This is the best known and perhaps the major pathway. Evidence for its role comes from studies on the incorporation of isotopic carbon dioxide and on the physiological effects of carbon dioxide mentioned above.

3. The direct condensation of two 2 carbon fragments in the Thun

35° 384) Distinctive features include the very low pH optimum—about 1.9 for shaken cultures—and the pronounced effects of sodium chloride and magnesium sulfate under certain conditions. The low pH optimum of the processes has been shown conclusively (334) to be determined by the fact that the enzyme system for itaconate formation is present only during cultivation at low pH.


Itaconic acid is formed *in vitro* from *cis*-aconitic acid by the enzyme *cis*-aconitic decarboxylase (11) since *cis*-aconitate is a component of the tricarboxylic acid cycle, the formation of itaconic acid may be regarded as an alternate metabolic pathway for glucose carbon (Figure 1).

A mutant of *Aspergillus terreus* obtained by ultraviolet irradiation forms a mixture of itatartaric acid and its lactone (492). The structure (Table 1) has an obvious resemblance to that of itaconate and it is reasonable to suppose that the two acids are biochemically related.

Other Acids Tartaric acid has been reported in small amounts in cultures of *Fusarium* spp. (Table 1) but the criteria of identification were not wholly satisfactory. Using color tests and copper reduction as criteria Bernhauer and Bockl (51) found that most strains of *Aspergillus niger* form tartrate during incubation of preformed mycelium with ethanol. The problem merits study with the more sensitive methods now available for isolation and determination of acids.

Glyoxylic acid ($\text{OHC}-\text{COOH}$) often accompanied by glycolic acid ($\text{HOCH}_2-\text{COOH}$) is formed in traces during carbohydrate breakdown by some fungi (Table 1) it has also been reported as being formed during the metabolism of acetate (55, 125, 403), ethanol (136) and citrate (125). It should be noted that substances giving the color reactions for glyoxylic acid are found in sterile autoclaved culture media (157) it has not always been clear that adequate controls have been included in experiments on the occurrence of glyoxylate.

Glyoxylate has been proposed as an intermediate in the oxidative formation of oxalate from acetate by the following reactions (403)



It is not yet certain that these reactions occur. The fact that glyoxylic and glycolic acids replace glycine in the nutrition of a mutant of *Neurospora crassa* (266) suggests a second pathway for their utilization and formation. Finally the ability of fungi under some conditions to split isocitrate to succinate and glyoxylate and to oxidize glycolate to glyoxylate has been shown unequivocally (405). Glyoxylate is also it should be noted a product of the action of allantoinase (Chapter 8).

The higher fatty acids above acetic are considered integrally with the metabolism of lipids

Oxalic Acid Oxalic acid is a common metabolic product of fungi (Table 1) especially of the Agaricales. Species of *Penicillium* and *Aspergillus* accumulate the largest amounts but substantial oxalate production occurs in a few other fungi e.g. *Sclerotium delphinii* (414). Sporophores of basidiomycetes often contain oxalate (17 575 576).

The physiology of oxalate formation deserves more study. The acid is formed from various carbon sources (49) the most favorable cultural conditions include a high carbohydrate concentration adequate aeration (49 545) a limited supply of inorganic nutrients (129 447) and most important a relatively high initial pH (46 49 299). The enzymatic basis of the pH requirement has been elucidated by Shimazono (470) at low pH an enzyme oxalic acid decarboxylase is formed and converts oxalate to formate and carbon dioxide. This finding suggests for the higher basidiomycetes at least that oxalate is a normal metabolite and accumulates under conditions of an enzyme deficiency. Cultures of *Aspergillus niger* utilize oxalate provided that the pH is below 5.5 (27).

In *Corticium centrifugum* oxalic acid production is increased by a growth limiting deficiency of thiamine (391). This relation is consistent with the general view that oxalic and other acids are formed maximally when available carbon cannot be used for growth.

The biochemical origin of oxalic acid has been investigated rather frequently but none of the pathways proposed has been proved (4 5 403). Recent results with isotopic carbon are consistent with the origin of oxalate from oxalacetate (88 138a 201 338). The possibility that glycine which in different organisms is metabolically related to other 2-carbon compounds is involved has not yet been explored. Progress on the problem depends on the preparation of active non-cellular preparations.

Itaconic and Itatartaric Acids Itaconic acid was discovered by Kinoshita (314 315) to be formed by a strain which he designated *Aspergillus itaconicus* n. sp. This strain has lost its capacity to form more than small amounts of the acid but strains of *Aspergillus terreus* have been found to produce large yields (105 384). Other strains of the same species form no itaconate and ultraviolet induced mutants yield more or less acid than the parent culture (319).

The optimum conditions for itaconic acid formation are rather different from those which give best growth of the organism (318 350).

352-384) Distinctive features include the very low pH optimum—about 1.9 for shaken cultures—and the pronounced effects of sodium chloride and magnesium sulfate under certain conditions. The low pH optimum of the processes has been shown conclusively (334) to be determined by the fact that the enzyme system for itaconate formation is present only during cultivation at low pH.


Itaconic acid is formed *in vitro* from *cis*-aconitic acid by the enzyme *cis*-aconitic decarboxylase (41) since *cis*-aconitate is a component of the tricarboxylic acid cycle; the formation of itaconic acid may be regarded as an alternate metabolic pathway for glucose carbon (Figure 1).

A mutant of *Aspergillus terreus* obtained by ultraviolet irradiation forms a mixture of itatartaric acid and its lactone (492). The structure (Table I) has an obvious resemblance to that of itaconate and it is reasonable to suppose that the two acids are biochemically related.

Other Acids Tartaric acid has been reported in small amounts in cultures of *Fusarium* spp. (Table I) but the criteria of identification were not wholly satisfactory. Using color tests and copper reduction as criteria, Bernhauer and Bockl (51) found that most strains of *Aspergillus niger* form tartrate during incubation of preformed mycelium with ethanol. The problem merits study with the more sensitive methods now available for isolation and determination of acids.

Glyoxylic acid ($\text{OHC}-\text{COOH}$) often accompanied by glycolic acid ($\text{HOCH}_2-\text{COOH}$) is formed in traces during carbohydrate breakdown by some fungi (Table I); it has also been reported as being formed during the metabolism of acetate (55, 125, 403), ethanol (136), and citrate (125). It should be noted that substances giving the color reactions for glyoxylic acid are found in sterile autoclaved culture media (157); it has not always been clear that adequate controls have been included in experiments on the occurrence of glyoxylate.

Glyoxylate has been proposed as an intermediate in the oxidative formation of oxalate from acetate by the following reactions (403):



It is not yet certain that these reactions occur. The fact that glyoxylic and glycolic acids replace glycine in the nutrition of a mutant of *Neurospora crassa* (566) suggests a second pathway for their utilization and formation. Finally, the ability of fungi under some conditions to split isocitrate to succinate and glyoxylate and to oxidize glycolate to glyoxylate has been shown unequivocally (405). Glyoxylate is also, it should be noted, a product of the action of allantoinase (Chapter 8).

The other acids listed in Table 1—glutaric, glutamic, dimethylpyruvic and ethylene oxide $\alpha\beta$ -dicarboxylic acids—have as yet no established physiological or biochemical significance. The first two have been reported only once from xylose cultures if their formation can be confirmed. A route from pentose to α -ketoglutaric acid is a possibility. Dimethylpyruvic acid accumulates in cultures poisoned with bisulfite. Suggestions on the biochemical origin of dimethylpyruvic acid are reviewed by Birkinshaw (63).

Ethylene oxide $\alpha\beta$ -dicarboxylic acid (*trans*-1-epoxysuccinic acid), chemically related to tartaric acid, is formed from a variety of carbon sources by *Aspergillus fumigatus* in up to 21 per cent yield (370).

During the breakdown of citric acid by *Aspergillus niger* two other acids, acetonedicarboxylic acid ($\text{HOOC}-\text{CH}_2-\text{CO}-\text{CH}_2-\text{COOH}$) and malonic acid ($\text{HOOC}-\text{CH}_2-\text{COOH}$) appear in the medium and it has been suggested that they derive from a route of citrate dissimilation otherwise unknown (125-495).

Conclusions Some generalizations can be made concerning the physiology and biochemistry of those acids—gluconic, citric, fumaric, succinic, lactic and oxalic—which are formed in large amounts and which have been studied intensively. Not all the data agree but in general it can be said:

1. The acids occur in small amounts at least as normal metabolites in organisms other than the fungi. This is also true of acetate, formate, glyoxylate and malate.

2. With few exceptions—and these are subject to confirmation—the acids which accumulate in fungus cultures are later utilized for energy and growth unless removed by a trapping agent, e.g. calcium ion, or unless their accumulation poisons the cell.

3. Most experiments agree that highest acid yields in terms of the fraction of sugar converted to product are obtained in media in which carbohydrate is abundant but in which some other factor, usually nutritional, restricts growth to a less than maximum amount.

4. Oxygen is essential for the synthesis of all but lactic acid; this requirement is of course obvious from the structure of the acids. Gluconate formation, catalyzed as it probably is by a flavin enzyme, is the most responsive to oxygen pressure.

5. Carbon dioxide enters by fixation into lactic, citric, fumaric and succinic acids and appears to be quantitatively important in providing the carbon of all of these except possibly lactate.

6. Genetic factors determine whether or not a clonal culture will accumulate acid under given external conditions.

We may visualize a metabolic pool of any given acid with carbon flowing into it from a high level of carbohydrate and flowing out into amino acids into respiratory carbon dioxide and into other undetermined channels. Under conditions of minimal demand on the pool for growth the accumulation of acid is determined by the available carbohydrate by the relative rates of the many reactions which produce and break down the acid and by the rate of exchange of acid between cell and environment. The scheme of Figure 1 applying specifically to citric acid is in principle a model for all acid accumulation. An acid may be more closely related to carbohydrate as is gluconic acid or it may be in a metabolic shunt removed by one or more enzymatic steps from the main pathway of carbohydrate dissimilation.

Physiologically a utilizable acid serves as a reserve carbon source although the concept is more or less meaningless under the usual conditions of acid accumulation in culture. A priori it would appear that the soluble acids having effects on both the pH and the osmotic concentration of the medium are less satisfactory reserve substances than the insoluble polysaccharides and fats.

The permeability of the cell membrane to organic acids deserves more study difficult as the problem is. Obviously a biochemical equilibrium will be reached very quickly within the cell and very little acid will accumulate if the formed acid cannot leave the cell. If it leaves rapidly in fact the concept of enzymatic equilibrium becomes almost inapplicable. Conversely if re entrance of external acid into the cell is slow its utilization as a source of carbon will be limited. It is conceivable that some of the environmental factors which favor acid accumulation act primarily on the rate of exchange between cell and environment and that differences between strains reflect membrane differences rather than enzymatic capabilities.

The genetic factor is so far a matter of observation only—acid formation is highly specific and individual strains differ from each other both quantitatively and qualitatively. These strain differences must not however be overemphasized so far only one instance of complete loss of ability to form an acid has been reported that of *Aspergillus fumigatus* (p. 141) and even the altered strain is able under some conditions to form the missing product. Quantitative changes are frequent especially when several acids compete for carbon. The qualitative constancy is particularly remarkable in view of the fact that in many of the fungi studied for acid production the conidia used for transfer of cultures represent isolations of nuclei from a heterocaryotic mycelium.

Virtually all modern theories of acid biosynthesis assume a preliminary conversion of carbohydrate to pyruvic acid except of course in the biosynthesis of the sugar acids. Whether this assumption represents the facts or merely the limitations of current biochemical knowledge is uncertain. The phosphogluconate pathway of glucose metabolism (Chapter 7) provides to the cell a group of compounds including the 4- and 7-carbon sugar phosphates previously unknown and quite possibly of direct importance in acid synthesis.

Table 1 summarizes the reported formation of acids by fungi and actinomycetes acting in culture on carbohydrates. So far as possible duplicating reports have been omitted and some of those included are based on questionable or at least inadequate evidence. Satisfactory proof for the identity of an acid should involve more than color tests or the chromatographic constant in a single solvent; on the other hand the results of these presumptive tests are often correct and reports based on them cannot therefore be dismissed out of hand. Ideally an acid or other metabolic product should be isolated as a crystalline derivative; if this is not possible several independent criteria should be utilized. Enzymatic identification promises to be the most sensitive and accurate for metabolites which are found only in micro amounts.

2 THE METABOLISM OF LIPIDS

The lipids may be defined as those substances which are soluble in fat solvents; a few like the lecithins are somewhat water soluble but most are insoluble in water. Lipids which are of importance to the physiology of the fungi include the true fats, the fatty acids, the compound lipids (phospholipids), the sterols, the carotenoids and triterpenes and a few hydrocarbons. The general chemistry of the lipids is reviewed by Deuel (164). Those lipids which occur as constituents of the cell are considered in Chapter 2; here we are primarily concerned with synthesis and breakdown although for convenience the occurrence of some lipids is included.

The Synthesis of Fats, Fatty Acids and Phospholipids The neutral fats are triglycerides of the higher fatty acids. These are synthesized by many fungi in amounts up to 50 per cent of the mycelial dry weight. One rather striking difference between the lipids of fungi and those of animals is the high content of free fatty acids in most fungal lipids (Chapter 2). This means that most experiments on the factors affecting gross lipid content deal indiscriminately with both neutral fats and fatty acids.

Virtually all modern theories of acid biosynthesis assume a preliminary conversion of carbohydrate to pyruvic acid except of course in the biosynthesis of the sugar acids. Whether this assumption represents the facts or merely the limitations of current biochemical knowledge is uncertain. The phosphogluconate pathway of glucose metabolism (Chapter 7) provides to the cell a group of compounds including the 4- and 7-carbon sugar phosphates previously unknown and quite possibly of direct importance in acid synthesis.

Table 1 summarizes the reported formation of acids by fungi and actinomycetes acting in culture on carbohydrates. So far as possible duplicating reports have been omitted and some of those included are based on questionable or at least inadequate evidence. Satisfactory *proof for the identity of an acid should involve more than color tests or the chromatographic constant in a single solvent*; on the other hand the results of these presumptive tests are often correct and reports based on them cannot therefore be dismissed out of hand. Ideally an acid or other metabolic product should be isolated as a crystalline derivative; if this is not possible several independent criteria should be utilized. Enzymatic identification promises to be the most sensitive and accurate for metabolites which are found only in micro amounts.

2 THE METABOLISM OF LIPIDS

The lipids may be defined as those substances which are soluble in fat solvents; a few like the lecithins are somewhat water soluble but most are insoluble in water. Lipids which are of importance to the physiology of the fungi include the true fats, the fatty acids, the compound lipids (phospholipids), the sterols, the carotenoids and triterpenes and a few hydrocarbons. The general chemistry of the lipids is reviewed by Deuel (164). Those lipids which occur as constituents of the cell are considered in Chapter 2; here we are primarily concerned with synthesis and breakdown although for convenience the occurrence of some lipids is included.

The Synthesis of Fats, Fatty Acids and Phospholipids The neutral fats are triglycerides of the higher fatty acids. These are synthesized by many fungi in amounts up to 50 per cent of the mycelial dry weight. One rather striking difference between the lipids of fungi and those of animals is the high content of free fatty acids in most fungal lipids (Chapter 2). This means that most experiments on the factors affecting gross lipid content deal indiscriminately with both neutral fats and fatty acids.

Virtually all modern theories of acid biosynthesis assume a preliminary conversion of carbohydrate to pyruvic acid except of course in the biosynthesis of the sugar acids. Whether this assumption represents the facts or merely the limitations of current biochemical knowledge is uncertain. The *phosphogluconate pathway of glucose metabolism* (Chapter 7) provides to the cell a group of compounds including the 4 and 7 carbon sugar phosphates previously unknown and quite possibly of direct importance in acid synthesis.

Table 1 summarizes the reported formation of acids by fungi and actinomycetes acting in culture on carbohydrates. So far as possible duplicating reports have been omitted and some of those included are based on questionable or at least inadequate evidence. Satisfactory proof for the identity of an acid should involve more than color tests or the chromatographic constant in a single solvent; on the other hand the results of these presumptive tests are often correct and reports based on them cannot therefore be dismissed out of hand. Ideally an acid or other metabolic product should be isolated as a crystalline derivative; if this is not possible several independent criteria should be utilized. Enzymatic identification promises to be the most sensitive and accurate for metabolites which are found only in micro amounts.

2 THE METABOLISM OF LIPIDS

The lipids may be defined as those substances which are soluble in fat solvents; a few like the lecithins are somewhat water soluble but most are insoluble in water. Lipids which are of importance to the physiology of the fungi include the true fats, the fatty acids, the compound lipids (phospholipids), the sterols, the carotenoids and triterpenes and a few hydrocarbons. The general chemistry of the lipids is reviewed by Deuel (164). Those lipids which occur as constituents of the cell are considered in Chapter 2; here we are primarily concerned with synthesis and breakdown although for convenience the occurrence of some lipids is included.

The Synthesis of Fats, Fatty Acids and Phospholipids The neutral fats are triglycerides of the higher fatty acids. These are synthesized by many fungi in amounts up to 50 per cent of the mycelial dry weight. One rather striking difference between the lipids of fungi and those of animals is the high content of free fatty acids in most fungal lipids (Chapter 2). This means that most experiments on the factors affecting gross lipid content deal indiscriminately with both neutral fats and fatty acids.

Virtually all modern theories of acid biosynthesis assume a preliminary conversion of carbohydrate to pyruvic acid except of course in the biosynthesis of the sugar acids. Whether this assumption represents the facts or merely the limitations of current biochemical knowledge is uncertain. The phosphogluconate pathway of glucose metabolism (Chapter 7) provides to the cell a group of compounds including the 4 and 7-carbon sugar phosphates previously unknown and quite possibly of direct importance in acid synthesis.

Table 1 summarizes the reported formation of acids by fungi and actinomycetes acting in culture on carbohydrates. So far as possible duplicating reports have been omitted and some of those included are based on questionable or at least inadequate evidence. Satisfactory proof for the identity of an acid should involve more than color tests or the chromatographic constant in a single solvent; on the other hand the results of these presumptive tests are often correct and reports based on them cannot therefore be dismissed out of hand. Ideally an acid or other metabolic product should be isolated as a crystalline derivative; if this is not possible several independent criteria should be utilized. Enzymatic identification promises to be the most sensitive and accurate for metabolites which are found only in micro amounts.

2 THE METABOLISM OF LIPIDS

The lipids may be defined as those substances which are soluble in fat solvents; a few like the lecithins are somewhat water soluble but most are insoluble in water. Lipids which are of importance to the physiology of the fungi include the true fats, the fatty acids, the compound lipids (phospholipids), the sterols, the carotenoids and triterpenes and a few hydrocarbons. The general chemistry of the lipids is reviewed by Ducloux (164). Those lipids which occur as constituents of the cell are considered in Chapter 2; here we are primarily concerned with synthesis and breakdown although for convenience the occurrence of some lipids is included.

The Synthesis of Fats, Fatty Acids, and Phospholipids The neutral fats are triglycerides of the higher fatty acids. These are synthesized by many fungi in amounts up to 50 per cent of the mycelial dry weight. One rather striking difference between the lipids of fungi and those of animals is the high content of free fatty acids in most fungal lipids (Chapter 2). This means that most experiments on the factors affecting gross lipid content deal indiscriminately with both neutral fats and fatty acids.




Figure 3 Glucose concentration and fat synthesis by *Aspergillus fischeri* plotted from data of Prill Wenck and Peterson (498)

(318) and effects of metals which are definite but not susceptible of generalization (426-466). Fat formation in most yeasts and fungi is optimal at neutral or slightly alkaline reaction (238-128-485-560).

A supply of oxygen is invariably essential for fat synthesis (50). However cultivation with shaking reduces the fat content but not always the efficiency of conversion of glucose to fat (561). Here the effect is probably that of encouraging respiration and growth and thereby reducing the amount of carbon available for lipid synthesis.

Maximum yields of fat have been of interest because of possible industrial applications (270). Rippel (446) calculated that considering the usual values for protein content and for the economic coefficient the maximum synthesis by microorganisms the fat coefficient is 15 gm of fat per 100 gm of sugar utilized. Values usually found in the yeasts and fungi are in the range 5-15 (158-318) but cultivation of *Rhodotorula gracilis* under conditions which restrict protein formation to a minimum allows a value of 18 to be attained in aerated culture (179). Since lipids are more reduced than carbohydrate much of the glucose carbon must be oxidized to carbon dioxide in order that the necessary hydrogen be supplied.

The composition of the fat of a given species is relatively constant but external conditions appear to affect it in two ways. First the ratio of unsaturated to saturated fatty acids is claimed to be greater at low than at high growth temperatures (413-508). No such effect however was found in careful studies on *Aspergillus fischeri* (428). The ratio of saturated to unsaturated fats is also affected by glucose concentra-

tion and by pH (318) and in an as yet undetermined manner by cell pigments (403). Second the content of free fatty acids is greater at high carbohydrate levels. The acid value is also stated to be affected by pH (425) but the observed effects may be attributable to other causes.

The relation of fat formation to growth is exemplified in the data of Figure 4. As discovered early by Lindner (345), two phases of metabolic activity are distinguishable in many organisms: a phase of growth and a phase of fat accumulation. This is exactly comparable to the sequence of events in citric acid accumulation (Figure 2). Separation of growth and fat synthesis is not, however, always so clear (429). Figure 5 shows parallel courses of dry weight and fat production.

In animal and bacterial systems it seems clear that fatty acids are synthesized ultimately from 2 carbon fragments specifically acetyl coenzyme A (114). In the fungi precursor studies reviewed by Foster (200) early suggested the involvement of 2 carbon compounds in fat synthesis. More conclusive evidence has been obtained with several isotope techniques. Thus in a mutant of *Neurospora crassa* which is unable to form acetate from carbohydrate the carbon of fatty acids is derived almost entirely from exogenous acetate (407). The importance of acetate in fatty acid synthesis appears also from studies on isotope distribution in the fatty acids of *Phycomyces blakesleeana* (43-45), *Fusarium lini* (146) and *Ustilago zeae* (190).

Whether the unsaturated acids are derived directly from the saturated by dehydrogenation is an open question. Two types of evidence—nutritional responses of a fatty acid requiring mutant




Figure 4. Growth, sugar utilization, and fat formation in cultures of *Rhodotorula gracilis*. Redrawn from Enebo, Anderson, and Lindin (179) by permission of Academic Press, Inc.

of *Neurospora crassa* (336) and direct analyses of the mycelium of *Aspergillus nidulans* (475)—indicate that the routes of formation of the two types are independent. Evidence to the contrary (146) is less convincing. The only fatty acid dehydrogenase known in the fungi acts only on the short chain acids (387). In animal tissues it seems likely that the saturated and the unsaturated fatty acids arise independently (216).

Phospholipid synthesis has been studied by Raveux (442) who found the phospholipid content of *Aspergillus niger* to be independent of the fat content and to parallel the available nitrogen of the medium. The time course of phospholipid formation has been determined in *A. niger*: lipoidal phosphorus of the mycelium rises to a peak and then declines during the later—probably autolytic—phases of culture (222).

The Breakdown of Fats and Phospholipids Although the neutral fats are relatively poor sources of carbon they are utilized by many fungi (Chapter 3). The first step in such utilization is the hydrolysis of the fat to glycerol and fatty acids (288); it is assumed that endogenous fats are split in the same way before utilization. The hydrolysis is catalyzed by lipase or lipases.

It is common practice to assay for lipase activity by measuring the hydrolysis of simple esters, e.g. ethyl acetate or *p*-nitrophenyl acetate. It is doubtful, however, whether the simple esterases so determined are active in hydrolysis of the true fats (7). Simple esterases occur in most fungi investigated; the esterase of *Leizites septaria* fails to hydrolyze fats (574). Lipase proper, as distinct from esterases, can probably best be studied with water-soluble synthetic substrates (60, 168).

Virtually all fungi which have been studied for lipase production appear to form the enzyme (187, 317). Lipase is primarily intracellular, but both it and esterase do appear in the medium late in the culture cycle (153, 488, 570). It has been suggested (198) that *Aspergillus niger* and *Penicillium roqueforti* form two lipases, one in the mycelium and the other released into the medium.

Data on the relation of pH to activity and on the stability of fungal lipases vary widely from one organism to another (195, 198, 317, 440, 471, 512) so much so that the formation of different enzymes seems likely. The amount of extracellular lipase formed by *Mucor mucedo* is influenced by the temperature of cultivation and is increased by inclusion of a lipid in the growth medium (488). Like lipase from other sources, fungal lipase under appropriate conditions catalyzes the



synthesis of esters e.g. *n* butyl oleate from a fatty acid and an alcohol (440)

The phosphatides—lecithins and cephalins—are complex phospholipids in which one of the fatty acids found in the true fats is replaced by phosphoric acid linked to a nitrogen base. They are hydrolyzed at different points in the molecule by the phospholipases A, B, and C (498). Phospholipase A is found in *Aspergillus oryzae* (148) and *Lycopodium giganteum* (211); phospholipase B in the mycelium of *Penicillium notatum* (191). Phospholipase C, identical with glycerophosphatase (Chapter 9), is found in a commercial enzyme preparation from *Aspergillus* sp. (3).

The Metabolism of Fatty Acids The fatty acids occur both free and combined in mycelium (Chapter 2); information on their biosynthesis has been summarized above. In this section we are concerned with the breakdown of exogenous fatty acids arising in natural substrates by lipase action or supplied in pure culture as such.

The breakdown of fatty acids in animal tissues is believed to occur by the so-called β -oxidation condensation route; the oxidation proceeds by the removal of two carbon atoms at a time, with subsequent recombination of the 2 carbon fragments to form acetoacetic acid. Acetoacetate is formed by fungi during the oxidation of butyric acid (149, 388) but the details of its origin are not known. We are concerned therefore primarily with information bearing on the β -oxidation phase rather than on possible recombination processes.


Beta oxidation in animals is essentially the conversion of a fatty acid of *n* carbon atoms ($R-CH_2-CH_2-COOH$) to one of *n* - 2 carbons ($R-COOH$) and acetate or a derivative of acetate. The key intermediate is a β keto fatty acid and the reactions may be summarized as follows:






Coenzyme A is required at two points in the activation of the original fatty acid, forming an acyl coenzyme A, and in the formation of acetate, which appears as acetyl-coenzyme A. Repetition of reactions 4 and 5 leads ultimately to complete conversion of the long chain fatty acid to 2 carbon compounds. The process is reviewed in detail by Green (245) and by Chaikoff and Brown (114).

Evidence for β -oxidation in fungi derives almost entirely from studies on the breakdown of fats and fatty acids by growing or resting

cells. Interest in the problem was first aroused by the finding that methyl ketones are present in certain of the fungal cheeses and in edible oils attacked by fungi (482-494). In a number of experiments it has been shown that *Penicillium glaucum* converts a fatty acid to a methyl ketone of one less carbon atom, i.e.




This reaction has been demonstrated in experiments on many fatty acids (2-509-511) e.g.



The same general reaction is found in *Penicillium notatum* and in *Aspergillus* spp. (149-380-569).

Thaler and co-workers (510) propose the following sequence of reactions in the formation of methyl ketones from fatty acids



The evidence for this sequence is that methyl ketones arise during the growth of *Penicillium glaucum* on α, β unsaturated acids and β hydroxy acids. Thus methylpropyl ketone is formed from hexanoic acid, 2-hexanoic acid and β -hydroxyhexanoic acid. Further, fungi are known to form β -keto acid decarboxylases (309).

This scheme up to the formation of the β -keto acid agrees with our current understanding of the process in animal cells, but until enzymatic studies are made the identity of the two systems remains hypothetical. The last step—formation of the methyl ketone—does not go on appreciably in the normal animal cell, which instead forms as mentioned above a 2-carbon compound and a new fatty acid. This suggests that methyl ketone formation by fungi is an abnormal or at least unusual reaction, perhaps to be attributed to toxicity of the substrate fatty acid (494). It is reasonable to suppose that complete oxidation of fatty acids in the fungi follows the animal system with liberation of acetyl-coenzyme A or free acetate as the last step. The

strongest evidence for this supposition is in the finding by Webley and others (544) that the oxidation of *m*-substituted phenyl fatty acids by *Nocardia opaca* an actinomycete yields end products in conformity with predictions based on β -oxidation. Comparable studies on the fungi have not yet appeared.

Indirect evidence for β -oxidation in the fungi is provided by the incorporation of fatty acid carbon into the side chain of penicillin (380-516). The most effective precursors are fatty acids which by their structure are not susceptible to complete β -oxidation.

It has been argued from simultaneous adaptation experiments that *Streptomyces aureofaciens* oxidizes octanoate by some mechanism other than β -oxidation (276). However it is now known that the adaptation technique is unreliable in fatty acid oxidation since the adaptive system may be that which activates the fatty acid. Failure of octanoate adapted cells to oxidize butyrate may therefore mean only that activation of the two acids is carried out by different enzymes.

Fatty acid oxidase systems probably occur in fungi and actinomycetes (387-543). In addition mycelial extracts of *Penicillium glaucum* and of *Aspergillus* spp. contain a lipoxidase capable of oxidizing unsaturated fatty acids (387).

The Metabolism of Steroids and Triterpenes Sterols occur in small amounts in the lipid fraction of most fungi. Ergosterol appears to be the compound of this group most often formed by fungi, but other sterols have been tentatively identified (Chapter 2). Cultural conditions which favor sterol formation are reviewed by Bernhauer (50). It appears that sterol and fat synthesis are competitive reactions; the effect of pH on their formation (428) is perhaps the strongest evidence for this competition. Biochemical studies on yeast and on fungi indicate that ergosterol is formed in large part from 2 carbon compounds, although isovaleric acid may also be involved (146-318-407). The amount of sterol formed varies among strains and among single spore cultures of a given strain (556).

The breakdown of exogenously supplied steroids has been studied in two connections: (1) the utilization of steroids for energy and carbon and (2) partial and limited transformations of the steroid molecule.

Cholesterol is oxidized in soil (521) by actinomycetes and by Gram negative bacteria; it is not clear which organisms are responsible for steroid breakdown in natural soils (462-523-525). Some fungi are apparently able to grow with ergosterol as carbon source (462) and some of the products of the action of *Proactinomyces erythropolis* on

steroids have been identified (522-524-526) but most of the pathway of steroid oxidation is unknown

Specific microbiological transformations of steroids i.e. reactions at single specific points on particular steroid molecules have been studied extensively first with bacteria and yeast (364-532) later with actinomycetes (271-329-330-555) and fungi (423). The pharmacological importance of these processes has stimulated since 1952 a very large volume of experimental work for which the fungi and actinomycetes have been the best material. The known processes may be classified as follows (557)

1 Hydroxylations specific introduction of the —OH group. Examples are shown in Table 2 it will be noted that among the most frequent positions of attachment are 6β , 11α , 11β and 17α

Table 2 The Hydroxylation of Steroids by Fungi and Actinomycetes*

Organism	Steroid Position Hydroxylated†
<i>Aspergillus</i> spp	6β 11α 17α 21
<i>Cephalothecium roseum</i>	6β 11α 17α
<i>Colletotrichum antirrhini</i>	15α
<i>Coniothyrium</i> sp	11β
<i>Cunninghamella blakesleeana</i>	11β
<i>Curvularia</i> spp	7α 8β (?) 11β 14α
<i>Dactylium dendroides</i> ‡	11α 17α
<i>Didymella radakii</i>	16α
<i>Gibberella baccata</i>	15α
<i>Helicostylum piriforme</i>	6β 8β 11α 14α
<i>Lenzites abietina</i>	6β 15β
<i>Mucor</i> spp	6β 8β (?) 14α
<i>Neurospora crassa</i>	8β or 9α
<i>Ophiobolus herpotrichus</i>	21
<i>Penicillium</i> spp §	11α
<i>Peziza</i> spp	7α
<i>Phycomyces blakesleeana</i>	7β 15β
<i>Proactinomyces roseus</i>	7β
<i>Rhizopus</i> spp	6β 7β 8β 11α
<i>Streptomyces</i> spp	6β 8β or 9β (?) 11β 16α
<i>Trichothecium roseum</i>	6β 17α

* Except as noted from Wettstein (557). A more complete list appears in the review of Eppstein et al. (182a)

† The specific compounds attacked and often the yield of product are given in the original papers cited by Wettstein (557)

‡ Dulaney et al. (173)

§ Dulaney et al. (172)

2 Hydrogenation of steroid aldehydes or ketones by bacteria yeast *Streptomyces* spp. and *Rhizopus nigricans*

3 Hydrogenation of ring carbon atoms by *Clostridium* sp. *Streptomyces* sp. and *Rhizopus nigricans*

4 Dehydrogenation of steroid secondary alcohols by bacteria and actinomycetes

5 Epoxidation of steroids containing an isolated double bond by *Curvularia lunata* and *Cunninghamella blakesleeana*

6 Dehydrogenation in ring A with or without degradation of the side chain by *Fusarium solani* *Colanectria decora* *Cylindrocarpum radicola* and other fungi


7 Oxidation of the side chain only e.g. by *Gliocladium catenulatum*

8 Ring cleavage with lactone formation e.g. by *Penicillium chrysogenum*

Specific enzyme systems must of course be responsible for these transformations and some knowledge of animal and bacterial enzymes is available (244-577). The 6β hydroxylation of progesterone by *Aspergillus ochraceus* is accelerated by zinc: it has been suggested that the metal is required for the synthesis of an induced enzyme which carries out the reaction (174).

Steroidal sapogenins e.g. diosgenin appear not to be hydroxylated by fungi or other microorganisms (376).

Chemical studies on several basidiomycetes have brought to light a new series of fungal metabolites the tetracyclic triterpenes. As terpenes they are related to the carotenoids but their structure also bears an obvious relation to that of the sterols: they are not however true sterols. Similar compounds are known in higher organisms e.g. betulin in plants and squalene in animals. The structure of eburonic acid formed by several Hymenomycetes (151-220-281) may be taken as an example of the group.




Other compounds of this type include the closely related polyporenic acids A B and C isolated from *Polyporus* spp and *Poria cocos* (91 92 151 154 257 260). The triterpenoids are biosynthetically related as judged by the occurrence of all three polyporenic acids in one species (154) and by the fact that three strains of *Poria cocos* form polyporenic acid B and a fourth forms eburicoic acid (151). In *Polyporus eucalyptorum* eburicoic acid constitutes at least 20 per cent of the dry weight of the mycelium (220) and must therefore be considered a major metabolite. The chemistry of these compounds has been reviewed by Jones and Halsall (306). triterpenoids occur also as lichen products (24).

The antibiotics cephalosporin P_1 and helvolic acid are tetracyclic compounds and may be related to the steroids but their complete structure is not known (100).

The Metabolism of Carotenoids The carotenoid pigments which appear among the unsaponifiable lipids include (1) the true carotenes e.g. β carotene and lycopene and (2) oxygen containing derivatives e.g. rhodoviolascins. All may be considered as built up of

eight isoprene units the carbon skeleton of which is $\text{C}=\text{C}(\text{C})-\text{C}=\text{C}$. In addition to the colored carotenoids colorless polyenes are found in the fungi and will be considered with the pigments.

The formula of β carotene illustrates the essential features of the carotenoid pigments.



Carotenoids are of widespread but not universal occurrence in the fungi including representatives of all the major taxonomic groups (232 236 264) Table 3 lists the carotenoids of a few fungi which

Table 3 Carotenoids of Some Fungi

Organism	Carotenoids
<i>Allomyces jamaicus</i>	γ carotene * β carotene lycopene phytofluene (178 529)
<i>Phycomyces blakesleeanus</i>	α carotene β carotene * γ carotene ξ -carotene † lycopene neurosporene phytofluene phytoene (233, 310)
<i>Neurospora crassa</i>	β carotene γ carotene * θ -carotene lycopene * spirilloxanthin (rhodoviolascin) * lycopanthin neurosporene * phytofluene phytoene unidentified compounds (264)
<i>Dacrymyces stillatus</i>	α carotene β carotene * γ carotene (?) ξ carotene † torulene cryptoxanthin zeaxanthin phytofluene phytoene (?) (234)
<i>Puccinia</i> spp	β carotene γ -carotene (?) lycopene (293)

* Quantitatively important

† See Havo (264) on fungal ξ -carotene

have been intensively studied. In contrast to higher plants β -carotene is not so nearly universal in the fungi and acidic carotenoids e.g. torularhodin are more common (232).

The intracellular distribution of fungal carotenoids has not been systematically investigated. In certain of the higher fungi carotenoid pigments occur both in rod shaped structures apparently part of the mitochondrial system and in oil droplets in the cytoplasm (266 267). The carotenoids of the male gametangia of *Allomyces* spp are found in cytoplasmic oil droplets (178) those of *Rhodotorula rubra* in granules in the pericapsular region (127).

The physiological conditions of carotenogenesis are to some degree specific for individual organisms. In *Phycomyces blakesleeanus* the time course of carotene formation (Figure 5) indicates that the bulk of the carotene is formed after growth has been completed and that carotenoid formation is independent of lipid formation. Independence of fat and carotene synthesis also appears from their divergent pH requirements (238). Carotenogenesis is affected of course by carbon and nitrogen nutrition but the data do not afford a basis for generalization. Cultivation of *Phycomyces blakesleeanus* at low temperature decreases total carotene synthesis but does not affect the




Figure 5 Growth fat formation and carotenogenesis in *Phytomyces blakesleeanus* Redrawn from Goodwin and Willmer (238) by permission of the Cambridge University Press

proportions of the different carotenoids formed (215) as it does in *Rhodotorula rubra* (392). Oxygen is required for carotene formation by *Neurospora crassa* (572) and presumably this is true of other fungi.

The mycelial carotenoids of *Neurospora crassa* do not develop in complete darkness although the conidial carotenoids do; this problem has been studied by Zalokar (572-573). Very brief exposure to light is sufficient to activate pigment synthesis which is proportional to light intensity up to a saturation value. The action spectrum of light activation is consistent with but does not of course prove the participation of a flavoprotein as the photoreceptor. Calculated quantum yields are so high as to make it necessary to postulate that activation is an indirect process. Carotenogenesis in other fungi is stimulated by light but does not require it (131-219). Synthesis by green plants does not require light.

Carotenoid formation is inhibited by streptomycin, chloramphenicol and isonicotinic hydrazide at concentrations which do not inhibit growth (235). Diphenylamine inhibits carotenogenesis in both bacteria and fungi (233-528). Its effect is to reduce the absolute and relative amounts of the more unsaturated carotenoids e.g. β -carotene and to increase the relative amounts of the more saturated e.g. phytofluene and neurosporene.

The biosynthesis of the carotenoids may be considered under three

heads (1) the origin of the isoprene skeleton (2) the mode of condensation of isoprene units to form the 40-carbon chain of the carotenoids and (3) the relation of the different carotenoids to one another


On grounds of comparative biochemistry and especially from studies on higher plants (304) the most likely route of synthesis of the isoprene skeleton involves the following steps (236 346)

- 1 Formation of acetoacetic acid from acetate and coenzyme A
- 2 Condensation of acetoacetate with acetyl coenzyme A to form β hydroxy β methylglutaryl coenzyme A
- 3 Conversion of β hydroxy β methylglutaryl-coenzyme A to β methylcrotonyl-coenzyme A This process involves more than one reaction and includes a decarboxylation

The condensation of two 5-carbon units to yield a 10 carbon chain followed by successive reactions with new 5 carbon units could in principle yield the 40-carbon chain of the carotenoids (346)

Evidence for the foregoing scheme in the fungi is somewhat limited but adequate to establish it as a working hypothesis Carotenogenesis in fungi is stimulated by β methylcrotonaldehyde (237) and by pantothenic acid and pantethine which enter into the coenzyme A molecule (249) The labeling of the β carotene of *Mucor hiemalis* grown on C^{14} acetate is qualitatively consistent with the proposed mechanism (247 248 248a) That fungi form carotenoids during cultivation on acetate (463) cannot be urged as proof for the hypothesis although again the fact is consistent with it

The final set of problems is associated with the relationship of the carotenoids to one another From genetic data on higher plants Porter and Lincoln (427) proposed a unilinear sequence of formation of the carotenoids by successive dehydrogenations that is that the unsaturated hydrocarbons are derived from the saturated as follows



Although this scheme fits the behavior of mutants of *Chlorella vulgaris* (138) *Rhodotorula rubra* (89) and *Neurospora crassa* (264 469) it cannot be regarded as proved Goodwin (232 233) has argued largely on the basis of inhibitor studies that the more saturated polyenes e.g. phytofluene are formed by one pathway the less saturated e.g. β -carotene by another both from a common precursor The effects of added methylheptenone and β ionone have been interpreted as

supporting the second hypothesis (361) It must be concluded that the question is still open

It is usually assumed that the oxidized carotenoids neutral and acidic xanthophylls arise by oxidation of the hydrocarbon polyenes (264)

Although β ionone increases carotenogenesis in *Phycomyces blakesleeanus* isotope evidence indicates that the molecule is not as one might expect incorporated as such into the β carotene structure (182)

Two functions have been suggested for the carotenoids a direct role as receptor for the light stimulus in phototropism and an undefined role in sexual differentiation

Phototropism is discussed later (Chapter 11) and it is concluded that there is no satisfactory proof of the direct involvement of carotenoids

Although the carotenoid pigments and the colorless polyenes of *Allomyces javanicus* are restricted to the male gametangia (178 529) an essential role in the sexual process is rendered less probable by the finding (527) that diphenylamine inhibits carotenogenesis in *A. javanicus* without appreciably affecting the sexual reaction It seems likely that the carotenes are side products of a metabolism which is in some respects at least distinctive to the male gametangium The (+) and (-) strains of some of the Mucorales show quantitative differences in carotenoid content studies as reviewed by Goodwin (232) have been confined to the fact itself and it would appear that in *Phycomyces blakesleeanus* the difference between (+) and (-) strains is not consistent (219) Zygosporangium formation or merely pairing of compatible strains in some of the Mucorales is accompanied by increased formation of carotene (29 270a)

The most serious deficiency in the study of fungal carotenoids is the failure to determine whether or not carotenes occur in association with proteins or lipoproteins Such an association is theoretically necessary if they are to function in metabolism e.g. in the reception of light stimuli Carotenoproteins are well known in animals (234) and the now classical studies on rhodopsin (147 537) provide a model for the investigation of other pigments

The Metabolism of Hydrocarbons and Hydrocarbon Derivatives
It has been mentioned (Chapter 3) that hydrocarbons support growth of some actinomycetes and fungi The oxidation of *n*-dodecane and other hydrocarbons by *Proactinomyces opacus* can be demonstrated manometrically (Figure 6) the organism also oxidizes long chain saturated fatty acids which may be intermediates in hydrocarbon oxidation Details of the breakdown of hydrocarbons are not known




Figure 6 The oxidation of hydrocarbons by *Fractinomyces opacus* Redrawn from Wehley and Dekock (543) by permission of the Cambridge University Press

ZoBell (577) reviews our knowledge of the process in bacteria. Intermediates appearing during the oxidation of naphthalene by *Nocardia* sp. suggest that benzenoid compounds may be formed (520).


Hydrocarbons produced by fungi fall into two groups: ethylene and its derivatives, and the substituted hydrocarbon polyenes and polyenes.

From the biological effects of volatile products of *Penicillium digitatum* it seemed probable that the organism produces ethylene (59, 378). This has been confirmed (571); ethylene is probably produced also by *Blastomyces dermatitidis* (394) and by *Fusarium oxysporum* v. *lycopersici* (167). However, the ethylene produced by fungus-infected plant tissues is believed to be a host product stimulated quantitatively by infection rather than a fungus metabolite (559).


An interesting volatile hydrocarbon, 2-methylbutene-2 (trimethyl ethylene), is formed by uredospores of *Puccinia graminis tritici* and is perhaps responsible for self-inhibition of germination of the rust spores (199).

Several antibiotics produced by fungi and actinomycetes are in two

related classes the polyenyne and the polyene (406). The polyenyne characterized by a system of conjugated triple bonds include among compounds of known structure, diatetrynes 1 and 2 of *Clitocybe diatreta* (10), agrocybin of *Agrocybe dura* (305-312) and mycomycin produced by *Nocardia acidophilus* (113). Some have nitrogenous substituents e.g., diatetryne 2



Mycomycin, on the other hand is an acid containing no nitrogen



Acetylenic compounds are also known in higher plants and in other fungi than those named here (9-11, 97). In an unidentified basidiomycete the formation of polyacetylenes early in the culture cycle is followed by their disappearance; this may be interpreted as a reutilization of the compounds (98).

The conjugated polyenes include a number of antibiotics the complete structure of which is not known; the polyene structure is inferred from their ultraviolet absorption spectra (28a, 530, 534, 558). All are fungistatic agents. The polyenyne except for mycomycin have been found only among basidiomycetes; the known polyene antibiotics on the other hand are produced primarily by actinomycetes. Grifolin formed by *Grifola confluenta* (275) is a substituted hydrocarbon with one pair of conjugated double bonds.







Species of *Corticium* produce two yellow pigments both containing a system of conjugated double bonds comparable to that of the carotenoid pigments (183, 246); other aspects of their structure however relate them to atromentin and thelephoric acid considered later with the aromatic compounds.

The sporophore of *Lactarius* spp. and probably the ascocarp of *Peziza* sp. contain a rubberlike polymer, a *cis* polyisoprene (491).

3 THE METABOLISM OF AROMATIC COMPOUNDS


Aromatic Compounds of Fungi. The fungi resemble the higher plants and differ from the bacteria in that many species convert carbohydrate to aromatic compounds not known to be essential metabolites. These compounds have been reviewed in detail by Birkinshaw (63), by Rustrick (130) and in lichens by Asahina and Shibata (24). Any classification of them must of necessity be based on their chemical structure and may unintentionally obscure physiological and biosynthetic relations. Thus a substituted benzene and the quinone

Table 4 Representative Aromatic Compounds Produced by Fungi


Group*	Compound	Organism(s)
I	Methyl cinnamate $\text{CH}=\text{CH}-\text{COOCH}_3$ 	<i>Lentinus lepideus</i> (74)
	Anisaldehyde CHO  OCH_3	<i>Trametes suaveolens</i> (64) <i>Daedalea juniperina</i> (69) <i>Polyporus ben oinus</i> (77)
	Gentisyl alcohol OH  CH_2OH	<i>Penicillium</i> spp (66 93 176 180)
	Homoprotocatechuic acid CH COOH  OH OH	<i>Polyporus tumulosus</i> (437)
3 5	Dihydroxyphthalic acid OH  COOH COOH HO	<i>Penicillium brev-compactum</i> (142 408)
	Gallic acid COOH  HO OH OH	<i>Phycomyces blakesleeana</i> (44)

* See text

related classes the polyenyne and the polyenes (406). The polyenyne characterized by a system of conjugated triple bonds include among compounds of known structure diatetrynes 1 and 2 of *Clitocybe diatreta* (10) agrocybin of *Agrocybe dura* (305-312) and mycomycin produced by *Nocardia acidophilus* (113). Some have nitrogenous substituents e.g. diatetryne 2



Mycomycin on the other hand is an acid containing no nitrogen



Acetylenic compounds are also known in higher plants and in other fungi than those named here (9-11, 97). In an unidentified basidiomycete the formation of polyacetylenes early in the culture cycle is followed by their disappearance; this may be interpreted as a reutilization of the compounds (98).

The conjugated polyenes include a number of antibiotics the complete structure of which is not known; the polyene structure is inferred from their ultraviolet absorption spectra (28a, 530, 534, 558). All are fungistatic agents. The polyenyne except for mycomycin have been found only among basidiomycetes; the known polyene antibiotics on the other hand are produced primarily by actinomycetes. Grifolin formed by *Grifola confluenta* (275) is a substituted hydrocarbon with one pair of conjugated double bonds.







Species of *Corticium* produce two yellow pigments both containing a system of conjugated double bonds comparable to that of the carotenoid pigments (183, 246); other aspects of their structure however relate them to atromentin and thelephoric acid considered later with the aromatic compounds.

The sporophore of *Lactarius* spp. and probably the ascocarp of *Peziza* sp. contain a rubberlike polymer, a *cis* polyisoprene (491).

3 THE METABOLISM OF AROMATIC COMPOUNDS

Aromatic Compounds of Fungi. The fungi resemble the higher plants and differ from the bacteria in that many species convert carbohydrate to aromatic compounds not known to be essential metabolites. These compounds have been reviewed in detail by Birkinshaw (63), by Raistrick (430) and in lichens by Asahina and Shibata (24). Any classification of them must of necessity be based on their chemical structure and may unintentionally obscure physiological and biosynthetic relations. Thus a substituted benzene and the quinone

Table 4 Representative Aromatic Compounds Produced by Fungi

Group*	Compound	Organism(s)
I	Methyl cinnamate $\text{CH}=\text{CH}-\text{COOCH}_3$ 	<i>Lentinus lepideus</i> (74)
	Anisaldehyde CHO  OCH_3	<i>Trametes suavis</i> (64) <i>Daedalea juniperina</i> (69) <i>Polyporus benzoinus</i> (77)
	Geranyl alcohol OH  CH_2OH	<i>Penicillium</i> spp (66 93 176 180)
	Homoprotocatechuic acid CH COOH  OH OH	<i>Polyporus tumulosus</i> (437)
	3,5-Dihydroxyphthalic acid OH  HO COOH COOH	<i>Penicillium brev-compactum</i> (142 408)
	Gallic acid COOH  HO OH OH	<i>Phycomyces Blakesleeanus</i> (44)

* See text

Table 4 (continued)







Group*	Compound	Organism(s)
I	<p>Gladiolic acid (tautomeric)</p> 	<i>Penicillium gladioli</i> (253 254)
	<p>Cyclopolic acid</p> 	<i>Penicillium cycloptum</i> (84)
	<p>Cyclopaldic acid</p> 	<i>Penicillium cycloptum</i> (84)
	<p>Chloramphenicol</p> 	<i>Streptomyces venezuelae</i> (444)
	<p>Mycophenolic acid</p> 	<i>Penicillium</i> spp (6 65 83 142)
	<p>Sulochrin</p> 	<i>Oospora sulphurea ochracea</i> (395 396)

Table 4 (continued)






Group*	Compound	Organism(s)
II	4 Methoxy 2 5 <i>p</i> -toluquinone 	<i>Coprinus similis</i> <i>Lentinus degener</i> (12)
	Spinulosin 	<i>Aspergillus fumigatus</i> <i>Penicillium</i> spp (16 79 94)
	Oosporein 	<i>Oospora colorans</i> (328) <i>Chaetomium aureum</i> (347)
	Atromentin 	<i>Paxillus atromentatus</i> (325)
	Telephonic acid 	<i>Telephora</i> spp <i>Hydnum ferrugineum</i> (326) <i>Lobaria retigera</i> (lichen) (23)

Table 4 (continued)












Group*	Compound	Organism(s)
III	Chrysophanic acid	<i>Penicillium islandicum</i> (289)
		
	Emodin	<i>Dermocybe sanguinea</i> (327)
		
IV	Asperthecin	<i>Aspergillus quadrilineatus</i> (290)
		
	Ravenelin	<i>Helminthosporium ravenelii</i> (389-431)
		
V	Pinselic acid	<i>Penicillium amarum</i> (390)
		
V	Javanicin	<i>Fusarium javanicum</i> (20 21-61a)
		

Table 4 (continued)

Group*	Compound	Organism(s)
VI	Stipitatic acid	<i>Penicillium stipitatum</i> (68 150 165)
		
	Puberulonic acid	<i>Penicillium puberulum</i> (25 81)
		
VII	Lecanoric acid	<i>Parmelia</i> spp (lichens) (24)
		
	Gyrophoric acid	<i>Gyrophora esculenta</i> and other lichens (24)
		
VIII	Nidulin	<i>Aspergillus nidulans</i> (163)
		

formed from it by oxidation are more closely related biologically than are two differently substituted benzene compounds. The following division of the aromatic products synthesized by fungi taken largely from Raistrick (430) should be considered as a formal classification only. Examples of the major types are illustrated in Table 4.

Group I substituted benzenes is the largest single group of aromatic compounds and presents a bewildering array of possible biological interrelations. Some of the members of this group are soluble metabolites of the culture filtrate; others occur in the mycelium and still others are associated with reproductive structures (214-243).

Members of Group II substituted benzoquinones often have an obvious metabolic relation as oxidation products to phenolic compounds of Group I. Many are derivatives of toluquinone.

Anthraquinones Group III are pigments and are known also in lichens and in higher plants (280). The fungal anthraquinones may be considered as derivatives of chrysophanic acid.

Group IV the xanthenes is represented in the fungi by only a few certain examples. Rubrofusarin formed by *Fusarium* spp. (389) is probably the same methyl xanthone as asperxanthone of *Aspergillus niger* (355).

Group V comprises the naphthoquinones. In addition to javanicin other fungal products of this type have been identified as products of *Fusarium* spp. (104-451) and other fungi (40-161). Solanione (552) and oxyjavanicin (21) are probably identical with respectively javanicin and fusarubin (63).

Derivatives of tropolone Group VI in Table 4 are only quasi-benzenoid inasmuch as the ring contains seven carbons. Such higher plant products as thujaplicin and colchicine are tropolones (412). An origin of tropolones from phenolic compounds has been suggested (371).


The depsides Group VII are known so far only from lichens; it may be presumed by analogy with other lichen products (112) that the fungus component of the lichen is responsible for their formation.

Nidulin illustrates the depsidones (Group VIII). The lichen depsidones are similar; some e.g. gangaleoidin and diploicin are chlorinated; most are not (24).

The physiological importance of the aromatic compounds is largely unknown; most effort to date having been concentrated on problems of identification. In what follows three topics will be briefly considered: pathways of biosynthesis, enzymatic transformations, and the possible metabolic significance of the aromatic compounds.

The Biosynthesis of Aromatic Compounds Although not all of the aromatic compounds of fungi have been proved to be formed from carbohydrate—some have been reported only from sporophores in nature—most of them are probably so formed this is evidenced for example by the very high yields often obtained on synthetic media (95 128 252)

The origin of the aromatic ring from glucose has been investigated most successfully in connection with the biogenesis of the aromatic amino acids and *p* aminobenzoic acid in bacteria. The current view of this process may be diagrammed as follows (162)



In this scheme an unknown compound V arises from carbohydrate and is converted through the known materials shown to a second unidentified compound Z₁ which is in turn ultimately transformed into the aromatic amino acids and *p* aminobenzoic acid the known later steps in the process are reviewed in Chapter 8. The initial steps of the reaction sequence are not known preliminary evidence (162) suggests that the carbon chain of glucose is ruptured before incorporation of glucose carbon into shikimic acid.

A mutant of *Neurospora crassa* which requires *p* aminobenzoic acid tyrosine phenylalanine and tryptophan for growth is similar in these requirements to the bacterial mutants which have provided much of the evidence for the scheme just outlined and appears to be blocked in the conversion of dehydroshikimic acid to shikimic acid (507). It accumulates protocatechuic acid which is then oxidized by ring fission (p 174). Protocatechuic acid is known as a metabolic product of *Phycomyces blakesleeanus* growing on a glucose asparagine medium.

(44) and is also formed by many fungi during the oxidation of quinic acid (53-57)

Isotope studies on *Neurospora crassa* indicate that as in bacteria 3 and 4 carbon fragments derived from glucose—presumably via the phosphogluconate oxidation pathway (Chapter 7)—combine to form shikimic acid (506)

As discussed later currently available information at least encourages the speculation that all aromatic compounds arise analogously to protocatechuic acid in *N. crassa* i.e. that the normal pathway to aromatic amino acids is blocked or saturated and that the carbon of shikimic acid or its precursors is diverted to the synthesis of new aromatic structures. However in *N. crassa* protocatechuic acid is oxidized with ring fission and does not give rise to other aromatic compounds

Another possibility has been considered by Elvensvard (176) that pyrones might be intermediates in the conversion of glucose to aromatic compounds by *Penicillium urticae*. Evidence so far is limited to the action of pyranes e.g. dehydracetic acid on the synthesis of aromatic compounds

Close structural relations among the aromatic compounds are numerous both within and between species. Many are discussed by Raistrick (450) a few examples may be cited

1 Simultaneous formation by one organism of a phenol and the corresponding quinone (14-180)

2 Occurrence in the same organism of a compound and its dihydro derivative e.g. cyclopolic and cyclopaldic acids (Table 4)

3 Formation of a compound and a hydroxy or methoxy derivative of it e.g. gladiolic and cyclopaldic acids (Table 4) and methyl cinnamate and methyl *p*-methoxycinnamate (74). The ability of fungi to introduce hydroxyl groups into a ring has been noted above in connection with steroid metabolism

4 The close structural relationship already mentioned of all of the fungal anthraquinones found in such diverse genera as *Aspergillus*, *Cortinarius*, *Helminthosporium* and *Penicillium* (289). A possible origin of the anthraquinones from benzophenone is discussed by Tatum (505)

The foregoing examples refer to more or less static relations. In a dynamic and more conclusive experiment it has been shown that the ratio of patulin to gentisyl alcohol is controlled by the concentration of iron in the medium (Figure 7). Similarly *Penicillium urticae* in a low zinc medium forms predominantly 6-methylsalicylic




Figure 7 The effect of iron on the formation of patulin and gentisyl alcohol by *Penicillium* sp. From data of Brack (93)

acid but at higher concentrations of zinc this compound is replaced by a mixture of gentisyl alcohol, patulin and toluquinone (176). This type of analysis in which an environmental modification causes a shift in metabolic products is extremely promising.


Penicillium brevicompactum produces several related phenols. Both the temporal sequence of appearance of these compounds (409) and the observation that the organism when fed one compound produces others (230) support the proposition that these structurally related molecules are related biologically as well.

A role of the polyphenol oxidases in the conversion of phenols to other compounds is probable but has not been investigated in the fungi. Mason (371) reviews several theories on the function of these enzymes in the origin of aromatic products of higher plants.

Enzymatic Transformations of Aromatic Compounds The major known types of attack on aromatic compounds may be summarized as (1) the oxidation of aliphatic side chains, (2) the oxidation of phenols, (3) the oxidative fission of the aromatic ring, and (4) the hydrolysis of depsides.

The side chain of phenylacetic acid is oxidized by *Penicillium chrysogenum*; benzaldehyde is formed during the oxidation and is


thought to be an intermediate (279). The process therefore resembles the oxidation of substituted benzenes by bacteria (479) and higher plants (313) in which benzoic acid is the ultimate product



However the entire sequence is not known in fungi and the formation from phenylacetate of 2-hydroxyphenylacetate and 2,5-dihydroxyphenylacetate point to other possible modes of attack by fungi and actinomycetes (297-320, 544). Benzoic acid is formed in culture by *Penicillium roscopurpureum* (427a).

The oxidation of vanillin and ferulic acid by fungi appears to be of the same general type as that of phenylacetate, the side chain being oxidized in one or more steps to a carboxyl group (268).

There occurs in bacterial metabolism an oxidative fission of the aromatic ring of catechol to yield *cis-cis* muconic acid (480)



This overall reaction has been hypothesized for *Aspergillus niger* (57) and *Penicillium* sp. accumulates *cis-cis* muconate during the oxidation of catechol (333). The aromatic ring of phenylacetic acid is split by *Penicillium chrysogenum* but the details of the process are not known except that an induced enzyme system is responsible and that compounds of the tricarboxylic acid cycle appear in the medium during oxidation (294-296).

It has been mentioned that a mutant of *Neurospora crassa* accumulates and then oxidizes protocatechuic acid. The ultimate product is β -keto adipic acid and the enzymatic steps involved are similar to but not identical with those known in *Pseudomonas fluorescens* (250-481). The first step is the conversion of protocatechuic acid to *cis-cis* β -carboxymuconic acid analogously to the reaction of Equation 12 by an inducible oxidase system (250-251).


The oxidation of phenols to quinones is a special problem and much more data are available on it than on other phases of aromatic metabolism.

The enzymes responsible for the oxidation of phenols are the polyphenol oxidases of which two have been separated

1 Tyrosinase which oxidizes both monophenols (tyrosine *p*-cresol etc.) and polyphenols (catechol pyrogallol etc.) Oxidation of tyrosine leads by a complex series of enzymatic and non-enzymatic reactions to the formation of black pigments the melanins (Chapter 8)

2 Laccase which oxidizes polyphenols but not monophenols In contrast to tyrosinase laccase is sensitive to carbon monoxide and has a somewhat different substrate range

Both enzymes are copper proteins The characteristic reaction often used for assay is the oxidation of catechol to *o* benzoquinone




Although other polyphenol oxidases probably occur in fungi (108 286) they have not been characterized Most assays do not indeed discriminate between laccase and tyrosinase

Interest in the polyphenol oxidases has been stimulated by the Bavendamm reaction (35) of the wood rotting fungi on an agar medium containing gallic or tannic acid the fungi causing white rots of wood and capable of utilizing lignin form a colored zone The brown rot fungi in general do not form the zone Several surveys of the wood rotting fungi have established that the correlation between zone production and type of rot is high although not perfect (86 160 397) The usual substrates do not distinguish between tyrosinase and laccase but work with specific substrates and on the distribution of the enzymes between cells and medium indicates that extracellular laccase rather than tyrosinase is responsible for the Bavendamm reaction (274 335 313 314)

Extracellular polyphenol oxidases are formed also by other basidiomycetes including parasitic and mycorrhizal species and both coprophilic and litter inhabiting saprophytes (340) Fungi other than basidiomycetes have been less well studied tyrosinase is found in *Clomerella cingulata* (500) and *Neurospora crassa* (210 287) and—judged by color reactions—in *Streptomyces scabies* (282) and in *Sclerotinia fructicola* and *Macrosporium sarcinaeforme* (445)

Polyphehol oxidases in the fungi are often present only at particular times in the culture cycle especially during autolysis (188-500) or only in particular morphological structures (342-344). The medium used affects the production of the enzymes both positively and negatively. Laccase formation by *Polyporus versicolor* is increased non specifically by constituents of the medium and specifically by certain aromatic compounds which probably act as inducers (189-190). Both carbon source and growth temperature affect enzyme synthesis in *Glomerella cingulata* (499). Negative effects of an interesting type have been found in studies on *Neurospora crassa* (287) in a medium containing the usual level of sulfate a sulfur containing dialyzable tyrosinase inhibitor is formed the result being that the conventional assay detects tyrosinase only in cultures grown in a low sulfate medium. This finding requires that earlier negative reports be re-evaluated.

The tannins are esters of polyhydroxy phenols with aromatic acids and are hydrolyzed by tannase e.g.




The enzyme an esterase also hydrolyzes other similar esters (498). The tannase of *Aspergillus niger* is inducible (322) its properties and purification are summarized by Dyckerhoff and Armbruster (175) and partial purification has been reported (519). Although most studies have been made on the tannase of *A. niger* the enzyme has been reported in a few other fungi (36-448-574).

The Significance of Aromatic Compounds Some of the aromatic compounds bulk rather large in the economy of the organism amounting to an appreciable fraction of the glucose consumed (140) and of the dry weight produced (128). Those which have been studied in relation to the history of the culture appear to be first formed and then utilized (93-95) to be more than inactive waste products of metabolism.

Possibly the most likely metabolic significance of these compounds is as products of an unbalanced nutrition and metabolism. Most of them have been isolated from fungi grown on media very high in carbon thus the glucose Czapek Dox medium often used has a carbon-nitrogen ratio of 60. Under these conditions it is not surprising

that carbon may flow into new metabolic pools simply because of an insufficiency of nitrogen. We may visualize this type of process schematically



If compound D for example cannot be synthesized because of a relative nitrogen deficiency enzymatic equilibria and factors of permeability and solubility will determine which of the potential alternative pathways is followed. In the specific instance of the benzenoid compounds and their oxidation products it is imaginable that the synthesis of aromatic amino acids stops because nitrogen is no longer available and that a precursor of these amino acids e.g. shikimic acid is then diverted to the synthesis of new compounds.

This view of the origin of aromatic compounds puts them in the same class as organic acids, fats and polysaccharides as products of what have been termed metabolic shunts (200). Physiologically the model proposed above requires that the ratio of metabolic product to mycelial nitrogen should increase with increasing glucose concentration at least over a certain range. This has not been tested; the ratio of product to dry weight of mycelium (213) is less informative because of the likelihood that carbon components of the mycelium may also be products of metabolic shunts.


It has often been suggested that fungal quinones play a specific role in respiration and the hypothesis is an attractive one. The frequent simultaneous occurrence of the reduced and oxidized forms in one organism suggests that the system could transport electrons and so function in terminal respiration. However, satisfactory proof of such a role in the intact cell has not been brought forward. Addition of quinones to a culture often elicits a respiratory response (214, 389) but this can only be taken as an indication that the compound could function, not that it does in fact function.

A respiratory role for the polyphenol oxidases of higher plants has often been proposed and the question is still open. In *Glomerella cingulata* it appears that tyrosinase cannot be an essential respiratory enzyme inasmuch as it is not detectable during the growth phase (500).

A pyridine nucleotide menadione reductase occurs in *Aspergillus niger* (561) and a pyridine nucleotide-quinone reductase in other fungi (565). In principle either of these could participate in a

Polyphenol oxidases in the fungi are often present only at particular times in the culture cycle especially during autolysis (188-500) or only in particular morphological structures (342-344). The medium used affects the production of the enzymes both positively and negatively. Laccase formation by *Polyporus versicolor* is increased non specifically by constituents of the medium and specifically by certain aromatic compounds which probably act as inducers (189-190). Both carbon source and growth temperature affect enzyme synthesis in *Glomerella cingulata* (499). Negative effects of an interesting type have been found in studies on *Neurospora crassa* (287) in a medium containing the usual level of sulfate a sulfur-containing dialyzable tyrosinase inhibitor is formed the result being that the conventional assay detects tyrosinase only in cultures grown in a low sulfate medium. This finding requires that earlier negative reports be re-evaluated.

The tannins are esters of polyhydroxy phenols with aromatic acids and are hydrolyzed by tannase e.g.




The enzyme an esterase also hydrolyzes other similar esters (498). The tannase of *Aspergillus niger* is inducible (322) its properties and purification are summarized by Dyckerhoff and Armbruster (175) and partial purification has been reported (519). Although most studies have been made on the tannase of *A. niger* the enzyme has been reported in a few other fungi (36-448-574).

The Significance of Aromatic Compounds Some of the aromatic compounds bulk rather large in the economy of the organism amounting to an appreciable fraction of the glucose consumed (140) and of the dry weight produced (128). Those which have been studied in relation to the history of the culture appear to be first formed and then utilized (93-95) to be more than inactive waste products of metabolism.

Possibly the most likely metabolic significance of these compounds is as products of an unbalanced nutrition and metabolism. Most of them have been isolated from fungi grown on media very high in carbon thus the glucose Czapek Dox medium often used has a carbon-nitrogen ratio of 60. Under these conditions it is not surprising

that carbon may flow into new metabolic pools simply because of an insufficiency of nitrogen. We may visualize this type of process schematically



If compound D for example cannot be synthesized because of a relative nitrogen deficiency enzymatic equilibria and factors of permeability and solubility will determine which of the potential alternative pathways is followed. In the specific instance of the benzenoid compounds and their oxidation products it is imaginable that the synthesis of aromatic amino acids stops because nitrogen is no longer available and that a precursor of these amino acids e.g. shikimic acid is then diverted to the synthesis of new compounds.

This view of the origin of aromatic compounds puts them in the same class as organic acids, fats and polysaccharides as products of what have been termed metabolic shunts (200). Physiologically the model proposed above requires that the ratio of metabolic product to mycelial nitrogen should increase with increasing glucose concentration at least over a certain range. This has not been tested; the ratio of product to dry weight of mycelium (243) is less informative because of the likelihood that carbon components of the mycelium may also be products of metabolic shunts.

It has often been suggested that fungal quinones play a specific role in respiration and the hypothesis is an attractive one. The frequent simultaneous occurrence of the reduced and oxidized forms in one organism suggests that the system could transport electrons and so function in terminal respiration. However satisfactory proof of such a role in the intact cell has not been brought forward. Addition of quinones to a culture often elicits a respiratory response (214-389) but this can only be taken as an indication that the compound could function, not that it does in fact function.

A respiratory role for the polyphenol oxidases of higher plants has often been proposed and the question is still open. In *Glomerella cingulata* it appears that tyrosinase cannot be an essential respiratory enzyme inasmuch as it is not detectable during the growth phase (500).


A pyridine nucleotide menadione reductase occurs in *Aspergillus niger* (564) and a pyridine nucleotide quinone reductase in other fungi (565). In principle either of these could participate in a

coupling mechanism between substrate dehydrogenation and a polyphenol oxidase system transferring electrons to oxygen

Many of the quinonoid compounds of fungi are more or less inhibitory to other organisms and it has been suggested (341) that this property may confer an ecological advantage. Whether highly reactive compounds like the quinones could persist long enough in soil to inhibit competing bacteria and fungi is however somewhat doubtful.

4 DERIVATIVES OF PYRONE

Several metabolic products may be considered together as derivatives of 1,4-pyrone and is related structurally to the pyranose form of glucose. Kojic acid is typical.




Kojic acid was first discovered as a product of *Aspergillus oryzae* by Saito (452). Its structure was determined by Yabuta (568). Many species of *Aspergillus* but only one of *Penicillium* produce it in culture (71, 227, 242, 303). Kojic acid is weakly antibacterial (381).

Kojic acid is produced from most common carbon sources (71, 311) although strains differ somewhat in this regard (404). The number of compounds providing carbon for kojic acid synthesis is less for a preformed mycelium than for growing cells (504).

The data of Figure 8 show that as would be expected there is over a certain range an inverse relation between nitrogen supplied to a culture and kojic acid production per unit mycelial weight. These data can best be interpreted as showing that kojic acid synthesis per cell is maximal when growth is limited by a nitrogen deficiency in accordance with the principle that metabolites accumulate when carbon is supplied in excess of that which can be incorporated into cellular material.

The formation of kojic acid is sensitive to pH; the yield from a preformed mycelium declines virtually to zero as the pH is raised from 2.2 to 6.0 (319). Ethylene chlorhydrin increases the formation of

Figure 8 Growth and kojic acid formation by *Aspergillus flavus* as functions of nitrogen supply. Drawn from tabular data of May Moyer Wells and Herrick (373)



kojic acid by *Aspergillus flavus* but the effect appears to be on the entire metabolism of the organism rather than specifically on kojic acid synthesis (374)

The very high yields of kojic acid—up to 57 per cent of the glucose consumed by a growing culture (373) and up to 78 per cent of that used by preformed mycelium (319)—suggest that the acid may be formed directly from glucose. This is borne out by the isotope studies of Arnstein and Bentley (18). Degradation of kojic acid derived from glucose-1- C^{14} shows that the bulk of the acid is formed from glucose without splitting of the carbon chain. The occurrence of small amounts of C^{14} in other positions is adequately explained by the postulate that a minor fraction of the carbon of kojic acid is derived from compounds of the phosphogluconate oxidation respiratory pathway (19, 283).

The details of the conversion of glucose to kojic acid are not known. Arnstein and Bentley (18) discuss some possibilities. The fact that a preformed mycelium does not require added phosphate for kojic acid synthesis (242) is irrelevant to the problem of whether or not phosphorylative steps are involved in synthesis since the endogenous phosphate reserves of the cell would without question be adequate for a cyclic process.


Patulin is a pyrone derivative the structure of which is largely known (181, 563). It has been isolated several times independently under different names—clavacin, claviformin, expansin and clavatin—from species of *Penicillium* and *Aspergillus*; its history and synonymy

are reviewed by Raistrick (430) As mentioned earlier the effects of metals on patulin production indicate that it is biosynthetically related to some of the aromatic compounds of fungi

Alternariol is a dibenzo 1,2 pyrone formed by *Alternaria tenuis* after long-continued growth on a glucose medium (435) There is no reason to believe that it is metabolically related to the 1,4 pyrones


5 DERIVATIVES OF FURAN

A number of fungal metabolites are structurally furan derivatives Most of these are derivatives of tetronic acid and have in common the structure



One organism *Penicillium charlesii* forms five related compounds of this type γ methyltetronic acid carlic acid carlosic acid carolinic acid and carolic acid (140-143) Other compounds of the same general class are terrestric acid of *Penicillium terrestre* (82) dehydro carolic acid of *P. cinerascens* (94) and penicillic acid of *Penicillium* spp and *Aspergillus* spp (6, 78, 227)

L Ascorbic acid well known as a product of higher plants has been isolated from the culture filtrate of *Aspergillus niger* (223) Its relation to the tetronic acids is shown by its structure



Studies on higher plants indicate that ascorbic acid is synthesized from glucose without rupture of the carbon chain and that L galactono- γ lactone is an important intermediate (367) It is also striking in view of the formation of oxalic acid by *Aspergillus niger* that oxalic acid is an end product of ascorbate metabolism in the guinea pig (99) Reports of ascorbic acid in fungi other than *Aspergillus niger* are summarized by Van Lanen and Tanner (531)

An ascorbic acid oxidase of the higher plant type containing copper has not been reported from fungi *Myrothetium verrucaria*


spores oxidize ascorbate but the reaction is believed to be mediated by a new enzyme probably located on the spore surface (365-366)

Two other derivatives of furan—not derivatives however of tetronic acid—are known: 2-hydroxy-5-furan-carboxylic acid from *Aspergillus* spp. (496-497) and zymonic acid from yeasts (493)

The thiophen or thiofuran ring differs from the furan in that a sulphur atom replaces the oxygen. A single compound of this type, junipal, is known in the fungi from *Daedalea juniperina* (69)

6 BIOLOGICAL METHYLATION


The transfer of intact methyl groups from one molecule to another has been most intensively studied in animal materials. The synthesis of methionine may be taken as an example



In this reaction details of which are neglected in the above overall formulation, betaine functions as the methyl donor. Methionine can in turn donate a methyl group to other molecules, e.g. in the stepwise methylation of aminoethanol to choline.

Studies on *Neurospora crassa* mutants which require methionine show that the final step in methionine synthesis is the conversion of homocysteine to methionine as in Equation 16 above (285). The methyl donor in the fungus system is not known.

In a particularly important application of biochemical genetics, Horowitz (284) showed that *Neurospora crassa* synthesizes choline by the stepwise methylation of aminoethanol that is



Here it is to be presumed that methionine or formate is the source

of methyl groups. The ultimate source for those fungi—perhaps all naturally occurring fungi—which do not require preformed methyl groups in the medium is of course carbohydrate that is most or all fungi can synthesize methyl groups from the products of glucose metabolism.

Early in the nineteenth century it was recognized that an arsenical poisoning of humans by volatile compounds is associated with the action of fungi on wallpaper or other building materials containing arsenical pigments. The history of this problem is reviewed by Challenger (117); the toxic material was finally identified as trimethylarsine (121). The ability of certain fungi to form methylated compounds of arsenic and other elements has since been investigated vigorously by the group at the University of Leeds. Some examples of the products identified are listed in Table 5; these and others are reviewed

Table 5 Methylated Products of Fungi

Substrate	Product	Organism
Na methylarsenate	Trimethylarsine	<i>Scopulariopsis brevicaulis</i> (121)
Na cacodylate	Trimethylarsine	<i>S. brevicaulis</i> (121)
Arsenous oxide	Trimethylarsine	<i>S. brevicaulis</i> (121)
Diethylarsonic acid	Methyldiethylarsine	<i>S. brevicaulis</i> (120)
H ₂ tellurite	Dimethyl telluride	<i>S. brevicaulis</i> , <i>Penicillium</i> spp. (62)
Na selenate	Dimethyl selenide	<i>Aspergillus niger</i> (170) <i>Penicillium</i> spp. (62) <i>Scopulariopsis brevicaulis</i> (123)
Na selenite	Dimethyl selenide	<i>S. brevicaulis</i> (123)
Ethyl disulfide	Methyl ethyl sulfide	<i>S. brevicaulis</i> (124)
Inorganic sulfate	Methyl mercaptan dimethyl sulfide	<i>Schizophyllum commune</i> (76, 119)

by Challenger (118). It should be noted that the methods used permit only identification of the final product. Obviously many steps in addition to methylation must be assumed in the conversion of the highly oxidized selenate ion for example to the reduced dimethyl selenide.

Evidence is now at hand that these products do in fact arise by transmethylation. Isotope experiments (122-170) show that the methyl groups of dimethyl selenide and trimethylarsine originate from methionine in *Scopulariopsis brevicaulis* and *Aspergillus niger*. Betaine does not serve as a methyl donor; carbon from formate and choline enters the system probably through methionine.

7 BIOLOGICAL CHLORINATION

Chloride ion is commonly present in culture media simply as an accompaniment of essential ions. With the discovery of geodin and erdin in *Aspergillus terreus* (433-434) it was realized that chlorine may be incorporated into organic compounds by fungi. Subsequent investigations have shown many chlorinated metabolites; a partial list includes

- 1 *p*-Methoxy tetrachlorophenol of *Drosophila subatrata* (8)
- 2 Chlorinated depsidones e.g. nidulin (Table 4). Chlorinated depsidones are also known in the lichens (24); they are presumably synthesized by the fungus component.
- 3 Griseofulvin, a complex benzenoid compound of uncertain structure produced by *Penicillium* spp. (96, 256-411)
- 4 Chlortetracycline (aureomycin), an antibiotic of *Streptomyces aureofaciens* (437)
- 5 Chloramphenicol (chloromycetin) (Table 4). Here the chlorine is attached to a side chain carbon; chloramphenicol is also unusual in its nitrobenzene structure.
- 6 Nalgiolaxin, a chlorinated anthraquinone of *Penicillium nalgiovenus* occurring in association with its dechloro analogue nalgiolaxin (436)
- 7 Caldariomycin of *Caldariomyces fumago* (*Fumago vagans*) probably a cyclopentane with two chlorine atoms attached to a ring carbon (141). Terrein of *Aspergillus terreus*, not a chlorine compound, is the only other cyclopentane derivative known in fungi (32, 255)

In a survey of 139 fungi it was found that strains of *Abisidia spinosa*, *Alternaria tenuis*, *Aspergillus* spp., *Caldariomyces fumago*, *Penicillium* spp., *Syncephalastrum racemosum* and *Trichoderma viride* incorporate 15 per cent or more of the available chlorine into organic combinations (141).

Frequently a compound and its dechloro derivative occur together in one organism, e.g. griseofulvin and dechlorogriseofulvin in *Penicillium* spp. (362). The antibiotics tetracycline and chlortetracycline are produced by different species of *Streptomyces*; the related oxytetracycline by a third (61). The substitution of bromide for chloride in the medium has specific effects of two kinds: either the organism forms the bromo analogue of a chlorine compound (363) or chlorination is suppressed and the dechloro derivative is formed (468). The incorporation of chlorine into tetracycline is also suppressed by copper-com-


plexing agents e.g. thiourea (168) which are known to interfere with iodination of tyrosine in the thyroid (551) presumably the chlorination step requires copper

8 PENICILLIN

The discovery by Fleming (197) of an organism producing penicillin initiated a new field of medical and biological investigation the study of the formation and clinical use of antibiotics. The history of the exploitation of Fleming's discovery has been told elsewhere (1) in general although extensive application of our knowledge of fungal physiology was essential to the development of methods of production the physiological studies were less dramatic than the genetic and few new principles of physiology or biochemistry have emerged from the work on penicillin and other antibiotics.

The penicillins are produced by several different fungi including *Penicillium* spp. *Aspergillus* spp. and a thermophilic organism tentatively identified as *Malbranchea pulchella* (441). Strains of the *Penicillium chrysogenum* series including *P. notatum* are the most active and supply the penicillin of commerce.

The penicillins are alike in their basic chemical structure but differ in the side chain (R in the formula)



The natural penicillins whose structures have been determined include

- 1 2-Pentenylpenicillin (penicillin I penicillin F)
- 2 Benzylpenicillin (penicillin II penicillin G)
- 3 *p*-Hydroxybenzylpenicillin (penicillin III penicillin V)
- 4 *n*-Amylpenicillin (dihydro penicillin F)
- 5 *n*-Heptylpenicillin (penicillin IV penicillin K)
- 6 4-Amino-4-carboxy *n*-butylpenicillin (cephalosporin N) (393)

It is likely that other penicillins formed only in small amounts have not been identified. Benzylpenicillin now available in crystalline form is the penicillin of medical practice.

Extensive information on the metabolism of penicillin-producing fungi reviewed by Chain (115) and Perlman (417) is available but it must be said that there is no correlation of any other metabolic

process with penicillin formation. It is possible to specify fairly exactly the optimal conditions for penicillin formation by a given strain but not possible to define the way in which these conditions act.

The metabolic changes which take place during penicillin production are conveniently divided into three phases: an initial phase of rapid growth and metabolic activity with assimilation of the more available nutrients; a second phase of slower growth with carbohydrate utilization; and a third phase of autolysis and low metabolic activity. These are illustrated in Figure 9. Penicillin formation begins late in the first phase and is most vigorous during the second. Rates differ but in broad outline this pattern is followed by different strains in both complex and synthetic media with various carbon sources and under diverse conditions of aeration (31, 209, 217, 239, 300, 324). Lactose or other slowly utilizable carbohydrates yield both a favorable rate of growth and a favorable pH for penicillin formation (385, 484).

Penicillin production is also directly affected by aeration (483). The metal requirement for maximal penicillin formation by a strain of




Figure 9 Metabolic changes accompanying penicillin formation by *Penicillium notatum* in shaken flasks in a corn steep liquor lactose medium. Redrawn from Koffler, Emerson, Perlman, and Buttriss (33) by permission of the Williams and Wilkins Company.

P. chrysogenum is somewhat higher particularly as to iron and phosphorus than that for growth (301)

Corn steep liquor a complex mixture of organic substances derived by bacterial action on maize (339) probably has more than one role in penicillin production its principal function however appears to be as a source of β phenylethylamine and related compounds (375) which are specific precursors of benzylpenicillin In the biosynthesis of mixed penicillins phenylacetic acid and its derivatives shift the balance of natural penicillins toward benzylpenicillin (273) Similarly *p* hydroxyphenylacetate increases the relative amount of *p* hydroxy benzylpenicillin Although only a small fraction of the phenylacetate supplied is incorporated into the molecule (295) isotope studies have shown that it is incorporated without dilution i.e. that the intact benzyl group enters as such into penicillin (38 240 467)

Not only are close relatives of phenylacetic acid incorporated into benzylpenicillin but many other substituted acetic acids can serve as precursors for new biosynthetic penicillins (37) Fatty acids especially those which are not rapidly broken down by the fungus are incorporated into both natural and new penicillins (380 516)

The sulfur of penicillin can be supplied by inorganic sulfate sulfur of methionine cysteine cystine or glutathione is however utilized in preference to that of inorganic sulfate (490)

The origin of the main part of the penicillin molecule has been clarified by Arnstein and Grant (22) Using C^{14} labeled valine and triply (N^{15} C^{14} S^{35}) labeled cystine they have shown that (1) cystine—probably after conversion to cysteine—is incorporated directly into the penicillin molecule and (2) the remaining five carbons are supplied by valine This is consistent with earlier work on the incorporation of small molecules into penicillin (368 489 517) The molecule of benzylpenicillin may be viewed therefore as made of three metabolic units one formed directly from phenylacetic acid one directly from cystine (probably via cysteine) and one supplied directly or indirectly by valine The steps by which these structures are combined remain for future investigation It is to be expected that energy is necessary for penicillin formation from these units and the inhibitory effect of 2,4-dinitrophenol suggests that the energy is derived from oxidative phosphorylations (419)

The extensive studies on penicillin precursors have made it clear that although a precursor may be specifically and directly incorporated into a molecule provision of the precursor does not necessarily increase the final yield of product It is only when the formation of the precursor is so slow as to be rate limiting to the total process that

such increases in yield are found. This general principle obviously applies to all biosynthetic studies with putative precursors.

Penicillinase is produced by many bacteria and probably by actinomycetes; it is not certain whether any fungus forms it (116). Its action is to convert penicillins to penicilloic acids by hydrolysis and opening of the β lactam ring.

Penicillin amidase found in the mycelium of *Penicillium chrysogenum* and *Aspergillus oryzae* splits off the benzyl side chain from benzylpenicillin like penicillinase; it attacks a peptide (amide) linkage (459).

BIBLIOGRAPHY

- 1 Abraham E. P. et al. 1949. In H. W. Florey et al. (eds) *Antibiotics* Vol. 9 p. 631-671. London: Oxford Univ. Press.
- 2 Acklin O. 1979. *Biochem. J.* 170: 45-54.
- 3 Akamatsu S. 1973. *Biochem. J.* 134: 186-187.
- 4 Allsopp A. 1937. *New Phytologist* 36: 327-336.
- 5 Allsopp A. 1940. *J. Exp. Botany* 1: 71-81.
- 6 Alsberg C. L. and O. F. Black. 1913. U.S. Dept. Agr. Bur. Plant Ind. Bull. 10: 1-48.
- 7 Ammon R. and M. Jaarma. 1950. In J. B. Sumner and K. Myrback (eds) *The Enzymes* Vol. 1 (1) p. 590-41. New York: Academic Press.
- 8 Anselmi M. 1950. *J. Am. Chem. Soc.* 72: 7943.
- 9 Anselmi M. 1953. *Arch. Biochem. Biophys.* 43: 17-13.
- 10 Anselmi M. 1955. *Science* 121: 607-608.
- 11 Anselmi M. and M. P. Cohen. 1954. *J. Biol. Chem.* 72: 318-326.
- 12 Anselmi M., A. Hervey, F. Kavanagh, J. Polatinck, and W. J. Robbins. 1948. *Proc. Natl. Acad. Sci. U.S.A.* 34: 498-502.
- 13 Angeletti A. and C. F. Cerruti. 1930. *Ann. chim. appl.* 9: 424-433.
- 14 Anslow W. K. and H. Raistrick. 1938a. *Biochem. J. (London)* 3: 687-696.
- 15 Anslow W. K. and H. Raistrick. 1938b. *Biochem. J. (London)* 3: 803-806.
- 16 Anslow W. K. and H. Raistrick. 1938c. *Biochem. J. (London)* 3: 2280-2289.
- 17 Armstrong G. I. 1929. *Protoplasma* 8: 222-250.
- 18 Arnstein H. R. V. and R. Bentley. 1953. *Biochem. J. (London)* 54: 493-508.
- 19 Arnstein H. R. V. and R. Bentley. 1956. *Biochem. J. (London)* 6: 403-411.
- 20 Arnstein H. R. V. and A. H. Cook. 1947. *J. Chem. Soc. (London)* 1947: 1021-1028.
- 21 Arnstein H. R. V., A. H. Cook, and M. S. Lacey. 1946. *Bull. J. Exp. Pathol.* 27: 319-325.
- 22 Arnstein H. R. V. and P. T. Grant. 1954. *Biochem. J. (London)* 57: 33-39.
- 23 Asahina Y. and S. Shibata. 1939. *Ber. deut. chem. Ges.* 72B: 1531-1533.
- 24 Asahina Y. and S. Shibata. 1953. *Chemistry of Lichen Substances*. Tokyo: Japan Society for the Promotion of Science, pp. 40.
- 25 Aulén Erdman G. 1951. *Acta Chem. Scand.* 5: 301-315.
- 26 Baba S. and K. Sakaguchi. 1941. *Bull. Agr. Chem. Soc. Japan* 18: 93-91.
- 27 Bach D. and J. Fournier. 1935. *Compt. rend. (Paris)* 201: 98-984.
- 28 Baker G. E. 1915. *Mycologia* 27: 580-600.

P. chrysogenum is somewhat higher particularly as to iron and phosphorus than that for growth (301)

Corn steep liquor a complex mixture of organic substances derived by bacterial action on maize (339) probably has more than one role in penicillin production its principal function however appears to be as a source of β phenylethylamine and related compounds (372) which are specific precursors of benzylpenicillin In the biosynthesis of mixed penicillins phenylacetic acid and its derivatives shift the balance of natural penicillins toward benzylpenicillin (273) Similarly *p* hydroxyphenylacetate increases the relative amount of *p* hydroxy benzylpenicillin Although only a small fraction of the phenylacetate supplied is incorporated into the molecule (292) isotope studies have shown that it is incorporated without dilution i.e. that the intact benzyl group enters as such into penicillin (38 210 467)

Not only are close relatives of phenylacetic acid incorporated into benzylpenicillin but many other substituted acetic acids can serve as precursors for new biosynthetic penicillins (37) Fatty acids especially those which are not rapidly broken down by the fungus are incorporated into both natural and new penicillins (380 516)

The sulfur of penicillin can be supplied by inorganic sulfate sulfur of methionine cysteine cystine or glutathione is however utilized in preference to that of inorganic sulfate (490)

The origin of the main part of the penicillin molecule has been clarified by Arnstein and Crant (22) Using C^{14} labeled valine and triply (N^{15} C^{14} S^{35}) labeled cystine they have shown that (1) cystine—probably after conversion to cysteine—is incorporated directly into the penicillin molecule and (2) the remaining five carbons are supplied by valine This is consistent with earlier work on the incorporation of small molecules into penicillin (368 489 517) The molecule of benzylpenicillin may be viewed therefore as made of three metabolic units one formed directly from phenylacetic acid one directly from cystine (probably via cysteine) and one supplied directly or indirectly by valine The steps by which these structures are combined remain for future investigation It is to be expected that energy is necessary for penicillin formation from these units and the inhibitory effect of 2,4-dinitrophenol suggests that the energy is derived from oxidative phosphorylations (119)

The extensive studies on penicillin precursors have made it clear that although a precursor may be specifically and directly incorporated into a molecule provision of the precursor does not necessarily increase the final yield of product It is only when the formation of the precursor is so slow as to be rate limiting to the total process that

such increases in yield are found. This general principle obviously applies to all biosynthetic studies with putative precursors.

Penicillinase is produced by many bacteria and probably by actinomycetes; it is not certain whether any fungus forms it (116). Its action is to convert penicillins to penicilloic acids by hydrolysis and opening of the β lactam ring.

Penicillin amidase found in the mycelium of *Penicillium chrysogenum* and *Aspergillus oryzae* splits off the benzyl side chain from benzylpenicillin. Like penicillinase it attacks a peptide (amide) linkage (159).

BIBLIOGRAPHY

- 1 Abraham E. I. et al. 1949. In H. W. Florey et al. (eds) *Antibiotics* Vol. 2 p. 651-671. London: Oxford Univ. Press.
- 2 Acklin O. 1909. *Biochem. Z.* 204: 23-24.
- 3 Akamatsu S. 1973. *Biochem. Z.* 147: 186-18.
- 4 Allsopp A. 1937. *New Phytologist* 36: 3: 3-6.
- 5 Allsopp A. 1940. *J. Exp. Botany* 1: 71-81.
- 6 Alsberg C. L. and O. F. Black. 1913. *U. S. Dept. Agr. Bur. Plant Ind. Bull.* 746: 1-19.
- 7 Ammon R. and M. Jaariva. 1940. In J. B. Sumner and K. Myrback (eds) *The Enzymes* Vol. 1 (1): p. 390-41. New York: Academic Press.
- 8 Anchel M. 1940. *J. Am. Chem. Soc.* 71: 2943.
- 9 Anchel M. 1953. *Arch. Biochem. Biophys.* 45: 127-15.
- 10 Anchel M. 1940. *Science* 121: 607-608.
- 11 Anchel M. and M. P. Cohen. 1954. *J. Biol. Chem.* 72: 318-36.
- 12 Anchel M., A. Hervey, F. Kavanagh, J. Polatnick and W. J. Robbins. 1948. *Proc. Natl. Acad. Sci. U. S. A.* 34: 498-502.
- 13 Angeletti A. and C. F. Cerruti. 1930. *Ann. chim. appl.* 20: 4: 1-453.
- 14 Anslow W. K. and H. Raistrick. 1938a. *Biochem. J. (London)* 32: 687-696.
- 15 Anslow W. K. and H. Raistrick. 1938b. *Biochem. J. (London)* 32: 803-806.
- 16 Anslow W. K. and H. Raistrick. 1938c. *Biochem. J. (London)* 32: 2287-2289.
- 17 Armstrong G. I. 1929. *Protoplasma* 8: 222-260.
- 18 Arnstein H. R. V. and R. Bentley. 1953. *Biochem. J. (London)* 51: 493-508.
- 19 Arnstein H. R. V. and R. Bentley. 1954. *Biochem. J. (London)* 51: 509-516.
- 20 Arnstein H. R. V. and R. Bentley. 1954. *Biochem. J. (London)* 51: 517-522.
- 21 Arnstein H. R. V. and R. Bentley. 1954. *Biochem. J. (London)* 51: 523-528.
- 22 Arnstein H. R. V. and A. H. Cook. 1947. *J. Chem. Soc. (London)* 194: 1071-1078.
- 23 Arnstein H. R. V., A. H. Cook and M. S. Lacey. 1946. *Brit. J. Exp. Pathol.* 7: 319-335.
- 24 Arnstein H. R. V. and P. T. Grant. 1954. *Biochem. J. (London)* 51: 353-359.
- 25 Asahina Y. and S. Shibata. 1939. *Ber. deut. chem. Ges.* 72B: 1531-1533.
- 26 Asahina Y. and S. Shibata. 1953. *Chemistry of Lichen Substances*. Tokyo: Japan Society for the Promotion of Science. pp. 240.
- 27 Aulic Erdmann G. 1941. *Acta Chem. Scand.* 5: 391-313.
- 28 Baba S. and K. Sakaguchi. 1940. *Bull. Agr. Chem. Soc. Japan* 18: 93-91.
- 29 Bach D. and J. Fournier. 1930. *Compt. rend. (Paris)* 201: 982-981.
- 30 Baker G. E. 1915. *Mycologia* 37: 587-600.

P. chrysogenum is somewhat higher particularly as to iron and phosphorus, than that for growth (301)

Corn steep liquor a complex mixture of organic substances derived by bacterial action on maize (339) probably has more than one role in penicillin production its principal function however appears to be as a source of β phenylethylamine and related compounds (375) which are specific precursors of benzylpenicillin In the biosynthesis of mixed penicillins phenylacetic acid and its derivatives shift the balance of natural penicillins toward benzylpenicillin (273) Similarly *p* hydroxyphenylacetate increases the relative amount of *p* hydroxy benzylpenicillin Although only a small fraction of the phenylacetate supplied is incorporated into the molecule (295) isotope studies have shown that it is incorporated without dilution i.e. that the intact benzyl group enters as such into penicillin (38 210 467)

Not only are close relatives of phenylacetic acid incorporated into benzylpenicillin but many other substituted acetic acids can serve as precursors for new biosynthetic penicillins (37) Fatty acids especially those which are not rapidly broken down by the fungus are incorporated into both natural and new penicillins (380 516)

The sulfur of penicillin can be supplied by inorganic sulfate sulfur of methionine cystine cysteine or glutathione is however utilized in preference to that of inorganic sulfate (490)

The origin of the main part of the penicillin molecule has been clarified by Arnstein and Grant (22) Using C^{14} labeled valine and triply (N^{15} C^{14} S^3) labeled cystine they have shown that (1) cystine—probably after conversion to cysteine—is incorporated directly into the penicillin molecule and (2) the remaining five carbons are supplied by valine This is consistent with earlier work on the incorporation of small molecules into penicillin (358 489 517) The molecule of benzylpenicillin may be viewed therefore as made of three metabolic units one formed directly from phenylacetic acid one directly from cystine (probably via cysteine) and one supplied directly or indirectly by valine The steps by which these structures are combined remain for future investigation It is to be expected that energy is necessary for penicillin formation from these units and the inhibitory effect of 2,4-dinitrophenol suggests that the energy is derived from oxidative phosphorylations (119)

The extensive studies on penicillin precursors have made it clear that although a precursor may be specifically and directly incorporated into a molecule provision of the precursor does not necessarily increase the final yield of product It is only when the formation of the precursor is so slow as to be rate limiting to the total process that

such increases in yield are found. This general principle obviously applies to all biosynthetic studies with putative precursors.

Penicillinase is produced by many bacteria and probably by actinomycetes; it is not certain whether any fungus forms it (116). Its action is to convert penicillins to penicilloic acids by hydrolysis and opening of the β lactam ring.

Penicillin amidase found in the mycelium of *Penicillium chrysogenum* and *Aspergillus oryzae* splits off the benzyl side chain from benzylpenicillin like penicillinase; it attacks a peptide (amide) link age (459).

BIBLIOGRAPHY

- 1 Abraham E P et al. 1949. In H W Florey et al (eds) *Antibiotics* Vol 9 p 631-671. London: Oxford Univ Press.
- 2 Acklin O. 1939. *Biochim Z* 204: 23-24.
- 3 Akamatsu S. 1953. *Biochem Z* 147: 186-187.
- 4 Allsopp A. 1937. *New Phytologist* 26: 37-38.
- 5 Allsopp A. 1939. *J Exp Botany* 1: 71-81.
- 6 Ahberg C L and O F Black. 1913. U S Dept Agr Bur Plant Ind Bull 270: 1-48.
- 7 Ammon R and M Jaarola. 1950. In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 390-411. New York: Academic Press.
- 8 Anchel M. 1951. *J Am Chem Soc* 73: 7943.
- 9 Anchel M. 1953. *Arch Biochem Biophys* 43: 17-13.
- 10 Anchel M. 1951. *Science* 121: 607-608.
- 11 Anchel M and M P Cohen. 1954. *J Biol Chem* 79: 318-36.
- 12 Anchel M, A Hervey, F Kavanagh, J Polatnick and W J Robbins. 1948. *Proc Natl Acad Sci U S A* 34: 498-502.
- 13 Angeletti A and C F Cerruti. 1950. *Ann chim appl* 20: 474-433.
- 14 Anslow W K and H Raistrick. 1938a. *Biochem J (London)* 32: 687-696.
- 15 Anslow W K and H Raistrick. 1938b. *Biochem J (London)* 32: 803-806.
- 16 Anslow W K and H Raistrick. 1938c. *Biochem J (London)* 32: 278-279.
- 17 Armstrong G I. 1939. *Protoplasma* 8: 227-260.
- 18 Arnstein H R V and R Bentley. 1953. *Biochem J (London)* 54: 493-508.
- 19 Arnstein H R V and R Bentley. 1956. *Biochem J (London)* 60: 403-411.
- 20 Arnstein H R V and A H Cook. 1947. *J Chem Soc (London)* 1947: 1071-1078.
- 21 Arnstein H R V, A H Cook and M S Lacey. 1946. *Proc J Exp Pathol* 27: 349-351.
- 22 Arnstein H R V and P T Grant. 1951. *Biochem J (London)* 57: 353-359.
- 23 Asahina Y and S Shibata. 1939. *Ber deut chem Ges* 7 B: 1531-1533.
- 24 Asahina Y and S Shibata. 1953. *Chemistry of Lichen Substances*. Tokyo: Japan Society for the Promotion of Science. pp 40.
- 25 Auldin Erdman G. 1931. *Acta Chem Scand* 5: 301-311.
- 26 Baba S and K Sakaguchi. 1940. *Bull Agr Chem Soc Japan* 18: 95-96.
- 27 Bach D and J Fournier. 1951. *Compt rend (Paris)* 201: 982-984.
- 28 Baker G E. 1915. *Mycologia* 37: 582-600.

- 28a Ball S G J Bessell and A Mortimer 1957 *J Gen Microbiol* 17 96-103
- 29 Barnett H L V G Lilly and R F Krause 1956 *Science* 123 141
- 30 Barron E S G and F Ghirelli 1953 *Biochim et Biophys Acta* 12 239-249
- 31 Bartholomew W H F O Karow M R Spar and R H Wilhelm 1950 *Ind Eng Chem* 42 1810-1815
- 32 Barton D H R and F Miller 1955 *J Chem Soc (London)* 1955 1028-1029
- 33 de Bary A 1886 *Botan Ztg* 41 393-401
- 34 Bauer R 1938 *Zentr Bakteriell Parasitenk Abt II* 98 153-167
- 35 Bavendamm W 1927 *Ber deut bot Ges* 45 357-367
- 36 Bazzigher G 1955 *Phytopathol Z* 23 265-282
- 37 Behrens O K 1949 In H T Clarke J R Johnson and R Robinson (eds) *The Chemistry of Penicillin* p 657-679 Princeton N J Princeton Univ Press
- 38 Behrens O K et al 1948 *J Biol Chem* 175 765-769
- 39 Belin P 1926 *Bull soc chim Biol* 8 1081-1102
- 40 Bendz G 1951 *Acta Chem Scand* 5 489-490
- 41 Bentley R and C P Thiessen 1957 *J Biol Chem* 226 673 687 689-690 703 709
- 42 Bergel F A L Morrison A R Moss and H Rinderknecht 1941 *J Chem Soc (London)* 1941 415-421
- 43 Bernhard K 1948 *Cold Spring Harbor Symposia Quant Biol* 13 26-28
- 44 Bernhard K and H Albrecht 1947 *Helv Chim Acta* 30 67-632
- 45 Bernhard K and H Albrecht 1948 *Helv Chim Acta* 31 2214-2220
- 46 Bernhauer K 1926a *Biochem Z* 172 313-323
- 47 Bernhauer K 1926b *Biochem Z* 172 321-349
- 48 Bernhauer K 1928 *Biochem Z* 197 218-286
- 49 Bernhauer K 1931 *Ergeb Enzymforsch* 3 185-216
- 50 Bernhauer K 1933 *Ergeb Enzymforsch* 9 297-360
- 51 Bernhauer K and N Bockl 1932 *Biochem Z* 253 16-24
- 52 Bernhauer K N Bockl and H Siebenauger 1937 *Biochem Z* 253 37-41
- 53 Bernhauer K and B Görlisch 1935 *Biochem Z* 280 394-395
- 54 Bernhauer K J Rauch and J N Miksch 1949 *Biochem Z* 350 178-183
- 55 Bernhauer K and Z Scheuer 1932 *Biochem Z* 253 11-15
- 56 Bernhauer K and H Thole 1936 *Biochem Z* 287 167-171
- 57 Bernhauer K and H H Waelsch 1932 *Biochem Z* 249 223-226
- 58 Bernhauer K and H Wolf 1928 *Z physiol Chem Hoppe Seyler's* 177 210-219
- 58a Bertrand D and A De Wolf 1955 *Compt Rend (Paris)* 241 1877-1880
- 59 Biale J B and A D Shepherd 1941 *Am J Botany* 28 263-270
- 60 Bier M 1955 In S P Colowick and N O Kaplan (eds) *Methods in Enzymology* Vol 1 p 627-647 New York Academic Press
- 61 Binkley S B 1955 *Ann Rev Biochem* 24 596-626
- 61a Birch A J and F W Donovan 1954 *Chem & Ind* 1954 1047-1048
- 62 Bird M L and F Challenger 1939 *J Chem Soc (London)* 1939 163-168
- 63 Birkinshaw J H 1953 *Ann Rev Biochem* 22 371-398
- 64 Birkinshaw J H A Bracken and W P K Findlay 1944 *Biochem J (London)* 38 151-157
- 65 Birkinshaw J H A Bracken E N Morgan and H Raistrick 1948 *Biochem J (London)* 43 216-223
- 66 Birkinshaw J H A Bracken and H Raistrick 1945 *Biochem J (London)* 37 726-728
- 67 Birkinshaw J H A Bracken and H Raistrick 1945 *Biochem J (London)* 39 70-72
- 68 Birkinshaw J H A R Chambers and H Raistrick 1942 *Biochem J (London)* 36 242-251

- 69 Birkinshaw J H and P Chaplen 1935 *Biochem J (London)* 60 253-261
- 70 Birkinshaw J H J H V Charles A C Hetherington and H Raistrick 1931 *Phil Trans Roy Soc London Ser B* 220 153-171
- 71 Birkinshaw J H J H V Charles C H Lilly and H Raistrick 1931 *Phil Trans Roy Soc London Ser B* 220 17-138
- 72 Birkinshaw J H J H V Charles and H Raistrick 1931 *Phil Trans Roy Soc London Ser B* 220 333-36
- 73 Birkinshaw J H J H V Charles H Raistrick and J A R Stoyke 1931 *Phil Trans Roy Soc London Ser B* 220 27-54
- 74 Birkinshaw J H and W P K Findlay 1940 *Biochem J (London)* 31 82-88
- 75 Birkinshaw J H W P K Findlay and R A Webb 1940 *Biochem J (London)* 31 906-916
- 76 Birkinshaw J H W P K Findlay and R A Webb 1940 *Biochem J (London)* 36 526-529
- 77 Birkinshaw J H E N Morgan and W P K Findlay 1932 *Biochem J (London)* 50 509-516
- 78 Birkinshaw J H A E Oxford and H Raistrick 1936 *Biochem J (London)* 30 391-411
- 79 Birkinshaw J H and H Raistrick 1931a *Phil Trans Roy Soc London Ser B* 220 215-234
- 80 Birkinshaw J H and H Raistrick 1931b *Phil Trans Roy Soc London Ser B* 220 331-333
- 81 Birkinshaw J H and H Raistrick 1932 *Biochem J (London)* 26 441-453
- 82 Birkinshaw J H and H Raistrick 1936 *Biochem J (London)* 30 2194-2200
- 83 Birkinshaw J H H Raistrick and D J Ross 1932 *Biochem J (London)* 50 630-634
- 84 Birkinshaw J H H Raistrick D J Ross and C E Stickings 1932 *Biochem J (London)* 50 610-628
- 85 Birkinshaw J H H Raistrick and G Smith 1941 *Biochem J (London)* 36 829-833
- 86 Boidin J 1911 *Rev mycol (Paris)* 16 175-197
- 87 Bolcato V and P Tono 1939 *Enzymologia* 7 146-166
- 88 Bomstein R A and M J Johnson 1952 *J Biol Chem* 198 143-153
- 89 Bonner J A Sandoval Y W Tang and L Zechmeister 1946 *Arch Biochem* 10 113-123
- 90 Boothroyd B J A Thorn and R H Haskins 1933 *Can J Biochem Physiol* 33 289-296
- 91 Bowers A T G Halsall E R H Jones and A J Lemin 1933 *J Chem Soc (London)* 1933 248-2560
- 92 Bowers A T G Halsall and G C Sayer 1934 *J Chem Soc (London)* 1934 3040-3084
- 93 Brack A 1947 *Helv Chim Acta* 30 1-8
- 94 Backen A and H Raistrick 1917 *Biochem J (London)* 41 569-574
- 95 Breen J J C Dacre H Raistrick and G Smith 1933 *Biochem J (London)* 60 618-666
- 96 Brian P W P J Curtis and H G Hemming 1949 *Trans Brit Mycol Soc* 32 30-33
- 97 Bu Lock J D E R H Jones and P R Leeming 1933 *J Chem Soc (London)* 1933 4240-476
- 98 Bu Lock J D and E F Leadbeater 1936 *Biochem J (London)* 62 476-480
- 99 Burns J J H B Buich and C G King 1951 *J Biol Chem* 191 501-514
- 100 Burton H S E P Abraham and H M E Caldwell 1936 *Biochem J (London)* 62 171-176

- 101 Butkewitsch W 1921 *Biochem Z* 151 171-190
- 102 Butkewitsch W and M W Fedoroff 1929 *Biochem Z* 207 30-318
- 103 Butkewitsch W (Butkevich V S) and A G Timofeeva 1931 *Mikrobiologiya* 3 574-581
- 104 Cajori I A T T Otrum and M A Hamilton 1951 *J Biol Chem* 208 107-114
- 105 Calam C T A C Oxford and H Raistrick 1939 *Biochem J (London)* 33 1488-1493
- 106 Cantino F C 1913 *Ann J Botany* 36 93-112
- 107 Cantino F C 1951 *Am J Botany* 38 59-583
- 108 Cantino F C and E A Horensstein 1953 *Physiol Plantarum* 8 189-221
- 109 Carson S F J W Foster W E Jefferson F F Phares and D S Anthony 1951 *Arch Biochem Biophys* 33 418-458
- 110 Carson S F F H Mosbach and E F Phares 1951 *J Bacteriol* 62 233-238
- 111 Crude L F Jr and S G Knight 1951 *J Bacteriol* 67 111-111
- 112 Castle H and F Kuhach 1919 *Arch Biochem* 23 158 160
- 113 Celmer W D and I A Solomon 1953 *J Am Chem Soc* 75 13 2 1316
- 114 Charkoff I L and E W Brown Jr 1951 In D M Crenberg (ed) *Chemical Pathways of Metabolism* Vol 1 p 27-317 New York Academic Press
- 115 Chain E 1919 In H W Florey et al (eds) *Antibiotics* Vol 2 p 749-61 London Oxford Univ Press
- 116 Chain E H W Florey N C Heatley and M A Jennings 1943 In H W Florey et al (eds) *Antibiotics* Vol 1 p 1090-1110 London Oxford Univ Press
- 117 Challenger F 1911 *Chem Revs* 36 315-361
- 118 Challenger F 1911 *Advances in Enzymol* 12 479-491
- 119 Challenger F and I T Charlton 1917 *J Chem Soc (London)* 131 474-479
- 120 Challenger F and L Ellis 1933 *J Chem Soc (London)* 1933 336-400
- 121 Challenger F C Higginbottom and L Ellis 1933 *J Chem Soc (London)* 1933 93-101
- 122 Challenger F D B Lisle and I B Dransfeld 1954 *J Chem Soc (London)* 1954 1160-1171
- 123 Challenger F and H E North 1931 *J Chem Soc (London)* 1931 68-71
- 124 Challenger F and A A Rawlings 1937 *J Chem Soc (London)* 1937 868-875
- 125 Challenger F V Subramaniam and T K Walker 1927a *J Chem Soc (London)* 1927 200-208
- 126 Challenger F V Subramaniam and T K Walker 1927b *Nature (London)* 119 674
- 127 Champeau M F and I J Luteraan 1916 *Ann parasitol humaine et comparée* 21 345-353
- 128 Charles J H V H Raistrick R Robinson and A R Todd 1933 *Biochem J (London)* 27 499-511
- 129 Chesters C G C and G N Robinson 1951a *Biol Revs Cambridge Phil Soc* 26 219-252
- 130 Chesters C G C and G W Robinson 1951b *J Gen Microbiol* 1 553-558
- 131 Chichester C O P S Wong and G Mackinney 1954 *Plant Physiol* 29 238-241
- 132 Chrzastecz T 1901 *Centr Bakteriell Parasitenk Abt II* 7 326-338
- 133 Chrzastecz T and R Schillak 1936 *Biochem Z* 288 3 3-368
- 134 Chrzastecz T and D Tiukow 1929 *Biochem Z* 201 106-121
- 135 Chrzastecz T and D Tiukow 1931 *Biochem Z* 240 137-148
- 136 Chrzastecz T, D Tiukow and M Zakomoray 1932 *Biochem Z* 250 24-269

- 137 Chazotte T and M Zakomoray 1933 *Biochem Z* 219 1-10
- 138 Claes H 1954 *Z Naturforsch* 9b 461-469
- 138a. Cleland W W and M J Johnson 1956 *J Biol Chem* 200 593-600
- 139 Clutterbuck P W 1936 *J Soc Chem Ind (London)* 35 557-617
- 140 Clutterbuck P W W N Haworth H Raistrick C Smith and M Stacey 1934 *Biochem J (London)* 28 94-110
- 141 Clutterbuck P W, S L Mukhopadhyay A E Oxford and H Raistrick 1940 *Biochem J (London)* 34 664-677
- 142 Clutterbuck P W A E Oxford H Raistrick and C Smith 1937 *Biochem J (London)* 26 1441-1458
- 143 Clutterbuck P W H Raistrick and F Reuter 1933 *Biochem J (London)* 29 300-311 871-883 1300-1309
- 144 Cochrane V W 1952 *J Bacteriol* 63 429-431
- 145 Cochrane V W and I Dummick 1949 *J Bacteriol* 58 3-30
- 146 Coleman R J M Cefola and F F Nord 1954 *Arch Biochem Biophys* 40 100-110
- 147 Collins F D 1954 *Biol Pers Cambridge Phil Soc* 99 423-431
- 148 Contardi A. and A. Ercoli 1933 *Arch sci Biol (Bologna)* 1 1-14
- 149 Coppock P D V Subramaniam and T H Walker 1958 *J Chem Soc (London)* 1958 1422-1427
- 150 Corbett R. E A W Johnson and A R Todd 1950 *J Chem Soc (London)* 1950 147-149
- 151 Cort L. A et al 1954 *J Chem Soc (London)* 1954 3713-3714
- 151a. Cort W M et al 1956 *Arch Biochem Biophys* 63 41-48
- 152 Grasmann J M 1954 *Am J Botany* 41 302-310
- 153 Crewther W G and F G Lennox 1955 *Australian J Biol Sci* 6 410-419
- 154 Cross L C C. G Elor L M Heilbron and E. R. H Jones 1940 *J Chem Soc (London)* 1940 631-636
- 155 Currie J N 1917 *J Biol Chem* 51 15-31
- 156 Currie J N and C Thom 1915 *J Biol Chem* 28 8-93
- 157 Dakin H D 1906 *J Biol Chem* 1 211-218
- 158 Damm H 1943 *Chemiker Ztg* 61 47-49
- 159 Dammann E O T Rotini and F F Nord 1958 *Biochem Z* 291 183-207
- 160 Davidson R W W A. Campbell and D Blaisdell 1958 *J Agr Research* 57 683-693
- 161 Davies J E F E King and J C Roberts 1955 *J Chem Soc (London)* 1955 2787-2786
- 162 Davis B D 1955 *Advances in Enzymol* 16 46-317
- 163 Dean F M J C Roberts and A Robertson 1954 *J Chem Soc (London)* 1954 1437-1439
- 164 Deuel H J Jr 1951 *The Lipids* Vol 1 New York Interscience Publishers pp 98
- 165 Dewar M J S 1915 *Nature (London)* 150 50-51
- 166 Dimond A E G H Plumb E M Stoddard and J G Horstall 1949 *Conn (New Haven) Agr Exp Sta. Bull* 531 1-69
- 167 Dimond A E and P E Waggoner 1953 *Phytopathology* 43 663-669
- 168 Dingle J and G L Solomons 1952 *J Appl Chem (London)* 2 395-399
- 169 Doelger W P and S C Prescott 1934 *Ind Eng Chem* 6 1149-1149
- 170 Dransfield P B and F Challenger 1951 *J Chem Soc (London)* 1951 1153-1160
- 171 Dulaney E L and D Perlman 1951 *Bull Torrey Bot Club* 4 501-511
- 172 Dulaney E L et al 1955a *Appl Microbiol* 3 336-340
- 173 Dulaney E L et al 1955b *Appl Microbiol* 3 341-344

- 171 Dulaney E L et al 1955 *Mycologia* 47 461-464
- 172 Dyckerhoff H and R Armbruster 1933 *Z physiol Chem Hoppe Seyler's* 219 38-56
- 176 Ehrensward G 1955 *Exp Cell Research Suppl* 3 102-109
- 177 Ehrlich F 1911 *Ber deut chem Ges* 44 3131-3132
- 178 Emerson R and D L Fox 1910 *Proc Roy Soc (London) B* 178 215-223
- 179 Enebo L L G Anderson and H Iundin 1916 *Arch Biochem* 11 383-395
- 180 Engel B G and W Brzeski 1915 *Helv Chim Acta* 30 1472-1478
- 181 Engel B G W Brzeski and I A Hattner 1919 *Helv Chim Acta* 37 1160-1175 1752-1758
- 182 Engel B G J Wursch and M Zimmermann 1953 *Helv Chim Acta* 36 1771-1776
- 183a Eppstein S H I D Meister H C Murray and D H Peterson 1956 *Vitamins and Hormones* 14 359-432
- 183 Erdman H 1918 *Acta Chem Scand* 2 209-219
- 184 Erikson D and J W Porteous 1953 *J Gen Microbiol* 8 464-464
- 185 Erkama J B Hågerstrand and S Junkkunen 1919 *Acta Chem Scand* 3 862-866
- 186 Erkama J J Hienkenen and B Hagerstrand 1919 *Acta Chem Scand* 3 858-861
- 187 Eyre J C 1930 *Ann Appl Biol* 19 351-369
- 188 Iähræus G 1952 *Physiol Plantarum* 5 281-291
- 189 Iähræus G 1954 *Physiol Plantarum* 7 701-712
- 190 Iähræus G and G Lindeberg 1953 *Physiol Plantarum* 6 150-158
- 191 Fairbairn D 1918 *J Biol Chem* 13 105-111
- 192 Galek R and S N Kapur 1921 *Ber deut chem Ges* 77B 920-923
- 193 Junk H G Haenseler and M Schmidt 1937 *Z Spiritusint* 60 71 6-17 81-82 (*Chem Abstr* 31 1936 1937)
- 194 Fiore J V 1918 *Arch Biochem* 1b 161-164
- 195 Fiore J V and F F Nord 1950 *Arch Biochem* 26 382-400
- 196 Fitz A 1873 *Ber deut chem Ges* 6 48-58
- 197 Fleming A 1909 *Brit J Exp Pathol* 10 20-236
- 198 Fodor I J and A Charn 1939 *Lymologia* 13 258-267
- 199 Forsyth F R 1955 *Can J Botany* 33 363-363
- 200 Foster J W 1949 *Chemical Activities of Fungi* New York Academic Press 1 p 618
- 201 Foster J W 1951 *Ann Rev Microbiol* 5 101-120
- 202 Foster J W and S F Carson 1950 *Proc Natl Acad Sci U S* 36 219-223
- 203 Foster J W S F Carson D S Anthony J B Davis W E Jefferson and M A Long 1919 *Proc Natl Acad Sci U S* 3 663-662
- 204 Foster J W S F Carson S Ruben and M D Kamen 1941 *Proc Natl Acad Sci U S* 27 590-596
- 205 Foster J W and J B Davis 1918 *J Bacteriol* 36 329-338
- 206 Foster J W and J B Davis 1919 *Arch Biochem* 11 135-142
- 207 Foster J W and S A Waksman 1939a *J Am Chem Soc* 61 197-197
- 208 Foster J W and S A Waksman 1939b *J Bacteriol* 37 599-617
- 209 Foster J W H B Woodruff and L F McDaniel 1943 *J Bacteriol* 46 421-435
- 210 Fox A S and W D Gray 1950 *Proc Natl Acad Sci U S* 36 538-540
- 211 Franciosi M 1935 *Fermentforschung* 11 433-501
- 212 Franke W and M Dettner 1939 *Ann Chem Liebigs* 511 117-150
- 213 Frey A 1931 *Arch Mikrobiol* 2 212-309
- 214 Friedheim E A H 1933 *Compt rend soc biol* 112 1050-1050
- 215 Friend J and T W Goodwin 1954 *Proc Chem J (London)* 57 431-437

- 216 Fruton J S and S Summons 1953 *General Biochemistry* New York John Wiley and Sons pp 940
- 17 Gailey F B J J Stefaniak B H Olson and M J Johnson 1916 *J Bacteriol* 52 199-140
- 18 Gardner J F L V James and S D Rubbo 1956 *J Gen Microbiol* 14 28-237
- 219 Garton G A T W Goodwin and W Lajnsky 1951 *Biochem J (London)* 48 154-163
- 20 Gascoigne R M J S E Holker B J Ralph and A Robertson 1951 *J Chem Soc (London)* 1951 2346-2359
- 21 Gastrock E A N Porges P A Wells and A J Moyer 1938 *Ind Eng Chem* 30 78-189
- 2 Gayet J 1919 *Bull soc chim biol* 31 19-195
- 3 Geiger Huber M and H Galli 1945 *Helv Chim Acta* 28 248-50
- 4 Gentile A C 1954 *Plant Physiol* 29 257-261
- 22 Gibbs M V W Cochran L M Paegle and H Wolin 1954 *Arch Biochem Biophys* 50 37-247
- 6 Gibbs M and R Gastel 1953 *Arch Biochem Biophys* 43 33-38
- 27 Gill Carey D 1949 *Brit J Exp Pathol* 30 119-123
- 229 God n P 1953a *Biochim et Biophys Acta* 11 114-118
- 9 God n P 1953b *Biochim et Biophys Acta* 11 119-121
- 230 God n P 1955 *Antonie van Leeuwenhoek J Microbiol Serol* 21 215-224
- 31 Golueke C G 1957 *J Bacteriol* 74 337-341
- 3 Goodwin T W 1959a *The Comparative Biochemistry of the Carotenoids* London Chapman and Hall pp 36
- 33 Goodwin T W 1952b *Biochem J (London)* 50 550-558
- 234 Goodwin T W 1953a *Biochem J (London)* 53 538-540
- 35 Goodwin T W 1953b *J Sci Food Agr* 4 209-20
- 236 Goodwin T W 1955 *Ann Rev Biochem* 24 497-520
- 237 Goodwin T W W Lajnsky and J S Willmer 1953 *Biochem J (London)* 53 208-21
- 238 Goodwin T W and J S Willmer 1952 *Biochem J (London)* 51 213-17
- 239 Gordon J J et al 1947 *J Gen Microbiol* 1 187-202
- 240 Gordon M S C Pan A Virgona and P Numerof 1953 *Science* 118 43
- 241 Gordon M A 1950 *Mycologia* 4 167-185
- 242 Gould B S 1938 *Biochem J (London)* 32 797-809
- 243 Gould B S and H Raistrick 1934 *Biochem J (London)* 28 1640-1656
- 244 Grant J K and A C Brownie 1955 *Biochim et Biophys Acta* 18 433-434
- 245 Green D E 1954 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 1 p 276 New York Academic Press
- 246 Gripenberg J 1959 *Acta Chem Scand* 6 580-586
- 247 Grob E C and R Butler 1954 *Helv Chim Acta* 37 1908-1912
- 48 Grob E C and R Butler 1955 *Helv Chim Acta* 38 1313-1316
- 48a Grob E C and R Butler 1956 *Helv Chim Acta* 39 1975-1980
- 249 Grob E C V G undbacher and W H Schopfer 1954 *Experientia* 10 38-39
- 250 Gross S R R D Caffo d and E L Tatum 1956 *J Biol Chem* 219 781-796
- 251 Gross S R and E L Tatum 1955 *Science* 129 1141
- 252 Grosser A H Kundtner Schwarzkopf and K Bernhauer 1950 *Z Naturforsch B* 5 8-30
- 53 Grove J F 1954 *Biochem J (London)* 50 648-666

- 24 Grove J F 1953 *Biochem J (London)* 54 664-673
- 25 Grove J F 1954 *J Chem Soc (London)* 1954 4693-4694
- 26 Grove J F and J C McGowan 1947 *Nature (London)* 160 574
- 27 Cudde J M T G Halsall R Hodges and E R H Jones 1954 *J Chem Soc (London)* 1954 3931-3939
- 28 Cyllenberg H and A Rastio 1950 *Physiol Plantarum* 5 367-371
- 29 Halliwell C 1953 *J Exp Botany* 4 369-376
- 30 Halsall T C and R Hodges 1954 *J Chem Soc (London)* 1954 2382-2391
- 31 Hamada M 1910 *Japan J Botany* 10 387-458
- 32 Harrold C F and M Kling 1952 *J Biol Chem* 194 399-406
- 33 Haskins R H J A Thorn and B Boothroyd 1955 *Can J Microbiol* 1 719-726
- 34 Haxo F 1955 *Fortschr der Chemie org Naturstoffe* 12 169-197
- 35 Hayasida A 1938 *Biochem* 7 193 169-178
- 36 Heim I 1916a *Compt rend (Paris)* 220 1351-1355
- 37 Heim P 1916b *Compt rend (Paris)* 223 1170-1172
- 38 Henderson M F K and V C Farmer 1955 *J Gen Microbiol* 12 37-46
- 39 Herrick H T and O E May 1908 *J Biol Chem* 77 182-192
- 40 Hesse A 1919 *Advances in Enzymol* 9 63-101
- 41 Hesseltine C W and R F Anderson 1957 *Mycologia* 49 449-452
- 42 Heughebaert C and M Welsch 1919 *Bull soc chim biol* 31 282-286
- 43 Hida T 1955 *J Shanghai Sci Inst Ser II* 1 201-214
- 44 Higuchi K F C Jarvis W H Peterson and M J Johnson 1946 *J Am Chem Soc* 68 1669-1670
- 45 Higuchi T and K Kitamura 1953 *J Japan Forestry Soc* 35 30-35
(*Chem Abstr* 48 12875 1954)
- 46 Hirata Y and K Nakanishi 1950 *J Biol Chem* 181 135-143
- 47 Hirsch H M and G I Wallace 1951 *Proc can biol* 10 191-214
- 48 Hiura M and Y Nagata 1954 *Research Bull Fac Agr Cifu Univ (Japan)* No 3 93-99
- 49 Hockenhull D J D M Herbert A D Walker G D Wilkin and F C Winder 1954 *Biochem J (London)* 56 73-82
- 50 Hockenhull D J D A D Walker G D Wilkin and F G Winder 1950 *Biochem J (London)* 50 605-609
- 51 Hoffman Ostenhoff O 1950 *Fortschr der Chemie org Naturstoffe* 6 154-211
- 52 Holker J S E et al 1953 *J Chem Soc (London)* 1953 24 22479
- 53 Hollis J P 1952 *Phytopathology* 42 273-276
- 54 Horecker B L M Gibbs H Klenow and P Z. Smyrniotis 1954 *J Biol Chem* 207 393-403
- 55 Horowitz N H 1946 *J Biol Chem* 160 413-419
- 56 Horowitz N H 1947 *J Biol Chem* 171 255-261
- 57 Horowitz N H and M Fling 1953 *Genetics* 38 360-374
- 58 Horowitz N H and S C Shen 1952 *J Biol Chem* 197 513-520
- 59 Horowitz Wassowa L M and M J Livschitz 1935 *Zentr Bakteriol Parasitenk Abt II* 92 424-435
- 60 Howard B H and H Raistrick 1950 *Biochem J (London)* 46 49-53
- 61 Howard B H and H Raistrick 1955 *Biochem J (London)* 59 475-481
- 62 Howell A Jr and L Pine 1956 *J Bacteriol* 71 47-53
- 63 Ingraham J L and R Emerson 1954 *Am J Botany* 41 146-150
- 64 Irvine G N M Colobchuk and J A Anderson 1954 *Can J Agr Sci* 31 231-239
- 65 Isono M 1953a *J Agr Chem Soc Japan* 27 198-207
- 66 Isono M 1953b *J Agr Chem Soc Japan* 27 25-29

- 996 Isono M 1953c *J Agr Chem Soc Japan* 27 297-301
- 297 Isono M 1954 *J Agr Chem Soc Japan* 28 472-479
- 98 Itto G 1933 *J Agr Chem Soc Japan* 9 52-56. (*Chem Abstr* 27 5141 1933)
- 299 Jacquot R 1933 *Ann fermentations* 4 84-294 316-60
- 300 Jarvis F G and M J Johnson 1947 *J Am Chem Soc* 69 3010-3017
- 301 Jarvis F G and M J Johnson 1950 *J Bacteriol* 59 51-60
- 30 Jefferson W E and J W Foster 1953 *J Bacteriol* 65 587-590
- 303 Jennings M A and T I Williams 1945 *Nature (London)* 155 30.
- 304 Johnston J A D W Racusen and J Bonner 1954 *Proc Natl Acad Sci U S A* 40 1031-1037
- 305 Jones E R H and J D Bu Lock 1953 *J Chem Soc (London)* 1953 3719-3720
- 306 Jones E R H and T G Halsall 1955 *Fortschr der Chemie org Naturstoffe* 12 44-130
- 307 Kardo Syssojewa H 1936 *Zentr Bakteriell Parasitenk Abt II* 93 264-277
- 308 Karow E O and S A Waksman 1947 *Ind Eng Chem* 39 821-8 5
- 309 Karrer P and F Haab 1948 *Helv Chim Acta* 31 795-798
- 310 Karrer P and E Krause Voith 1948 *Helv Chim Acta* 31 802-803
- 311 Katagiri H and K Kitahara 19 9 *Bull Agr Chem Soc Japan* 5 38-47 (*Chem Abstr* 24 3813-3814 1930)
- 31 Kavanagh F A Hervey and W J Robbins 1950 *Proc Natl Acad Sci U S A* 36 10 -106
- 313 Kanten R H 1953 *Biochem J (London)* 55 350-360
- 314 Kinoshita K 1931a *Botan Mag (Tokyo)* 45 45 61
- 315 Kinoshita K 1931b *Acta Phytotom (Japan)* 271 8.
- 316 Kirsh D 1935a *Botan Ca* 97 321-333
- 317 Kirsh D 193 b *J Biol Chem* 108 421-430
- 318 Kleinzeller A 1948 *Advances in Enzymol* 8 39-341
- 319 Kluyver A J and L H C Perquin 1933 *Biochem Z* 266 8 -95
- 30 Kluyver A J and J C M van Zijp 1951 *Antonie an Leeuwenhoek J Microbiol Serol* 17 315-324
- 321 Knobloch H and H Mayer 1941 *Biochem Z* 307 285-292
- 3 Knudson L 1913 *J Biol Chem* 14 159 181 185- 02
- 323 Koffler H R L Emerson D Perlman and R H Burris 194 *J Bacteriol* 50 517-548
- 34 Koffler H S G Knight W C Frazier and R H Burris 1946 *J Bacteriol* 51 385-392
- 325 Kogl F and H Becker 1928 *Ann Chem Liebigs* 465 211-242
- 326 Kogl F H Erxleben and L Jüncke 1930 *Ann Chem Liebigs* 48 105-119
- 37 Kogl F and J J Postowsky 1925 *Ann Chem Liebigs* 444 1-7
- 28 Kogl F and G C van Wessem 1944 *Rec t av chim* 63 5-12
- 329 Kráml A and J Hor ath 1948 *Nature (London)* 16 619
- 330 Kráml A and J Horvath 1949 *Natu e (London)* 163 219
- 331 Kresling E and E Stern 1936 *Zentr Bakteriell Parasitenk Abt II* 95 37-310
- 332 Kusnetsov S J 1931 *Zent Fakteriell Parasitenk Abt II* 83 37-52
- 333 Landa S V Solin and J Palatý 1953 *Chem listy* 47 1066-1070 (*Chem Abstr* 48 4043 1954)
- 334 Larsen H and K E Esmhjellen 19 5 *Biochem J (London)* 60 135 139
- 335 Law R 19 0 *Ann Botany* 14 69-78
- 336 Lein J T A Puglisi and P S Lein 1953 *Arch Biochem Biophys* 45 434-442
- 337 Lewis K F and S Wenhause 1951a *J Am Chem Soc* 73 2500- 503

- 338 Lewis K. F. and S. Weimhouse 1931b *J Am Chem Soc* 73 2906-2909
- 339 Liggett R. W. and H. Koffler 1918 *Bacteriol Revs* 12 297-311
- 340 Lindeberg G. 1918 *Physiol Plantarum* 1 196-203, 401-409
- 341 Lindeberg G. 1919 *Siensk Botan Tidskr* 45 438-447
- 342 Lindeberg G. 1920 *Nature (London)* 166 739
- 343 Lindeberg G. and G. Fähræus 1920 *Physiol Plantarum* 5 277-283
- 344 Lindeberg G. and G. Holm 1922 *Physiol Plantarum* 7 100-114
- 345 Lindner P. 1922 *Z angew Chem* 35 110-114
- 346 Lipmann F. 1934 *Science* 190 855-863
- 347 Lloyd G. A. Robertson G. B. Sankey and W. B. Whalley 1935 *J Chem Soc (London)* 1935 2103-216
- 348 Lockwood L. B. and G. E. N. Nelson 1916 *Arch Biochem* 10 363-374
- 349 Lockwood L. B. K. B. Raper A. J. Moyer and R. D. Coghill 1915 *Am J Botany* 32 214-217
- 350 Lockwood L. B. and M. D. Reeves 1915 *Arch Biochem* 6 455-469
- 351 Lockwood L. B. J. J. Stubbs and C. F. Senseman 1938 *Zentr Bakteriell Parasitenk Abt II* 98 167-171
- 352 Lockwood L. B. and G. E. Ward 1915 *Ind Eng Chem* 37 403-406
- 353 Lockwood L. B. C. F. Ward and O. F. May 1936 *J Agr Research* 53 819-837
- 354 Loesche H. W. von 1915 *Chem Eng News* 23 1930-1939
- 355 Lund N. A. A. Robertson and W. B. Whalley 1933 *J Chem Soc (London)* 1933 2454-2459
- 356 Lutz L. 1931 *Bull soc chim biol* 13 436-457
- 357 Lutz G. 1934 *Phytopathol Z* 7 583-638
- 358 Maas Forster M. 1935 *Arch Mikrobiol* 22 115-144
- 359 McCallan S. E. A. and F. Wilcoxon 1936 *Contribs Boyce Thompson Inst* 8 151-165
- 360 McCown J. M. 1930 *Thesis Univ of California* pp 116
- 361 Mackinnon C. C. O. Chichester and P. S. Wong 1933 *Arch Biochem Biophys* 53 419-483
- 362 MacMillan J. 1933 *J Chem Soc (London)* 1933 1697-1702
- 363 MacMillan J. 1934 *J Chem Soc (London)* 1934 2585-2587
- 364 Mamoli L. and A. Vercellone 1937 *Z physiol Chem Hoppe Seyler's* 245 93-95
- 365 Mandels G. R. 1935a *Arch Biochem Biophys* 42 164-173
- 366 Mandels G. R. 1935b *Arch Biochem Biophys* 41 362-377
- 367 Mapson L. W. 1935 *Vitamins and Hormones* 13 71-100
- 368 Martin E. et al 1953 *J Biol Chem* 203 239-250
- 369 Martin S. M. P. W. Wilson and R. H. Burris 1950 *Arch Biochem* 26 103-111
- 370 Martin W. R. and J. W. Foster 1935 *J Bacteriol* 70 403-414
- 371 Mason H. S. 1935 *Advances in Enzymol* 16 103-184
- 372 May O. E. H. T. Herrick C. Thom and M. B. Church 1927 *J Biol Chem* 10 417-422
- 373 May O. E. A. J. Moyer I. A. Wells and H. T. Herrick 1931 *J Am Chem Soc* 53 774-782
- 374 May O. E. G. E. Ward and H. T. Herrick 1932 *Zentr Bakteriell Parasitenk Abt II* 86 179-181
- 375 Mead T. H. and M. V. Stack 1948 *Biochem J (London)* 42 xviii
- 376 Mininger R. I. M. E. Wall R. C. Dworschack and R. W. Jackson 1936 *Arch Biochem Biophys* 60 427-439
- 377 Miyaji K. 1930 *Research Bull Gifu Imp Coll Agr (Japan)* No 10 1-269 277
- 378 Miller E. V. J. R. Winston and D. F. Fisher 1940 *J Agr Research* 60 269 277
- 379 Mollard M. 1922 *Compt rend (Paris)* 174 881-883

- 380 Mortimer D C and M J Johnson 1952 *J Am Chem Soc* 74 4098-4102
- 381 Morton H E W Kocholaty R Junowicz Kocholaty and A Keiner 1915
J Bacteriol 50 519-584
- 382 Moses V 1955 *J Gen Microbiol* 13 235-251
- 383 Moyer A J 1955 *Appl Microbiol* 1 1-7
- 384 Moyer A J and R D Coghill 1945 *Arch Biochem* 7 167-183
- 385 Moyer A J and R D Coghill 1946 *J Bacteriol* 51 57-78
- 386 Moyer A J E J Umberger and J J Stubbs 1940 *Ind Eng Chem* 32
1319-1383
- 387 Mukherjee S 1951 *Arch Biochem Biophys* 33 364-366
- 388 Mukherjee S 1952 *Arch Biochem Biophys* 35 23-33 34-59
- 389 Muil R P and F F Nord 1944 *Arch Biochem* 4 419-433
- 390 Munekata H 1953 *J Biochem (Tokyo)* 40 451-460
- 391 Nagata Y A Masuda and M Hiura 1954 *Research Bull Fac Agr Gifu
Univ (Japan)* No 3 100-101
- 392 Nakayama T G Mackinnon and H J Phaff 1954 *Antonie van Leeuwen
hoek J Microbiol Serol* 20 217-228
- 393 Newton G G F E P Abraham and C W Hale 1954 *Biochem J (Lon
don)* 58 103-111
- 394 Nickerson W J 1948 *Arch Biochem* 17 225-233
- 395 Nishikawa H 1939 *Acta Phytochim (Japan)* 11 167-185
- 396 Nishikawa H 1940 *Bull Agr Chem Soc Japan* 16 97-98
- 397 Nobles M K 1948 *Can J Research C* 26 281-431
- 398 Nord F F 1940 *Chem Revs* 26 422-442
- 399 Nord F F and W Engel 1938 *Biochem Z* 296 153-170
- 400 Nord F F and L J Sciarini 1946 *Arch Biochem* 9 419-437
- 401 Nord F F and J C Vitucci 1947 *Arch Biochem* 14 29-241
- 402 Nord F F and J C Vitucci 1948 *Advances in Enzymol* 8 253-298
- 403 Nord F F and S Weiss 1951 In J B Sumner and K. Myrback (eds) *The
Enzymes* Vol 2 (1) p 681-790
- 404 Ohara Y 1952 *J Agr Chem Soc Japan* 26 547-551
- 405 Olson J A 1954 *Nature (London)* 174 692-696
- 406 Oroshnik W L C Vining A D Mebane and W A Taber 1955 *Science*
121 147-149
- 407 Otike R C E L Tatum I Zabun and K Bloch 1951 *J Biol Chem* 189
479-413
- 408 Oxford A E and H Raistrick 1932 *Biochem J (London)* 26 1902-1906
- 409 Oxford A E and H Raistrick 1933 *Biochem J (London)* 27 1473-1478
- 410 Oxford A E and H Raistrick 1934 *Biochem J (London)* 28 1321-1374
- 411 Oxford A E H Raistrick and F Simonart 1939 *Biochem J (London)*
33 240-248
- 412 Pauson P L 1955 *Chem Revs* 55 9-136
- 413 Pearson L K and H S Raper 1927 *Biochem J (London)* 21 875-879
- 414 Perlman D 1948 *Am J Botany* 35 350-363
- 415 Perlman D 1949 *Am J Botany* 36 180-184
- 416 Perlman D 1950a *Am J Botany* 37 237-241
- 417 Perlman D 1950b *Botan Rev* 16 449-503
- 418 Perlman D 1950c *Bull Torrey Botanical Club* 77 103-109
- 419 Perlman D 1951 *Am J Botany* 38 650-658
- 420 Perlman D D A Kita and W H Peterson 1948 *Arch Biochem* 11 123-
129
- 421 Perlman D and E O'Brien 1953 *Bacterial Proc (Soc Am Bacteriologists)*
1953 20
- 422 Perworszansky W W 1930 *Zentr Bakteriell Pathogen Abt II* 81 372-
390
- 423 Peterson D H and H C Murray 1952 *J Am Chem Soc* 74 1871-1872

- 424 Plotho O von 1910 *Arch Mikrobiol* 11 33-42
- 425 Pontillon C 1930 *Compt rend (Paris)* 191 1148-1151
- 426 Porges N 1932 *Botan Gaz* 91 197-205
- 427 Porter J W and R E Lincoln 1950 *Arch Biochem* 27 390-403
- 427a Posternak T 1910 *Helv Chim Acta* 23 1046-1053
- 428 Prill E A P R Wenck and W H Peterson 1935 *Biochem J (London)* 29 21-33
- 429 Raaf H 1941 *Arch Mikrobiol* 12 131-162
- 430 Raistrick H 1950 *Proc Roy Soc (London) B* 136 481-508
- 431 Raistrick H R Robinson and D E White 1936 *Biochem J (London)* 30 1303-1314
- 432 Raistrick H and P Simonart 1933 *Biochem J (London)* 27 698-633
- 433 Raistrick H and G Smith 1935 *Biochem J (London)* 29 606-611
- 434 Raistrick H and G Smith 1936 *Biochem J (London)* 30 1315-1322
- 435 Raistrick H C E Stickings and R Thomas 1953 *Biochem J (London)* 55 421-433
- 436 Raistrick H and J Ziffer 1951 *Biochem J (London)* 49 563-574
- 437 Ralph B J and A Robertson 1950 *J Chem Soc (London)* 1950 3380-3383
- 438 Ramachandran K and T K Walker 1951 *Arch Biochem Biophys* 31 224-233
- 439 Ramachandran K and T K Walker 1952 *Arch Biochem Biophys* 35 195-203
- 440 Ramakrishnan C V and B N Banerjee 1952 *Arch Biochem Biophys* 37 131-135
- 441 Raper K B and C. Thom 1949 *A Manual of the Penicillia* Baltimore The Williams and Wilkins Co pp 87
- 442 Raveux R 1948 *Bull soc chim biol* 30 346-357 357-366
- 443 Raveux R 1954 *Compt rend (Paris)* 238 1150-1152
- 444 Rebstock M C H M Crooks Jr J Controulis and Q R Bartz 1949 *J Am Chem Soc* 71 2458-2462
- 444a Ribéreaux Gayon J et al 1955 *Bull soc chim biol* 37 105-106
- 445 Rich S and J G Horsfall 1954 *Proc Natl Acad Sci U S A* 40 139-145
- 446 Rippel A 1940 *Arch Mikrobiol* 11 271-284
- 447 Rippel A and G Behr 1934 *Arch Mikrobiol* 5 561-577
- 448 Rippel A and J Keseling 1930 *Arch Mikrobiol* 1 60-77
- 449 Robinson G N 1954 *J Gen Microbiol* 11 412-419
- 450 Rotini O T E Dammann and F F Nord 1936 *Biochem Z* 288 414-420
- 451 Ruelius H W and A Gauhe 1950 *Ann Chemie Liebigs* 569 38-59
- 452 Saito K 1907 *Botan Mag (Tokyo)* 21 7-11
- 453 Saito K 1911 *Centr Bakteriell Parasitenk Abt II* 29 289-290
- 454 Sakaguchi K T Asai and H Munekata 1942 *Bull Agr Chem Soc Japan* 18 60
- 455 Sakaguchi K and S Baba 1942a *Bull Agr Chem Soc Japan* 18 82-86
- 456 Sakaguchi K and S Baba 1942b *Bull Agr Chem Soc Japan* 18 95
- 457 Sakaguchi K and T Inoue 1940 *Bull Agr Chem Soc Japan* 16 158
- 458 Sakaguchi K T Inoue and S Tada 1939 *Zentr Bakteriell Parasitenk Abt II* 100 302-307
- 459 Sakaguchi K and S Murao 1950 *J Agr Chem Soc Japan* 23 411
- 460 Sakaguchi K H Takahashi and J Kataoka 1951 *J Agr Chem Soc Japan* 25 180-185
- 461 Satina S and A F Blakeslee 1928 *Proc Natl Acad Sci U S A* 14 308-316
- 462 Schatz A K Savard and I J Pintner 1949 *J Bacteriol* 58 117-125
- 463 Schopfer W H and F C. Grob 1952 *Experientia* 8 140
- 464 Schreyer R 1928 *Biochem Z* 209 131-138
- 465 Schreyer R 1931 *Biochem Z* 240 295-305

- 466 Schulz G 1937 *Planta* 27 196-218
- 467 Sebek O K 1953 *Proc Soc Exp Biol Med* 84 110-112
- 468 Sekizawa Y 1955 *J Biochem (Tokyo)* 42 217-219
- 469 Sheng T C and C Sheng 1952 *Genetics* 37 264-69
- 470 Shimazono H 1955 *J Biochem (Tokyo)* 42 321-340
- 471 Shipe W F Jr 1951 *Arch Biochem* 30 162-169
- 472 Shu P and M J Johnson 1948a *Ind Eng Chem* 40 1702-1703
- 473 Shu P and M J Johnson 1948b *J Bacteriol* 55 577-583
- 474 Simonart P and P Godin 1951 *Bull soc chim Belges* 60 446-449
- 475 Singh J and T K Walker 1956 *Biochem J (London)* 62 986-989
- 476 Sisler H D and C E Cox 1954 *Am J Botany* 41 338-345
- 477 Smith R A and I C Gunsalus 1953 *Nature (London)* 173 774-775
- 478 Smith V M 1949 *Arch Biochem* 22 975-987
- 479 Stanier R Y 1948 *J Bacteriol* 55 447-493
- 480 Stanier R Y and O Hayaishi 1951 *Science* 114 396-399
- 481 Stanier R Y and J L Ingraham 1951 *J Biol Chem* 210 890-898
- 482 Starkle M 1951 *Biochem Z* 151 371-415
- 483 Stefaniak J J F B Garley C S Brown and M J Johnson 1946 *Ind Eng Chem* 38 666-671
- 484 Stefaniak J J F B Garley F G Jarvis and M J Johnson 1946 *J Bacteriol* 52 119-128
- 485 Steinberg M P and Z J Ordal 1954 *J Agr Food Chem* 2 873-877
- 486 Steiner M and H Hememann 1954 *Naturwiss* 41 40-41
- 487 Stephens C R et al 1954 *J Am Chem Soc* 76 5 68-5375
- 488 Stern A M Z J Ordal and H O Halvorson 1954 *J Bacteriol* 68 24-27
- 489 Stevens C M P Vohra and C W DeLong 1954 *J Biol Chem* 211 297-300
- 490 Stevens C M P Vohra E Inamine and O A Roholt Jr 1953 *J Biol Chem* 201 1001-1006
- 491 Stewart W D W L Wachtel J J Shipman and J A Yanko 1954 *Science* 127 1271-1272
- 492 Stodola F H M Friedkin A J Moyer and R D Coghill 1954 *J Biol Chem* 161 139-149
- 493 Stodola F H O L Shotwell and L B Lockwood 1955 *J Am Chem Soc* 77 5415-5418
- 494 Stokoe W N 1958 *Biochem J (London)* 27 80-93
- 495 Subramanian V H B Stent and T K Walker 1959 *J Clin Soc (London)* 1959 2482-2497
- 496 Sumiki Y 1959 *Bull Agr Chem Soc Japan* 5 10-13
- 497 Sumiki Y 1951 *Bull Agr Chem Soc Japan* 7 62-63
- 498 Sumner J B and G F Somers 1953 *Enzymes* 3d ed New York Academic Press pp 462
- 499 Sussman A S P Coughley and J C Strain 1955 *Am J Botany* 42 810-813
- 500 Sussman A S and C L Markert 1953 *Arch Biochem Biophys* 45 31-40
- 501 Takahashi T and T Asai 1953 *J Ag Chem Soc Japan* 8 6 638 (Chem Abstr 26 5602 1953)
- 502 Takahashi T and T Asai 1953 *Zentr Baktériol Parasitenk Abt II* 80 81-81
- 503 Takahashi T and K Sakaguchi 1957 *Bull Agr Chem Soc Japan* 3 59-6
- 504 Tamura H 1957 *Acta Phytochem (Japan)* 6 1-19
- 505 Tatum E L 1944 *Ann Rev Biochem* 13 567-604
- 506 Tatum E L and S R Gross 1956 *J Biol Chem* 219 19-80
- 507 Tatum E L S R Gross G Ehrensvard and L Garajda 1951 *Proc Natl Acad Sci U S A* 40 971-976
- 508 Terroine E F R Bonnet G Kopp and J Véchor 1957 *Bull Soc et m biol* 9 605-610

- 509 Thaler H and G Geist 1939 *Biochem Z* 309 191-196
- 510 Thaler H A Schoitmayer I Stählin and H Beck 1949 *Biochem Z* 308 79-98
- 511 Thaler H and I Stählin 1949 *Biochem Z* 309 81-86
- 512 Thibodeau R and H Macy 1942 *Missouri Agr Exp Sta Tech Bull* 152 1-36
- 513 Thies W 1930 *Zentr Bakteriell Parasitenk Abt II* 82 321-347
- 514 Thies W 1931 *Ber deut chem Ges* 61B 214-218
- 515 Thom C and J N Currie 1916 *J Agr Research* 7 1-15
- 516 Thorn J A and M J Johnson 1900 *J Am Chem Soc* 72 2032-2038
- 517 Tome J H D Zook R B Wagner and R W Stone 1953 *J Biol Chem* 203 251-255
- 518 Tomlinson N J J R Campbell and P C Trussell 1951 *J Bacteriol* 61 17-25
- 519 Tóth C and G Bárány 1943 *Enzymologia* 11 19-23
- 520 Treccani V N Walker and C H Wiltshire 1954 *J Gen Microbiol* 11 311-318
- 521 Turfitt G F 1913 *Biochem J (London)* 3 115-117
- 522 Turfitt G E 1914a *Biochem J (London)* 38 492-496
- 523 Turfitt G F 1914b *J Bacteriol* 47 487-493
- 524 Turfitt C E 1916 *Biochem J (London)* 40 79-81
- 525 Turfitt G E 1917 *J Bacteriol* 54 557-566
- 526 Turfitt C F 1918 *Biochem J (London)* 42 376-383
- 527 Turian C 1952 *Experientia* 8 309
- 528 Turian C and F T Haxo 1952 *J Bacteriol* 63 690-691
- 529 Turian C and F T Haxo 1954 *Botan Gaz* 115 254-260
- 530 Tytell A A et al 1955 *Antibiotics Annual 1954/55* 716-718
- 531 Van Lanen J M and F W Tanner Jr 1948 *Vitamins and Hormones* 6 163-274
- 532 Vercellone A and L Mamoli 1937 *Z physiol Chem Hoppe Seyler's* 218 27-29
- 533 Verujki D 1884 *Ann Inst Pasteur* 1 369-391
- 534 Vining L C W A Taber and F J Gregory 1955 *Antibiotics Annual 1954/55* 980-987
- 535 Wain R I and F H Wilkinson 1946 *Ann Appl Biol* 33 401-405
- 536 Waksman S A and I J Hutchings 1951 *J Am Chem Soc* 73 545-547
- 537 Wald G 1911 *Science* 113 287-291
- 538 Walker T K 1919 *Advances in Enzymol* 9 531-584
- 539 Walker T K V Subramaniam and F Challenger 1927 *J Chem Soc (London)* 1927 3044-3054
- 540 Ward G E L B Lockwood O E May and H T Herrick 1935 *Ind Eng Chem* 27 318-329
- 541 Ward G E L B Lockwood O E May and H T Herrick 1936 *J Am Chem Soc* 58 1286-1288
- 542 Ward G E L B Lockwood B Tabenkin and P A Wells 1938 *Ind Eng Chem* 30 1233-1235
- 543 Webley D M and P C DeLoock 1952 *Biochem J (London)* 51 371-375
- 544 Webley D M R B Duff and V C Farmer 1955 *J Gen Microbiol* 13 361-369
- 545 Wehmer C 1891 *Botan Ztg* 49 233-246 249-257 271-280 289-298 30-313 321-332 337-346 353-363 369-374 385-396 401-407 417-428 433-439 449-455 465-478 511-518
- 546 Wehmer C 1893a *Bull soc chim biol* 3 Sér 9 728-730
- 547 Wehmer C 1893b *Compt rend (Paris)* 117 332-333
- 548 Wehmer C 1897 *Chem Ztg* 21 1022-1023

- 549 Wehmer C 1918 *Ber deut chem Ges* 51B 1663-1668
- 550 Wehmer C 1918 *Biochem Z* 197 418-432
- 551 Weiss B 1933 *J Biol Chem* 101 31-43
- 552 Weiss S J V Fiore and F F Nord 1947 *Arch Biochem* 15 3 6-398
- 553 Wells P A A J Moyer and O E May 1936 *J Am Chem Soc* 58 555-558
- 554 Wells P A et al 1937 *Ind Eng Chem* 29 653-656
- 555 Welsch M and C Heugshem 1948 *Compt rend soc biol* 140 10 4-1076
- 556 Wenck P R W H Peterson and H C Greene 193 *Zentr Bakteriol Parasitenk Abt II* 9 324-330
- 557 Wettstein A 1935 *Experientia* 11 465-479
- 558 Whitfield G B et al 1955 *J Am Chem Soc* 77 4190-4801
- 559 Williamson C E 1950 *Phytopathology* 40 205-208
- 560 Witter R F and E Stott 1946 *Arch Biochem* 9 531-539
- 561 Woodbaine M M E Gregory and T A Walker 1951 *J Exp Botany* 2 201-211
- 562 Woodruff H B and J W Foster 1943 *Arch Biochem* 9 301-313
- 563 Woodward R B and G Singh 1949 *J Am Chem Soc* 71 708 709
- 564 Wosilait W D and A Nason 1951 *J Biol Chem* 200 2 5-9 0
- 565 Wosilait W D A Nason and A J Terrell 1954 *J Biol Chem* 206 271-280
- 566 Wright B E 1951 *Arch Biochem Biophys* 31 332-333
- 567 Wunschendorf H and C Killian 1928 *Compt rend (Paris)* 187 572-514
- 568 Yabuta T 1954 *J Chem Soc (London)* 125 575-587
- 569 Yamamoto G 1950 *Symposium Enzyme Chemistry (Japan)* 5 74-81 (*Chem Abstr* 45 8079 1951)
- 570 Yasuda S H Hori and A Habiya 1951 *J Agr Chem Soc Japan* 25 357-361
- 571 Young R E H A Pratt and J B Dale 1951 *Plant Physiol* 6 304-310
- 572 Zalokar M 1954 *Arch Biochem Biophys* 50 71-80
- 573 Zalokar M 1955 *Arch Biochem Biophys* 56 318-325
- 574 Zeller S M 1916 *Ann Missouri Botan Garden* 3 439-509
- 575 Zellner J 1907 *Chemie der höheren Pilze* Leipzig W Engelmann pp 2 7
- 576 Zellner J and E Zikmunds 1930 *Monatsh Chem* 56 200-203
- 577 Zofell C E 1950 *Advances in Enzymol* 10 443-486

7. Respiration

Respiration in the most general sense includes all cellular oxidations which yield energy or could yield energy to the cell. In this chapter after touching on a few general problems we shall consider primarily those reactions which are or may be part of organized respiratory systems. Many individual reactions described in the preceding chapter undoubtedly yield energy but their relation to the major pathways of respiration is not yet known.


Respiration may be aerobic or anaerobic. Aerobic respiration is defined by the fact that the ultimate hydrogen acceptor is molecular oxygen. In anaerobic respiration or fermentation hydrogen from substrate is transferred to compounds other than oxygen (107). Use of the term fermentation to describe an aerobic process is misleading.

The maximum energy available from a reaction or sequence of reactions is of course the free energy change. Thus complete aerobic oxidation of glucose to carbon dioxide and water involves a change in free energy of approximately $-700,000$ calories per mole under standard conditions. Some fraction of this energy is trapped by the cell and used for energy requiring reactions. So far as we now know the size of the fraction so trapped depends on the ability of the organism to couple the oxidation to phosphorylation of adenosine mono-phosphate or adenosine diphosphate to yield adenosinetriphosphate (ATP). In turn the hydrolysis of ATP is coupled to energy requiring reactions. The common practice of describing ATP and similar com-

pounds as having high energy phosphate bonds is ambiguous (103)

In general most of the oxidative phosphorylations associated with glucose or other substrate oxidations are carried out at the cofactor level electrons from substrate are passed along a chain of respiratory carriers to oxygen and in the process in ways still unknown in detail the energy released in electron transfer is used to form ATP Substrate level phosphorylations occur in the anaerobic conversion of glucose to ethanol and carbon dioxide—the Embden Meyerhof path way—and possibly in the oxidative decarboxylation of α ketoglutaric acid

As described briefly in Chapter 9 fungi during growth transform inorganic into organic phosphate Organic phosphorus compounds which have been identified in fungi and actinomycetes include sugar phosphates (38 91 134) adenosine phosphates (9 162 219) and various of the other known nucleotides (9) A mycelial extract of *Rhizopus nigricans* has myokinase activity (228) catalyzing the reversible transphosphorylation



The utilization of ATP to phosphorylate organic molecules is implicit in the two major respiratory pathways for glucose both of which start with glucose 6 phosphate There is very suggestive evidence that ATP generated by oxidative processes is involved in the streaming of the myxomycete *Physarum polycephalum* as it is in muscular contraction (3 177 303 301 305)

1 PROBLEMS OF METHOD

The respiration of whole cells and of extracts is usually measurable in terms of gas exchange—oxygen uptake or carbon dioxide formation The basic instrument in most laboratories is the Warburg Barcroft constant volume manometer described in the manuals of Dixon (77) and Umbreit et al (308) Macrorespirometers with larger capacity have been designed (170 188 282 328) and a few studies on fungi have been performed with more sensitive manometric apparatus (78 214 324) An amperometric procedure for measuring oxygen utilization has been applied successfully in studies on *Penicillium chrysogenum* (52)

Dye reduction methods have a more limited application but are useful in some problems The best known is that of Thunberg (298)

which depends on the reduction of methylene blue coupled to oxidation of substrate

Substrate + oxidized dye \rightarrow oxidized substrate + reduced dye (2)

The Thunberg method has been used to demonstrate the presence of respiratory enzymes in cell free extracts from fungi (182-226). Methylene blue is often inhibitory (271) and other dyes e.g. 2,6-dichlorophenolindophenol are in general to be preferred (30-207, 216-225).

Spectrophotometric methods are both more powerful and more quantitative. The most generally useful of these relies upon the fact that the reduced forms of the coenzymes diphosphopyridine nucleotide (DPN) and triphosphopyridine nucleotide (TPN) absorb strongly at $340\text{ m}\mu$; the change in density can be followed easily and rapidly after addition of substrate to a suitable system (312). Examples of the use of this method in studies of fungi and actinomycetes include work on the enzymes of *Neurospora crassa* (81-301), *Aspergillus niger* (148), *Fusarium lini* (62) and *Streptomyces coelicolor* (66).

Spectrophotometric measurements can of course be used whenever a component of the system changes its light absorption as a consequence of an enzymatic reaction. Oxidation and reduction of cytochrome c can be followed at $550\text{ m}\mu$ (46-47, 71-280) or reducible dyes may be used (50).

Like all other sensitive measurements the spectrophotometric requires a certain amount of judgment. Too often a crude enzyme preparation contains unsuspected activities which complicate a given determination. For example an extract may reduce both DPN and TPN when fructose 1,6-diphosphate is added. The DPN reduction is straightforward: aldolase forms triose phosphate which is then oxidized with concurrent reduction of the coenzyme. The reduction of TPN is not however as might first be thought the same process: instead fructose 1,6-diphosphate is first converted to glucose 6-phosphate which is oxidized by glucose-6-phosphate dehydrogenase and TPN to 6-phosphogluconate. The crude experiment falsely suggests that triose phosphate dehydrogenase can react with either DPN or TPN.

These manometric and spectrophotometric methods are general in their application and therefore stand in contrast to the problem of preparing cells and extracts of fungi for respirometric study. In a certain sense each organism is a problem in itself but it may be useful here to mention some of the methods which have been used in particular instances. Discs of growing mycelium from a surface

culture may be transferred to a respirometer flask (36 78 214 313) but most workers have preferred to use when possible the more homogeneous mycelium obtained from shaken or aerated cultures (62 70 102 163). The mycelium from such cultures is often in the form of small pellets; if these are too large (above 0.1 mm. for *Mycrothecium verrucaria* (70)) the rate of diffusion of oxygen into the interior of the pellet may be limiting. Either surface or shake culture mycelium may be homogenized briefly but there is a considerable risk that mechanical injury may reduce respiratory capacity (70 80).

The preparation of active cell free extracts from fungi is an even more vexing problem. We may list here in outline some of the methods which have been successful.

- 1 Hand grinding with abrasives (7 44 50 62 71 140 182 301)
- 2 Acetone drying (14 129 212 288)
- 3 Physical drying with or without a preliminary freezing (89 155 201 218 331)
- 4 Mechanical disruption in a ball mill (240) or in a motor-driven homogenizer (45 274)
- 5 Freezing followed by grinding with abrasives or by homogenization (190 201 208 274)
- 6 Sonic oscillation (210 261)
- 7 Disruption in the Hughes press (142) or in a small-orifice press (52 200 228)
- 8 Treatment of frozen mycelium with dilute ammonium hydroxide followed by mild pressure (148)

Of these the most generally useful appear to be hand grinding with sand or powdered glass, sonic oscillation and treatment in the Hughes or similar presses. Hugo (143) reviews the methods available for the preparation of extracts from bacteria.

Suspending media for extraction may be water, phosphate buffer or salt mixtures. Protective materials are often essential—both cysteine (62 209) and ethylenediaminetetraacetate (110) protect against inactivation of enzymes by heavy metals liberated during the extraction process.

During the preparation of cell free extracts the terminal respiratory system is usually disrupted and must be replaced by an added hydrogen carrier or hydrogen acceptor. Methylene blue, 2,6-dichlorophenol, indophenol and other dyes have often been used; phenazine derivatives (76), ferricyanide (134 235) and manganese dioxide (133) are of value for the same purpose.

2 FACTORS IN RESPIRATION

In general it is probable that the most important factor in respiration is the species. Few comparative data are available but we may assume that fungi which can grow on a given substrate can also oxidize it (291) so that the omnivorous saprophytes e.g. *Aspergillus* spp. may be expected to oxidize a greater variety of compounds than do nutritionally more specialized forms. Within a given species age and developmental status are the primary non-environmental factors the permeability of the cell is also presumed to be important by the usual indirect evidence. The principal external factors are acidity temperature oxygen pressure carbon dioxide and accessory nutrients.


Age and Developmental Status As mentioned in Chapter 1 the highest respiratory activity is usually found during the period of most rapid growth. Subsequent to this period respiration especially endogenous respiration falls off with increasing age. This may of course reflect only the fact that more and more dead cells accumulate in the older mycelium quite possibly the respiration of the growing tips of the thallus is independent of the age of the culture so long as nutrients are available.

The data of Figure 1 are typical and many others of the same type have been reported in studies of fungi and aerobic actinomycetes (52 70 95 96 115 134 163 249). *Ustilago zeae* appears unusual in that once the maximum respiratory rate is reached it is maintained (260).

Spore respiration is considered briefly in Chapter 12 there is as yet no reason to think that it differs fundamentally from that of mycelium although it would not be surprising to find generally higher activity in germinating spores than in older cultures. The yeast phase of *Blastomyces brasiliensis* exhibits an endogenous oxygen uptake greater than that of the mycelial form (215) and the same relation prevails between cell types of *Mucor guilliermondii* (179). In *Blastocladiella emersonii* the two different types of mature thallus differ in their content of important respiratory enzymes and presumably in their respiratory pathways (44). Oxygen consumption of *Dictyostelium discoideum* is in part a function of developmental stage (116).

Cell Permeability It is difficult to distinguish between intrinsic inability of a cell to respire an exogenous substrate and failure of the compound to penetrate to the interior of the cell. As in the utilization of organic acids for growth (Chapter 3) it is often found that a low pH is required for or accelerates the oxidation of organic acids

Figure 1 The effect of culture age on the endogenous respiration of *Fusarium solani* mycelium. Curve 1 dry weight of culture; curve 2 respiratory rate in μl oxygen per milligram dry cells per hour.



(93) or that provision of an ester of the acid increases its biological effect (13). Or intact cells of a fungus may fail to oxidize a substrate while cell free extracts do so rapidly (13, 65, 89, 258). All of these types of evidence suggest that the intact cell is impermeable to the substrate. In general ionizable substances e.g. phosphorylated compounds and organic acids are the most likely to be excluded by a permeability barrier.

However failure of exogenous substrate to be utilized cannot always be ascribed to membrane impermeability. Thus in yeast frozen cells are permeable to citrate but metabolize it only after further treatment with chloroform (90). It is possible for enzymes in the cell to be inaccessible by virtue of subcellular organization; in most cases we cannot distinguish between this situation and true membrane effects.

Induction of Respiratory Enzymes The enzymes of whatever fundamental respiratory pathways an organism has are probably always present whatever the substrate. But it is often observed that some compounds are oxidized only after a long lag or only if the cells have previously been grown in contact with them. This suggests that certain enzymes, probably those concerned in the preliminary steps of respiratory breakdown, are not constitutive but are induced by substrate. The oxidation of phenylacetic acid is an example (Chapter 6) and attack on other aromatic compounds by *Hormodendrum* sp.

appears also to require the formation of inducible enzymes (127). In *Streptomyces coelicolor* gluconic acid and pentoses are oxidized only by specifically adapted cells (66). In none of these cases just cited however has the specific enzyme been isolated and studied as such.

Acidity and Respiration In most experiments reported the pH of the suspending medium has little effect on endogenous metabolism over the range pH 5 to 8 (3, 70, 187) and the data of Figure 2 are typical of many. Exceptions to this general rule have however occasionally been found (93, 127, 214). Acidity is more apt to be important in substrate respiration e.g. the oxidation of phenylacetic acid by *Penicillium chrysogenum* (135) and of glucose by *Streptomyces griseus* (131). Since the internal pH of the intact cell is probably protected against all but the most extreme changes in the medium it seems likely that pronounced pH effects on respiration of undissociated compounds or of very weak acids signify that surface located enzymes are involved in the total process of metabolism.

Phosphate buffers are the most generally satisfactory for respirometric studies others particularly acetate are often toxic (Chapter 1).

Temperature Most respirometric studies of saprophytic fungi and actinomycetes are performed at moderate temperature 20–30°C. higher temperatures are more suitable for the fungi pathogenic to animals (215). Carbon dioxide evolution from growing mycelium of *Aspergillus niger* has a high Q_{10} at low temperature e.g. 3.12 over the range 7–17° the Q_{10} drops as the temperature is raised and is only 1.7 over the range 27–37° (217).

Oxygen and Carbon Dioxide Pressure The response of aerobic respiration to oxygen pressure varies with the organism studied (3, 72).





Figure 2 pH and the endogenous respiration of *Streptomyces coelicolor* (curve 1) *Myrothecium verrucaria* (curve 2) and *Fusarium solani* (curve 3) mycelium

Figure 3 The effect of oxygen pressure on the rate of respiration of *Aspergillus niger* Redrawn from Y Shu Journal of Agricultural and Food Chemistry Vol 1 p 1119 November 2 1953 Copyright 1953 by the American Chemical Society and reprinted by permission of the copyright owner



259 260 291) Respiration depends immediately upon dissolved oxygen which in turn is proportional to the partial pressure of oxygen in the gas phase. Finn (83) defines C_{it} as the concentration of dissolved oxygen above which cell respiration is independent of the oxygen concentration; data from several sources show that C_{it} for *Penicillium chrysogenum* is about 0.022 millimoles per liter (equivalent to an oxygen tension of 16 mm of mercury); that for yeast and *Escherichia coli* considerably lower. Amperometric measurements show that the rate of oxygen utilization is independent of oxygen pressure down to very low values (52). It follows therefore that the requirement of many fungi e.g. *Aspergillus niger* (Figure 3) for relatively high oxygen pressure for maximal respiratory rate means that other factors must be operative. Studies on the respiration of mycelium of *Myrothecium verrucaria* indicate that the rate of oxygen consumption by the rather active cells is limited in part by the rate at which oxygen reaches the interior of the cell mass (70).

The oxygen uptake of *Micromonospora vulgaris* is reduced by high oxygen pressure (315) but this effect is not usually encountered in the fungi.

Little evidence is available on the influence of carbon dioxide pressure on fungus respiration. Most respirometric techniques involve removal of metabolic carbon dioxide during measurement of oxygen uptake. In many bacteria oxygen uptake is in fact depressed by such removal of carbon dioxide (119). Two methods are available to determine if this occurs with any given organism: use of Warburg's indirect method (308) or use of diethanolamine as a carbon dioxide buffer (166 231).

Nutrients Potassium, phosphorus and magnesium are usually supplied to respiring cells although it is likely that endogenous reserves

of these and other minerals are sufficient. Aerobic oxidation with exogenous substrate is generally markedly increased if a source of nitrogen is provided (199, 265, 275); this of course converts the preparation at least potentially to a growing culture carrying on amino acid or even protein synthesis with an attendant demand upon energy and upon metabolic intermediates of the citric acid cycle. The effect of such organic materials as yeast extract may be the same but there is the added complication that respirable materials are also being supplied.

3 ENDOGENOUS RESPIRATION

Endogenous respiration or autorepiration defines the metabolism of the cell in the absence of an external substrate insofar as it is expressed by gas exchange. In yeast there is normally no anaerobic endogenous carbon dioxide liberation (270) although 2,4-dinitrophenol initiates fermentation of carbohydrate reserves (251). *Fusarium* on the other hand has a vigorous endogenous fermentation (98).

The substrate of endogenous respiration in fungi is not known; respiratory quotients are consistent with lipid oxidation (186, 230) but suitable analytical data to discriminate between this and other possibilities are not available. Cell free extracts of *Neurospora crassa* are stimulated in endogenous respiration by triphosphopyridine nucleotide (301).

Scopulariopsis brevicaulis cells are unable to assimilate nitrogen in the absence of exogenous substrate, i.e. the endogenous respiration cannot be coupled to amino acid or protein synthesis (186). This is in contrast to both *Chlorella vulgaris* (284) and *Torulopsis utilis* (332).

The high endogenous rate characteristic of many fungi and actinomycetes complicates experimental study; obviously if some limiting terminal system is fully saturated by the endogenous system then an addition of substrate will not cause any further increase in oxygen uptake. However it cannot be assumed that under these circumstances the exogenous substrate is inert—conceivably some of it will be oxidized in competition with the oxidation of endogenous materials. Partial responses are also possible: if for example the terminal system were 50 per cent saturated by endogenous metabolism addition of substrate could at most double the oxygen uptake rate. But—and this may be a real problem—one can imagine that in fact the substrate is competing on any equal terms with the endogenous metabolism for

the terminal system. Were this so, not 50 per cent but 75 per cent of the total metabolism would be substrate metabolism. put another way, the endogenous system would be suppressed 50 per cent.

Barker (11) proposed a method to determine whether or not substrate utilization does suppress endogenous respiration. The method relies essentially on the assumption that the oxygen uptake per mole of substrate consumed is not affected by the concentration of substrate. Application of this technique to *Penicillium chrysogenum* indicates that glucose does not suppress endogenous respiration (272). In yeast, on the other hand, suppression of endogenous respiration by acetate occurs with agar grown cells but not with cells grown in a liquid medium (322).

A second approach to this problem is to use cells the endogenous reserves of which have been labeled by cultivation on a carbon 14 substrate (37, 244). It is essential that all of the reserves be labeled by growth from a very early stage on the isotopic substrate; a short exposure to labeled acetate, for example, may not succeed in marking all of the reserves. This method has received only a limited trial to date (25, 63, 101, 128, 203) and it is of course open to the criticism that not enough is known of the composition and dynamics of the many metabolic pools involved. Experimentally, it is only necessary to determine what the total activity of carbon dioxide from labeled cells is and whether or not this activity is affected by the simultaneous oxidation of substrate.


Amino acids have been reported as affecting endogenous respiration both positively (17) and negatively (93). These findings cannot at present be interpreted since the presence of amino nitrogen in the system and the possibility of amino acid toxicity unduly complicate the picture.

Experimentally, it is often desirable or even essential to reduce the level of endogenous respiration. Starving the cells for a period of time prior to measuring their respiratory response to an exogenous substrate has been successful with some fungi and actinomycetes (31, 70, 316, 320). Others, however, lose substrate respiratory capacity during the starvation period (14, 134, 195). There is also the danger that the starvation period may result in developmental changes in the organism (145). Since endogenous respiration is affected by the composition of the growth medium (251), the better approach would seem to be an attempt to find a medium in which the cells do not accumulate reserve materials in large amounts (102). Many of the media commonly used for fungi are so concentrated that accumulation of such reserves is almost forced upon the organism.

4 OXIDATIVE ASSIMILATION

Oxidation of carbon compounds by many organisms is not accompanied by the oxygen uptake theoretically necessary to account for the known disappearance of substrate. This has been explained as the result of the assimilation of part of the utilized carbon into reserves presumably carbohydrate in nature of the cell (59).

Oxidative assimilation occurs in yeast (231-322), actinomycetes (254) and fungi (70-80, 253). Its extent may be rather high for *Penicillium chrysogenum* (272); the data fit the equation



Thus in this organism about 83 per cent of the glucose disappearing is assimilated.

Fermentative i.e. anaerobic assimilation is known in yeast and bacteria (297). In what is presumably the same process *Aspergillus niger* forms intracellular mannitol and carbohydrate anaerobically (255). Fermentative accumulation may account for the finding (154, 247) that *A. niger* forms less ethanol from glucose than the theoretical amount expected from a fermentation of the yeast type.

5 THE EMBDEN MEYERHOF PATHWAY

The anaerobic breakdown of glucose by yeast is diagrammed in part in Figure 4. This series of reactions is of very general although probably not universal occurrence in microorganisms (79-118). In muscle the reactions termed glycolysis are similar except that pyruvic acid is reduced to lactic acid in the final step and that the starting material is glycogen.

Since the filamentous fungi require oxygen for growth (Chapter 1) there has been a tendency to discount the possible importance of the Embden Meyerhof pathway. First it should be realized that the products of glucose metabolism by other routes may feed into the pathway, thus the phosphogluconate oxidation pathway as discussed later forms triose phosphate. Second the Embden Meyerhof pathway is not restricted to anaerobic conditions; it is entirely possible for a portion of it to function simultaneously with aerobic processes or to be coupled to terminal aerobic reactions. Finally some fungi seem to be able to ferment sugar anaerobically even though they require oxygen for growth.

Although in the bacteria there are routes of ethanol formation other




Figure 4 Schematic outline of the Embden Meyerhof pathway of yeast. Abbreviations: ATP = adenosine triphosphate; ADP = adenosine diphosphate; DPN = diphosphopyridine nucleotide; DPNH = reduced diphosphopyridine nucleotide. Enzymes are numbered as follows:

- | | |
|---------------------------------|--|
| 1 Hexokinase | 7 Phosphoglyceryl kinase |
| 2 Phosphohexoisomerase | 8 Phosphoglyceryl mutase |
| 3 Phosphofructokinase | 9 Enolase |
| 4 Aldolase | 10 Phosphoenolpyruvic transphosphorylase |
| 5 Triosephosphate isomerase | 11 Pyruvic carboxylase |
| 6 Triosephosphate dehydrogenase | 12 Alcohol dehydrogenase |

than the Embden Meyerhof we may consider ethanol production as at least suggestive of the pathway. Formation of ethanol from glucose is rather common among the fungi. Early work chiefly with species of *Aspergillus* *Fusarium* *Mucor* and *Penicillium* is reviewed by Wehnert (317) Czapek (69) and Foster (84). A number of the wood destroying basidiomycetes form detectable amounts of ethanol in culture (178 221 232 233 319) other ethanol producers include *Ashbya gossypii* (194) *Rhizopus* spp (121 310 314) *Diplodia tubericola* (121) *Dematium pullulans* (278) *Mortierella* spp (233) *Neurospora crassa* (232) *Stemphylium radicinum* (178) and probably several of the Fungi Imperfecti (20). The most active producers are in the genus *Fusarium* and as will be seen later a functional Embden Meyerhof pathway is present in *F. lini*.

Under the usual experimental conditions ethanol is oxidized concurrently with its formation so that an accurate stoichiometry is impossible (5 194 235). However under anaerobic conditions not permitting either growth or ethanol oxidation it has been possible to show that the production of alcohol and carbon dioxide from glucose follows the quantitative predictions of the Embden Meyerhof pathway (62 290 302). Species of a given genus vary considerably in their accumulation of alcohol under growth conditions this has been studied especially in *Aspergillus* (21 292 333).

Ethanol production by fungi is generally favored by restricted aeration (168 285 290) although even under conditions of rapid aeration some fungi form large amounts of alcohol (176 314). The accumulation of ethanol like that of other carbon containing metabolites is greatest in media relatively high in carbohydrate (146) and low in nitrogen and minerals (243).

Ethanol oxidation in *Fusarium lini* is a special case of a more general capacity to oxidize primary and secondary alcohols (111 112) ethanol is oxidized to acetaldehyde 2 butanol to 2 butanone and 2,3 butylene glycol to acetoin.

Acetaldehyde the precursor of ethanol in the yeast fermentation is formed sometimes in appreciable amounts by growing cultures or resting preformed mats of several fungi (16 22 27 171 176 211 221 255 257 279 286). Most of these forms are ethanol producers and the acetaldehyde may be formed either as a precursor of ethanol or during oxidation of it (112).

In the yeast fermentation small amounts of glycerol are formed more is produced in the presence of sulfite or at alkaline pH. The same observations have been made with fungi (211 287).


Pyruvic acid is frequently produced in isolable amounts by fungi

and actinomycetes (60 131 237 238 257) the presence of nitrite increases pyruvate formation by *Fusarium* spp (220) Added thiamine decreases pyruvate accumulation (273 323) presumably by reason of its function in diphosphothiamine the coenzyme of pyruvic carboxylase


The formation of ethanol acetaldehyde and pyruvic acid is consistent with the functioning of a pathway of the yeast type as is the appearance of other minor products (93 109 237 255 281) but does not prove the occurrence of the pathway in any sense

The occurrence of lactic acid among fungi has been reviewed in the preceding chapter Among the phycomycetes the only true fungi to form large amounts of lactic from glucose there appear to be two different types of metabolic pattern The lower phycomycetes so far investigated anaerobically convert virtually all of the glucose consumed to lactic acid (40 41 114 116) This immediately suggests comparison to the metabolism of muscle and of the homofermentative bacteria in both of which glucose is converted anaerobically to two moles of lactic acid via the Embden Meyerhof pathway The comparison must of course be only tentative in the absence of further data the demonstration of a diphosphopyridine linked lactic dehydrogenase would be of especial interest

Members of the genus *Rhizopus* constitute the second group of lactate forming fungi Under anaerobiosis glucose is converted to lactic acid in 50 per cent yield Waksman and Foster (310) suggest the over all reaction



The data of Table I show that *R. oryzae* ferments labeled glucose as expected from the Embden Meyerhof sequence that is the path of glucose carbon is presumably in outline



This implies that all products arise from a common pool of triose phosphate leaving open the problem of how the 1:1 ratio of lactic acid to ethanol (Equation 4) is maintained The conversion of pyruvate to

lactate has been shown (84) and carbon 14 data suggest that a direct reduction occurs (18) *Leuconostoc mesenteroides* yields the same products as *Rhizopus* spp but the mechanism of this heterolactic fermentation is entirely different (75)

Table 1 The Distribution of Glucose Carbon among the Fermentation Products of *Rhizopus oryzae**

Exp	Substrate	CO ₂	Ethanol		Lactic Acid		
			CH ₄	CH ₂ OH	COOH	CHOH	CH ₃
1	Glucose 1 C ¹⁴	—	—	—	0.5	0.5	16.0
2	Glucose 1 C ¹⁴	0.5	14.1	0.9	0.2	0.2	3.8
3	Glucose 3,4 C ¹⁴	13.0	0	0	13.2	0.35	0.5
4	Glucose 3,4 C ¹⁴	—	—	—	14	0.2	0.2

* Figures represent millimicrocuries of carbon 14 in each carbon atom. From M. Gibbs and R. Gastel, *Arch. Biochem. Biophys.* 43: 33-38 (1953) by permission of The Academic Press, Inc.

Purely enzymatic studies can establish only the possibility of the Embden Meyerhof pathway since most of the enzymes of the sequence are also used for disposing of the triose phosphate formed in the phosphogluconate oxidation pathway and since the enzymes even if present are not necessarily functional. Pyruvic carboxylase has been detected in several fungi (62, 88, 148, 230, 274). *Aspergillus niger* extracts contain virtually the complete battery of enzymes necessary for the conversion of glucose to ethanol and carbon dioxide (148); this is consistent with the data cited earlier indicating that some species of *Aspergillus* produce ethanol. Several of the same enzymes appear in extracts of *Penicillium chrysogenum* (264) and *P. notatum* (182). Aldolase, one of the enzymes which is characteristic of the Embden Meyerhof sequence, is found in *Aspergillus* spp. (7, 15a, 149, 150), *Neurospora crassa* (277), *Fusarium lini* (62), *Penicillium* spp. (182, 264), *Streptomyces coelicolor* (61) and *Tilletia caries* (212).

Ethanol production in *Fusarium* spp. has already been mentioned. Carbon balance studies, enzymatic analyses, and isotope distribution experiments agree that anaerobically the Embden Meyerhof pathway is functional and that no other anaerobic mechanism need be postulated (62, 125). Balance and fermentation data on *Ashbya gossypii* also implicate the yeast system (194).

Streptomyces coelicolor is unable to form either carbon dioxide or acids anaerobically; it contains, however, the enzymes necessary to convert glucose to pyruvic acid by way of the Embden Meyerhof pathway. The explanation for its failure to ferment seems to be that



no enzymes are present which can anaerobically metabolize pyruvate and regenerate diphosphopyridine nucleotide from its reduced form (61). This biochemical lesion occurs also in certain of the bacteria (132). These data suggest that failure of certain of the true fungi, e.g. *Mycrothecium verrucaria* to ferment glucose may ultimately be traced to the same enzymatic deficiency.

In summary the cumulative evidence for the Embden Meyerhof pathway in at least some of the filamentous fungi is impressive. There are however several different variations of the pattern. *Fusarium* spp. and probably *Aspergillus* spp. appear to resemble yeast most closely. The enzymatic basis of the two types of lactic acid formation in the phycomyces remains unknown and at present it is only on the basis of comparative biochemistry that we regard these as subtypes of the Embden Meyerhof sequence. Finally it is possible that non fermentative forms can utilize the pathway by coupling it to an aerobic process in which reduced coenzyme is regenerated by an oxidase system and pyruvate is metabolized via the citric acid cycle.

6 THE PHOSPHOGLUCONATE OXIDATION PATHWAY

Studies carried out during the decade 1930-1940 by Dickens, Lipmann and Warburg and more recently by Horecker, Racker and many others have established the occurrence of a second sequence of respiratory reactions variously known as the hexosemonophosphate shunt, the direct oxidative pathway and the pentose phosphate cycle. The system appears to be widespread in nature occurring in animals, plants and microorganisms (238).

For convenience and to avoid unnecessary detail we may, following Racker (236), divide the pathway into two phases. The first is an oxidative phase in which glucose 6-phosphate is transformed to pentose phosphate via 6-phosphogluconic acid. Carbon atom 1 of glucose, the aldehyde carbon, is converted to carbon dioxide in the process.



The key enzymes of this phase are glucose-6-phosphate dehydrogenase (Equation 6) and 6-phosphogluconic acid dehydrogenase (Equation 7); both are linked to triphosphopyridine nucleotide (TPN). Reac

lactate has been shown (84) and carbon 14 data suggest that a direct reduction occurs (48). *Leuconostoc mesenteroides* yields the same products as *Rhizopus* spp but the mechanism of this heterolactic fermentation is entirely different (75).

Table 1 The Distribution of Glucose Carbon among the Fermentation Products of *Rhizopus oryzae**

Exp	Substrate	CO ₂	Ethanol		Lactic Acid		
			CH ₃	CH ₂ OH	COOH	CHOH	CH ₃
1	Glucose 1 C ¹⁴	—	—	—	0.5	0.5	16.0
2	Glucose 1 C ¹⁴	0.5	14.1	0.9	0.2	0.2	3.8
3	Glucose 3,4 C ¹⁴	13.0	0	0	13.2	0.35	0.5
4	Glucose 3,4 C ¹⁴	—	—	—	14	0.2	0.2

* Figures represent millimicrocuries of carbon 14 in each carbon atom. From M. Gibbs and R. Gastel, *Arch. Biochem. Biophys.* 43: 33-38 (1953), by permission of The Academic Press, Inc.

Purely enzymatic studies can establish only the possibility of the Embden Meyerhof pathway since most of the enzymes of the sequence are also used for disposing of the triose phosphate formed in the phosphogluconate oxidation pathway and since the enzymes even if present are not necessarily functional. Pyruvic carboxylase has been detected in several fungi (62, 88, 148, 230, 271). *Aspergillus niger* extracts contain virtually the complete battery of enzymes necessary for the conversion of glucose to ethanol and carbon dioxide (148); this is consistent with the data cited earlier indicating that some species of *Aspergillus* produce ethanol. Several of the same enzymes appear in extracts of *Penicillium chrysogenum* (264) and *P. notatum* (182). Aldolase, one of the enzymes which is characteristic of the Embden Meyerhof sequence, is found in *Aspergillus* spp (7, 15a, 149, 150), *Neurospora crassa* (277), *Fusarium lini* (62), *Penicillium* spp (182, 264), *Streptomyces coelicolor* (61) and *Tilletia caries* (212).

Ethanol production in *Fusarium* spp has already been mentioned. Carbon balance studies, enzymatic analyses and isotope distribution experiments agree that anaerobically the Embden Meyerhof pathway is functional and that no other anaerobic mechanism need be postulated (62, 125). Balance and fermentation data on *Ashbya gossypii* also implicate the yeast system (191).

Streptomyces coelicolor is unable to form either carbon dioxide or acids anaerobically; it contains, however, the enzymes necessary to convert glucose to pyruvic acid by way of the Embden Meyerhof pathway. The explanation for its failure to ferment seems to be that

pathway. One group of methods is based upon the fact that operation of the phosphogluconate oxidation pathway results in a preferential liberation of the aldehyde carbon of glucose as carbon dioxide. The limitation of this approach is that glucose goes through many more metabolic pools and is more diluted in the Embden Meyerhof than in the oxidative pathway before liberation of carbon dioxide; this results in a bias which may exaggerate the contribution of the oxidative pathway. Application of various modifications of this method indicates that there is a substantial flow of carbon over some pathway preferentially liberating carbon 1 of glucose as carbon dioxide in *Penicillium chrysogenum* (80-124), *Aspergillus oryzae* (6), *Streptomyces coelicolor* (66) and *Neurospora crassa* (293). A modification of this method indicates that in growing mycelium of *Penicillium chrysogenum* about 60 per cent of the glucose carbon goes over the oxidative pathway (124) while in *Fusarium lini* a minimum of 17 per cent of the glucose catabolic mechanism involves preferential liberation of carbon 1 (125).

A more promising method involves determination of acetate, ethanol, pyruvate or compounds derived therefrom during aerobic respiration of specifically labeled glucose (26, 318). Application of this technique to *Penicillium chrysogenum* suggests that the Embden Meyerhof pathway is the principal route of glucose breakdown (172). Still other methods may be applicable but have not yet been tested in the fungi (18, 19, 23, 24, 105, 139, 156, 236, 268). Wood (327) discusses the methods critically and points out their limitations.


In *Aspergillus niger* the distribution of isotope in citric acid indicates that the Embden Meyerhof sequence accounts for 80 per cent of the total aerobic glucose catabolism (261). Similarly in *Ustilago zeae* the labeling of the carbohydrate moiety of the glucolipid ustilagic acid is consistent with the resynthesis of glucose from triose phosphate and consequently with the operation of the Embden Meyerhof pathway (29). The labeling of kojic acid formed by *Aspergillus flavus* from pentose is on the other hand suggestive of a preliminary conversion of pentose through the non-oxidative phase of the phosphogluconate pathway to hexose (8).

The various approaches which attempt to utilize labeled sugars each has its own difficulties and ambiguities and we should regard the data collected so far as preliminary and tentative. Some attempt should be made to see whether two different methods agree with regard to the same material before reliance can be placed on any single method. There seems little doubt that the phosphogluconate oxidation pathway is utilized by many fungi and actinomycetes and it is not unreasonable

tion 7 is probably the sum of two reactions. It is assumed that the aerobic reoxidation of reduced coenzyme (TPNH) occurs and that in the process the energy of the oxidation is utilized for synthesis of adenosine triphosphate or related compounds. The phosphogluconate pathway is an aerobic system; evidence for an analogous fermentative pathway is summarized by Gunsalus et al. (118).

The first pentose phosphate formed, i.e. the product of Equation 7, is ribulose 5 phosphate; this rapidly comes into enzymatic equilibrium with two other pentose phosphates, ribose 5 phosphate and xylulose 5 phosphate.

The second phase of the sequence is a complex series of non-oxidative conversions of the pentose phosphates into other phosphorylated sugars. Direct intermediates in this phase include sedoheptulose 7 phosphate, glyceraldehyde 3 phosphate, a tetrose-4 phosphate (probably erythrose-4 phosphate) and fructose 6 phosphate. Since glucose 6 phosphate is regenerated, this second phase may be summarized



The key enzymes in this phase are transketolase and transaldolase. In the presence of enzymes of the Embden-Meyerhof sequence, still other sugar phosphates, fructose 1,6-diphosphate and sedoheptulose 1,7-diphosphate, appear. In addition, a greater or less amount of the triose phosphate formed may be converted to pyruvic acid and thereby removed from the system under discussion.

It was natural to investigate the fungi for an aerobic system of this type, but so far comparatively little progress has been made toward an organized and coherent picture of the role of this cycle in the oxidative metabolism of the fungi. The evidence at hand includes three types of study: demonstration of particular enzymes; demonstration of one or more of the characteristic sugar phosphates; and study of the distribution of isotope from specifically labeled glucose.

The two dehydrogenases of the oxidative phase of the pathway have been identified in *Neurospora crassa* (213), *Tilletia caries* (212) and *Streptomyces coelicolor* (66). Extracts of *Aspergillus niger* contain glucose 6 phosphate dehydrogenase (147).

The conversion of pentose phosphate to hexose phosphate and the formation of sedoheptulose phosphate and other known intermediates are easily demonstrated in cell-free preparations of both fungi and actinomycetes (64, 212, 263).


A variety of methods depending on the formation of metabolites from carbon-14 labeled glucose have been devised in an attempt to assay the quantitative importance of the phosphogluconate oxidation.

The importance of the oxidation of free glucose is difficult to assess. A relatively large number of fungi form gluconic acid (Chapter 6) and we may reasonably assume that the same reaction is involved in all. However, the known glucose ferredoxin hydrogenase has been found only in the fungi mentioned above and in *Penicillium glaucum* apparently not occurring in *Aspergillus fumigatus*, *Mucor racemosus*, *Rhizopus nigricans* or *Dematiium pullulans* (92). As mentioned earlier, the availability of the energy of the reaction to the cell is uncertain; the hydrogen peroxide formed could, in the presence of catalase, serve as an oxidizing agent for other cell metabolites (157-291). The further metabolism of gluconic acid is similarly obscure; the likeliest possibilities are that the gluconate is phosphorylated and enters the hexose monophosphate pathway or that there is a continued non-phosphorylative metabolism after the pattern of the bacterial genera *Pseudomonas* and *Acetobacter* (297-329). Discovery of 2-keto-gluconic acid in small amounts in the culture fluid of *Penicillium brevis compactum* (Chapter 6) suggests that the bacterial pathway may be worth investigation.

8. PENTOSE FERMENTATION AND OXIDATION

The fermentation of pentoses by fungi has been studied primarily as it occurs in the genus *Fusarium*, although other fungi can carry out at least a slow fermentation of pentose (221). Early results and theories on the reaction are reviewed by Foster (84) and are of historical interest only.

Resting cells of *Fusarium lini* ferment xylose in accordance with the equation



With xylose 1- C^{14} as substrate, virtually all of the label appears in the methyl carbon of acetate (98). This system resembles that of *Lactobacillus* spp. (169) with the difference that the lactic acid characteristic of the bacterial fermentation is replaced by ethanol and carbon dioxide. In *Lactobacillus pentosus* preparations the pentose chain is split between carbons 2 and 3, yielding triose phosphate and acetyl phosphate (35-123). Presumably in *F. lini* the triose phosphate is then metabolized to ethanol and carbon dioxide—the enzymes necessary for this transformation are all present (62).


The pentose fermentation by *F. lini* is striking in another connection: there does not seem to be any conversion of pentose to hexose by the non-oxidative reactions of the phosphogluconate oxidation path-

to anticipate that for some it may be a major pathway of respiration or of pentose biogenesis. It is also clear that in some organisms both the oxidative and the Embden Meyerhof systems are present and presumably complete for glucose metabolism (64). It is to be expected that cells in different physiological states will strike different balances between the two systems.

7 THE NON PHOSPHORYLATIVE OXIDATION OF GLUCOSE

The single reaction of this type known is the oxidation of free glucose to gluconic acid (the first product is in fact glucono δ lactone) by the enzyme usually described as glucose oxidase or notatin. It is debatable whether this reaction can be classed as respiratory since we know neither the subsequent metabolic history of the gluconic acid nor the way in which energy from the oxidation can be used by the cell. However it seems likely that further study will reveal a genuine respiratory pathway.

The enzyme responsible for the oxidation of free glucose was discovered by D. Muller (205, 206) in *Aspergillus niger* and named glucose oxidase. Earlier results of Maximov (193) probably involve this enzyme. Interest in it was revived by the later discovery that an antibiotic effect of *Penicillium resticulosum* and *P. notatum* is traceable to the hydrogen peroxide generated by the reaction



As an antibiotic and before its identity with the enzyme was known the material was variously named penicillin B, notatin and penatin.

The enzyme is a flavoprotein with two flavin adenine dinucleotide groups per molecule (158). It is highly specific for β -D-glucose; purified preparations are virtually inactive toward other common sugars (160) although 2-deoxyglucose is oxidized (267). The specificity has been taken advantage of in a manometric assay for glucose (159) and in the demonstration of the mutarotase of *Penicillium notatum* (161).

Studies with isotopic oxygen show that the enzyme transfers hydrogen from glucose to oxygen (15). It is therefore a dehydrogenase rather than an oxidase and the name glucose aerodehydrogenase is to be preferred.

The enzyme differs from the glucose dehydrogenases of liver (120) and of *Pseudomonas fluorescens* (330), neither of which is a flavo protein. Claims that other glucose dehydrogenases are present in fungi (207, 224, 225) have been disputed (91) and it appears that only one such enzyme is involved in the organisms studied.




Figure 5 Schematic and unimplified outline of the citric acid cycle. Abbreviations: DPN = diphosphopyridine nucleotide; DPNH = reduced diphosphopyridine nucleotide; TPN = triphosphopyridine nucleotide; TPNH = reduced triphosphopyridine nucleotide. Enzymes are numbered as follows:

- | | |
|-----------------------------------|--------------------------|
| 1 Condensing enzyme | 5 Complex system |
| 2 Aconitase | 6 Succinic dehydrogenase |
| 3 Isocitric dehydrogenase | 7 Fumarase |
| 4 Oxalosuccinic decarboxylase (?) | 8 Malic dehydrogenase |

Free acetate may enter the cycle and be oxidized by 2 moles of oxygen.

The occurrence of the cycle is well established in animal cells, and there is good evidence that it is at least potentially functional in yeast, bacteria, and higher plants. There is, however, a real question whether its major metabolic importance is in supplying the carbon skeleton of amino acids (167-248) or whether it has an additional role in providing energy to the cell (74-283). With the probable exception of the oxidative decarboxylation of α -ketoglutaric acid, the phosphorylations associated with the citric acid cycle are at the cofactor level.

The organic acids of the citric acid cycle which are formed in substantial amounts by fungi include succinic, citric, fumaric, and acetic

way as described earlier for other fungi and actinomycetes and as is common in yeast and bacterial pentose metabolism (499)


The oxidative breakdown of pentose by fungi has not been intensively investigated. It has been mentioned in the preceding chapter that there is one report of the oxidation of pentoses to pentonic acids the more recent and better documented report (175) that *Pseudomonas* spp. form pentonic acids from pentose suggests that a re-examination of the problem in fungi would be of interest. *Streptomyces coelicolor* extracts oxidize xylose and ribose only if the cells have been grown on the particular sugar. Indirect evidence suggests that these pentoses must be phosphorylated and that the kinase responsible is an inducible enzyme not present in glucose-grown cells (66).

9 THE CITRIC ACID CYCLE

In animal tissues the functioning of a respiratory cycle—the citric acid cycle (also called the tricarboxylic acid cycle or the Krebs cycle)—is well established. The historical development of the concept has been reviewed by Krebs (165); other valuable general reviews are those of Wood (326) and Ochoa (222).

In Figure 5 the principal reactions of the citric acid cycle are outlined. It can be seen first that the net effect of one turn of the cycle is the complete oxidation of a 2-carbon fragment $\text{CH}_3\text{CO}-$ to carbon dioxide and water. This acetyl fragment is known to be attached to coenzyme A, forming acetyl coenzyme A. The first reaction in the cycle is the transfer of the acetyl fragment to oxaloacetic acid, forming citric acid. In subsequent reactions two molecules of carbon dioxide are split off and oxaloacetic acid is regenerated. In principle, therefore, only one molecule of oxaloacetate is required to initiate the cycle, after which the reaction becomes self-sustaining. In fact, of course, carbon is drawn off especially for the synthesis of amino acids, so that steady-state operation of the cycle under normal conditions presumably requires that 4-carbon dicarboxylic acids be formed *de novo* continually. Mechanisms are known in bacteria by which pyruvic acid, phosphoenolpyruvic acid, and propionic acid can be carboxylated by carbon dioxide fixation (118); in addition, the possibility of malate synthesis from acetate has been raised by work with bacteria (p. 227).

Viewed strictly as a respiratory system, the net effect of the cycle shown in Figure 5 is the oxidation of pyruvic acid:



the role of an interrupted tricarboxylic acid cycle in the biogenesis of citric acid (242)

Extensive studies of the incorporation of isotopic carbon into the citric acid molecule lend further and decisive support to the idea that bulk citric acid synthesis in *Aspergillus niger* occurs via the citric acid cycle. The labeling of citrate formed in the presence of $C^{14}O$ is consistent with fixation of carbon dioxide into oxalacetate and condensation of the oxalacetate with an acetyl fragment (191, 192, 198). Lewis and Weinhouse (174) showed experimentally that the symmetrical citric acid molecule is asymmetrically synthesized as first suggested by Ogston (223).

Experiments with the metabolism of labeled acetic acid (28, 49, 173) similarly implicate the known initial reactions of the citric acid cycle in the synthesis of citrate by *A. niger*. A study of the incorporation of glucose-3,4- C^{14} into citrate provides evidence that pyruvate arising from glucose is converted to oxalacetate and an acetyl fragment and that these condense to yield citrate with almost theoretical incorporation of carbon dioxide and little or no randomization of the carboxyl carbons of oxalacetate (58). The last observation is important in showing that oxalacetate does not equilibrate with a symmetrical acid like fumaric or succinic.

Although it has been suggested (85) that a condensation of 2-carbon fragments may yield the 4-carbon dicarboxylic acid skeleton for citrate synthesis, work with isotopic acetate indicates that this pathway is at best of minor significance (28, 173) and the studies of Cleland and Johnson (58) with labeled glucose afford no evidence at all for such a reaction.

For *Penicillium chrysogenum* there is good evidence of the formation of citrate from acetyl phosphate and oxalacetate and for the presence of enzyme systems oxidizing citrate, isocitrate, α -ketoglutarate, succinate and malate (51, 126, 229). A partially purified succinic dehydrogenase is inhibited by malonate (110) and acetate oxidation is depressed by the same poison (13). The incorporation of acetate carbon into glutamic acid conforms to that expected if the citric acid cycle is responsible for the metabolism of acetate (113).

Extracts of *Streptomyces coelicolor* carry out the major reactions of the cycle including the formation of citrate, the oxidation of α -ketoglutarate and other acids and the decarboxylation of oxalacetate (65). The incorporation of labeled carbon dioxide and acetate into amino acids by *S. griseus* is also consistent with the operation of the cycle (39, 104).

acids malic acid appears usually in smaller quantities (Chapter 6) Although their occurrence is suggestive of the cycle it is in no sense proof Nor can it be argued from the mere fact that intact cells oxidize these organic acids that the cycle is present

Alpha ketoglutaric acid is formed in detectable amounts by *Penicillium chrysogenum* from lactic acid (136) by *Aspergillus oryzae* from glucose (252) and by *A. niger* from endogenous reserves (311) Its role in glutamic acid synthesis is considered later (Chapter 8)

Individual reactions of the cycle have been studied especially in seven organisms *Blastocladiella emersonii* *Ashbya gossypii* *Neurospora crassa* *Aspergillus niger* *Penicillium chrysogenum* *Streptomyces coelicolor* and *Zygorhynchus moelleri* For some of these more or less data is also available from isotope distribution experiments

Most of the known enzymes of the cycle have been demonstrated in *Blastocladiella emersonii* as has also the formation of citrate (43 46 47) The cycle however appears to be at best weakly functional in this organism and development of the resistant sporangia is accompanied by loss of most of the enzyme activities of the cycle (43) Cantino and Horeinstein (45) have suggested that in *B. emersonii* a reductive succinate ketoglutarate isocitrate (S K I) cycle is operative the most striking feature of this proposal is that light which accelerates growth is visualized as providing the energy for the carboxylation of succinate and α ketoglutarate


In *Ashbya gossypii* acetate oxidation is accelerated by dicarboxylic acids citrate is synthesized from acetate and oxalacetate and citrate accumulation is increased by malonate (195)

Several of the characteristic enzyme activities of the cycle have been demonstrated in extracts of *Neurospora crassa* (82 213 258 275) Incorporation of carbon 14 from several substrates into aspartic and glutamic acids indicates that in this fungus the citric acid cycle plays a major role in the synthesis of these amino acids (248)

Aspergillus niger is of particular interest in that some strains accumulate citric acid (Chapter 6) and in that tracer studies have been made Enzymatic analyses are the most complete for any fungus and show that all of the known enzymes of the citric acid cycle are present in extracts made from mycelium (190 239 240 241) The enzyme systems acting upon tricarboxylic acids are rather complex there is evidence for an aconitase of the usual type an aconitic hydratase converting aconitate only to citrate (73 210) and two isocitric dehydrogenases (241) Excellent agreement between the enzymatic content of cells and the accumulation of citric acid affords further evidence for

The formation of 1 carbon dicarboxylic acids has often been ascribed to the Thunberg condensation solely on the basis that resting or growing cells given acetate form one or more of the dicarboxylic acids. By itself this observation is of no significance and cannot be offered as evidence for the Thunberg reaction.


The most effective evidence for some sort of Thunberg reaction in fungi comes from studies on the formation of fumaric acid by *Rhizopus nigricans* and consists of the finding that the organism forms fumarate from ethanol in yields so high as to make it unlikely that the citric acid cycle is the pathway responsible (89). The reaction may be written







As indicated in the equation the methyl carbons of ethanol appear as the methine carbons of fumarate. However in this instance the distribution of isotope is not of critical value i.e. the labeling results do not require us to postulate the Thunberg reaction (2, 293, 309). The entrance of formic acid into the methyl group of pyruvate also complicates the interpretation of the isotope data (151, 152).

It has been suggested that an acetate dehydrogenase of *Aspergillus fumigatus* carries out the Thunberg reaction (12) but the products of the reaction have not been isolated.

Studies on bacteria offer a possible model for the results on *Rhizopus nigricans*. The enzyme malate synthetase of *Escherichia coli* carries out the reaction (325)



Kornberg and Madsen (164) find the same enzyme in *Pseudomonas* sp and suggest the following sequence of reactions



This complex system requires condensing enzyme (15a) aconitase (15b) isocitritase (15c) some mechanism generating acetyl coenzyme A (15d) malate synthetase (15e) and malic dehydrogenase (15f). Al

Most of the enzymes of the cycle are demonstrable in extracts of *Zygorhynchus moelleri* (201) and isotope distribution experiments indicate that the cycle plays a major role in energy metabolism (202)


Apart from these more intensively studied species individual enzyme activities have been determined in many other fungi and actinomycetes. These include oxalacetic decarboxylase (87 306) α ketoglutaric acid oxidase (46) oxalosuccinic decarboxylase (183) enzyme systems oxidizing succinic acid (46 89 137 182 216) and isocitric dehydrogenase and condensing enzyme (12)

In conclusion it is evident that the enzymes of the citric acid cycle are of quite general occurrence among fungi and actinomycetes and indeed that further enzyme surveys can add little to our understanding. The more important question is the physiological role of the citric acid cycle. Two such roles may be envisaged: provision of energy to the cell and provision of carbon skeletons for amino acid synthesis. The second of these is reasonably certainly a real function of the cycle in *Neurospora crassa*, *Penicillium chrysogenum* and *Streptomyces griseus*. A role in energy metabolism is more difficult to establish but would be of great interest. Citric acid accumulation seems to result directly from an interruption of the cycle under special nutritional conditions at the citric acid stage.

We must also recall from Chapter 6 that isocitritase catalyzing the conversion of isocitrate to succinic and glyoxylic acids is known to occur in fungi; its activity may contribute to some of the complexities of the metabolic data.

10 DICARBOXYLIC ACIDS IN RESPIRATION


Apart from their role in the citric acid cycle it has long been speculated that the 4 carbon dicarboxylic acids (succinic, fumaric, malic and oxalacetic) may be components of another respiratory cycle in which acetate is oxidized to carbon dioxide. The initial reaction of such a cycle would be the reductive condensation of acetate first suggested by Thunberg (298)



Further reactions can be imagined whereby the succinate so formed is converted to 1 mole of acetate and 2 moles of carbon dioxide, the acetate re-entering the cycle. The net effect would be the oxidation of one mole of acetate for each turn of the cycle. Possible dicarboxylic acid cycles are described by Ajl (1). Ochoa (222) and Umbreit (307). Siegel (262) discusses energy relations.

The formation of 4 carbon dicarboxylic acids has often been ascribed to the Thunberg condensation solely on the basis that resting or growing cells given acetate form one or more of the dicarboxylic acids. By itself this observation is of no significance and cannot be offered as evidence for the Thunberg reaction.


The most effective evidence for some sort of Thunberg reaction in fungi comes from studies on the formation of fumaric acid by *Rhizopus nigricans* and consists of the finding that the organism forms fumarate from ethanol in yields so high as to make it unlikely that the citric acid cycle is the pathway responsible (89). The reaction may be written







As indicated in the equation the methyl carbons of ethanol appear as the methine carbons of fumarate. However in this instance the distribution of isotope is not of critical value i.e. the labeling results do not require us to postulate the Thunberg reaction (2 283 309) the entrance of formic acid into the methyl group of pyruvate also complicates the interpretation of the isotope data (151 152).

It has been suggested that an acetic dehydrogenase of *Aspergillus fumigatus* carries out the Thunberg reaction (12) but the products of the reaction have not been isolated.

Studies on bacteria offer a possible model for the results on *Rhizopus nigricans*. The enzyme malate synthetase of *Escherichia coli* carries out the reaction (325)



Kornberg and Madsen (164) find the same enzyme in *Pseudomonas* sp and suggest the following sequence of reactions



This complex system requires condensing enzyme (15a) aconitase (15b) isocitritase (15c) some mechanism generating acetyl coenzyme A (15d) malate synthetase (15e) and malic dehydrogenase (15f). At

though the accumulation of malate or of fumarate by such a system can be imagined the cyclical character of the system as written would not be maintained. Most of the enzymes required for Equation 13 are known in fungi and it will be of considerable interest to see if some such series of reactions participates in energy metabolism or in the synthesis of dicarboxylic acids. A comparable reaction system in *Tetrahymena pyriformis* converts succinate reversibly to acetyl phosphate (256).

11 CARBON DIOXIDE FIXATION

Evidence for a physiological role of carbon dioxide in the growth of fungi appears first in relatively crude experiments on *Fusarium* spp (180) retardation of early growth in the absence of carbon dioxide has been observed in many other fungi (215 246 250). The carbon dioxide requirement is enhanced by mineral deficiencies (269).

An apparent effect of carbon dioxide on dehydrogenations (130) has been shown to be exerted in fact on endogenous methylene blue reduction (97).

Carbon dioxide stimulation of growth could of course be ascribed to other phenomena than fixation. However in *Penicillium chrysogenum* there is good evidence that fixation is related to the essentiality of carbon dioxide (106). In this organism fixation is negligible unless both carbon and nitrogen sources are provided this situation coupled with the fact that carbon dioxide is not essential in a complex medium and that aspartic and glutamic acids are labeled very early by C^{14} from isotopic carbon dioxide suggests strongly that the physiological role of carbon dioxide is in the synthesis of essential amino acids via the citric acid cycle. Carbon dioxide fixation in *Neurospora crassa* is similarly enhanced by provision of nitrogen (275) and carbon (128).

It has been mentioned earlier in this chapter that under appropriate conditions there is extensive incorporation of carbon dioxide into citric acid. Labeled carbon dioxide is also incorporated into the carboxyl group of pyruvic acid by *Rhizopus nigricans* (86) in the carboxyl of lactate by *R. oryzae* (100) in oxalic acid by *Aspergillus niger* (174) in succinic acid by *Streptomyces coelicolor* (60) in streptomycin by *S. griseus* (144) and in unidentified cell constituents by several fungi (181).

In organisms generally there are several different pathways by which carbon dioxide is fixed (10 118 309). Indirect and not entirely conclusive evidence suggests that fixation occurs into oxalacetate in

Rhizopus nigricans (86) and *Neurospora crassa* (276) We have already noted the proposal that the light stimulated carbon dioxide fixation in *Blastocladiella emersonii* reflects the reductive carboxylation of succinate and α ketoglutarate (45) The pathway of fixation can presumably only be detected with finality in experiments with active cell free preparations

12 TERMINAL RESPIRATION

In aerobic organisms the ultimate oxidizing agent is molecular oxygen the reaction or sequence of reactions by which the electrons withdrawn from the substrate are transferred to oxygen is denoted terminal respiration Conventionally the process is visualized as transfer of hydrogen from substrate to oxygen The transfer is mediated by a system in which substances of increasingly positive oxidation reduction potential form a chain of carriers

The simplest system involves only one carrier the flavoprotein prosthetic group of an enzyme examples are the amino acid oxidases and glucose aerodehydrogenase More usually the pyridine nucleotides—diphosphopyridine nucleotide or triphosphopyridine nucleotide—are the first hydrogen carriers of the sequence Perhaps the most common terminal pathway involves a pyridine nucleotide a flavoprotein and the cytochrome system in succession that is the flow of hydrogen (electrons) is over the pathway

Substrate \rightarrow pyridine nucleotide \rightarrow flavoprotein \rightarrow cytochromes
 \rightarrow oxygen (16)

The cytochrome system is complex and there is considerable doubt whether what is known of animal systems can be applied directly to microorganisms The flavoprotein is probably always a metallo enzyme (141)

Under some conditions e.g. a high partial pressure of oxygen flavo proteins may react directly with oxygen without the intervention of the cytochrome system (295)

The cytochrome system appears to be general in the fungi as judged in the first instance by spectroscopic observations (32 33 56 117 122 197 227 291 300) However it should be noted that experience with bacterial cytochromes indicates that it is not possible from simple spectroscopic measurements to state with precision whether the components of a new system are the same as those of the more familiar yeast and animal systems (266 321) Consequently it is well to reserve judgment for the time being as to whether the fungal cyto

chromes are the same as those of yeast and to be content with the generalization that cytochromes are generally present in fungi. The cytochrome *c* of *Ustilago sphaerogena* for example has a spectrum like beef heart cytochrome *c* but has a higher molecular weight and a lower isoelectric point (208). In this organism too the conditions of culture especially the supply of zinc and of thiamine markedly affect the concentrations of the cytochromes (117).

Exogenously supplied cytochrome from animal sources functions as an electron carrier for the aerobic lactic dehydrogenase of *Penicillium chrysogenum* (50) and for the succinoxidase systems of all forms tested (13 190 258).

Cytochrome oxidase may be assayed either manometrically or spectrophotometrically (Figure 6) usually with animal cytochrome *c* as substrate. The enzyme operationally defined by these assays seems to be of universal occurrence in the fungi; experiments have covered representatives of all of the major taxonomic groups (31 32 33 46 47 57 71 117 213 230 300). As in other organisms cytochrome oxidase activity is associated with the insoluble and sedimentable fraction of homogenates (57 71 122 258 299). In bacteria this fraction is apparently the cytoplasmic membrane or a similar structure (266).

The flavin components of the terminal system which proceeds over the cytochromes (Equation 16) have not been systematically investigated. Reductases linking cytochrome *c* with the pyridine nucleotide coenzymes and with succinate occur in *Neurospora tetrasperma* (57).




Figure 6 Cytochrome oxidase in *Blastocladiella* sp. Curve 1 in presence of cyanide curve 2 action of the enzyme. Redrawn from E. C. Cantino.

The Role of Metabolism and alpha ketoglutarate Oxidase in the Growth and Differentiation of the Aquatic Phycomycete *Blastocladiella* *Emersonii*. Transactions of The New York Academy of Sciences Series II Vol 15 No 5 p 159 163 (1953) by permission of the New York Academy of Science.

It is of some interest whether the cytochrome system is responsible for the entire terminal respiration or whether other systems may also be important. Elsewhere (Chapter 6) it has been noted that a role of the polyphenol oxidases or of quinone reductases is not yet established. The other alternative is a flavin system and this possibility has received some attention.

The cytochrome system is poisoned by cyanide, azide, and carbon monoxide. In general, the aerobic respiration of intact cells of fungi is also inhibited by these poisons (31, 108, 189, 191, 324). However, both in *Myrothecium verrucaria* (71) and *Ustilago sphaerogena* (117) cell respiration is not affected by cyanide or carbon monoxide at concentrations which inhibit cytochrome oxidase. This suggests—although there are other possibilities—that some system other than the cytochrome is operative in intact cells. Cells of *Gelasinospora tetrasperma* on the other hand are as sensitive to these poisons as is the extracted oxidase (31); in this and in some other fungi (32–57) the activity of the cytochrome system is sufficient to explain mycelial respiration.

There is some possibility that slow growing strains of *Neurospora crassa* possess a terminal respiratory system other than the cytochromes. The poky strain, the distinctive features of which are inherited cytoplasmically, is relatively insensitive to cytochrome inhibitors, more responsive than the normal strain to oxygen pressure, and has a high content of flavin adenine dinucleotide (122, 301). A similar but gene determined strain lacks cytochrome oxidase activity and is insensitive to azide and cyanide (300). Finally, although the poky character is suppressed by a nuclear gene, the cytochrome system is not increased along with the return to normal growth rate; this suggests that the effect of the gene is to open up some pathway alternative to the cytochrome route of terminal respiration (196). These observations all point toward a terminal system involving flavin enzymes.

Tamaya (291) summarizes the Japanese literature which suggests that in *Aspergillus oryzae* the respiration of mycelium from submerged (aerated or shaken) culture is mediated by the cytochromes, that of mycelium from still culture by some other, possibly flavin system. This perhaps merits reexamination with the improved methods now available, although there are some contradictions (313).

Catalase, an iron-containing enzyme of the hydroperoxidase group (296), is present in all fungi which have been examined for it. Catalyzing the breakdown of hydrogen peroxide, the enzyme may serve only to protect the organism against hydrogen peroxide or may have a broader function in the oxidation of organic molecules (34, 157). The

catalase of *Aspergillus oryzae* appears in the medium during autolysis (67) and the enzyme may have more than one component (68-153). Peroxidases which utilize hydrogen peroxide for the oxidation of other compounds are especially active in some of the wood destroying basidiomycetes (184-185). *Neurospora tetrasperma* contains a cytochrome c peroxidase (57).

BIBLIOGRAPHY

- 1 Aji S J 1951 *Bacteriol Revs* 15 211-244
- 2 Aji S J and D T O Wong 1955 *Arch Biochem Biophys* 54 474-485
- 3 Allen P J and W H Price 1950 *Am J Botany* 37 393-400
- 4 Altermatt H A F J Simpson and A C Neish 1955 *Can J Biochem and Physiol* 33 615-621
- 5 Anderson A K 1924 *Stud Biol Sci Univ Minnesota* 5 237-280
- 6 Arnstein H R V and R Bentley 1953a *Biochem J (London)* 51 493-508
- 7 Arnstein H R V and R Bentley 1953b *Biochem J (London)* 51 508-516
- 8 Arnstein H R V and R Bentley 1956 *Biochem J (London)* 62 403-411
- 9 Ballo A C Casinori and G Serlupi Crescenzi 1956 *Biochim et Biophys Acta* 20 414-415
- 10 Bandurski R S 1955 *J Biol Chem* 217 137-150
- 11 Barker H A 1956 *J Cellular Comp Physiol* 8 231-250
- 12 Barron E S G and F Chretien 1953 *Biochim et Biophys Acta* 10 239-249
- 13 Beevers H E P Goldschmidt and H Kofler 1952 *Arch Biochem Biophys* 39 236-238
- 14 Bentley M L 1953 *J Gen Microbiol* 8 365-377
- 15 Bentley R and A Neuberger 1949 *Biochem J (London)* 43 584-590
- 15a Bentley R and C P Thiessen 1957 *J Biol Chem* 226 689-700
- 16 Bernhauer K and H Theelen 1932 *Biochem Z* 253 30-36
- 17 Bernheim F 1940 *J Bacteriol* 44 555-559
- 18 Bernstein I A 1956 *J Biol Chem* 221 873-878
- 19 Bernstein I A et al 1955 *J Biol Chem* 215 137-152
- 20 Birkinshaw J H J H V Charles A C Heitherington and H Raistrick 1931 *Phil Trans Roy Soc (London) Ser B* 220 99-125
- 21 Birkinshaw J H J H V Charles H Raistrick and J A R Staley 1931 *Phil Trans Roy Soc (London) Ser B* 220 27-54
- 22 Birkinshaw J H and H Raistrick 1931 *Phil Trans Roy Soc (London) Ser B* 220 331-353
- 23 Bloom B 1955 *J Biol Chem* 215 467-472
- 24 Bloom B F Eisenberg Jr and D Steffen Jr 1955 *J Biol Chem* 215 461-466
- 25 Blumenthal H J H Kofler and E P Goldschmidt 1950 *Science* 116 475-477
- 26 Blumenthal H J A F Lewis and S Weinhouse 1954 *J Am Chem Soc* 76 6093-6097
- 27 Bolcato V and P Tono 1939 *Enzynologia* 7 146-156
- 28 Bornstein R A and M J Johnson 1952 *J Biol Chem* 198 143-153
- 29 Boothroyd B J A Thorn and R H Haskins 1955 *Can J Biochem and Physiol* 33 289-296
- 30 Bose S 1917 *J Indian Chem Soc* 24 307-337
- 31 Boulter D 1957 *J Gen Microbiol* 16 305-316

- 32 Boulter D and A Burges 1955 *Experientia* 11 188-189
- 33 Boulter D and E Derbyshire 1957 *J Exp Botany* 8 313-318
- 34 Brown D H and E C Cantino 1955 *Am J Botany* 42 337-341
- 35 Burma D F and B L Horecker 1957 *Biochim et Biophys Acta* 24 660-661
- 36 Burnet J H 1953 *New Phytologist* 52 58-64
- 37 Burns R H 1949 In W W Umbreit R H Burris and J F Stauffer (eds) *Manometric Techniques and Tissue Metabolism* 2nd ed p 96-97 Minneapo-
lis Burgess Publishing Co
- 38 Buston H W and A H Khan 1956 *J Gen Microbiol* 14 655-660
- 39 Butterworth E M C M Gilmour and C H Wang 1955 *J Bacteriol* 63 725-727
- 40 Cantino E C 1949 *Am J Botany* 36 95-110
- 41 Cantino E C 1951 *Antonie van Leeuwenhoek J Microbiol Serol* 14 59-96
- 42 Cantino E C 1953 *Trans N Y Acad Sci* 15 159-163
- 43 Cantino E C 1956 *Mycologia* 48 225-10
- 44 Cantino E C and E A Horenstein 1955 *Physiol Plantarum* 8 189-211
- 45 Cantino E C and E A Horenstein 1956 *Mycologia* 48 777-799
- 46 Cantino E C and M T Hyatt 1953a *Am J Botany* 40 688-694
- 47 Cantino E C and M T Hyatt 1953b *J Bacteriol* 66 712-720
- 48 Carson S F J W Foster W E Jefferson E F Phares and D S Anthony 1951 *Arch Biochem Biophys* 33 418-458
- 49 Carson S F F H Mosbach and E F Phares 1951 *J Facteriol* 6 235-38
- 50 Casida L E Jr and S G Knight 1954a *J Facteriol* 67 171-175
- 51 Casida L E Jr and S G Knight 1954b *J Bacteriol* 67 68-661
- 52 Chain E B and G Guiland 1954 *Rend ist sup r s ita* (English ed) 17 5-60
- 53 Challenger F L Klein and T K Walker 1931 *J Chem Soc (London)* 1931 16-23
- 54 Chance B 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 2 (1) p 48-453 New York Academic Press
- 55 Chattaway F W C C Thompson and A C F Parlow 1954 *Biochim et Biophys Acta* 14 583-584
- 56 Chauvet J 1943 *Enzymologia* 10 57-69
- 57 Cheng S C 1954 *Plant Physiol* 29 458-467
- 58 Cleland W W and M J Johnson 1954 *J Biol Chem* 208 649-689
- 59 Clifton C E 1952 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 2 (2) p 912-928 New York Academic Press
- 60 Cochrane V W 1955 *J Bacteriol* 63 459-461
- 61 Cochrane V W 1955 *J Bacteriol* 69 256-263
- 62 Cochrane V W 1956 *Mycologia* 48 1-12
- 63 Cochrane V W and M Gibbs 1951 *J Bacteriol* 61 305-307
- 64 Cochrane V W and P L Hawley 1956 *J Bacteriol* 71 308-314
- 65 Cochrane V W and H D Peck Jr 1953 *J Bacteriol* 66 37-44
- 66 Cochrane V W H D Peck Jr and A Harrison 1953 *J Bacteriol* 66 17-23
- 67 Crewther W G and F G Lennox 1953a *Australian J Biol Sci* 6 410-427
- 68 Crewther W G and F G Lennox 1953b *Australian J Biol Sci* 6 428-436
- 69 Crapek F 1955 *Biochemie der Pflanzen* 3rd ed Vol 1 p 316-338 Jena
Gustav Fischer
- 70 Darby R T and D R Goddard 1950a *Am J Botany* 37 39-387
- 71 Darby R T and D R Goddard 1950b *Physiol Plantarum* 3 43-446
- 72 DeBoer S R 1958 *Rec trav Botan Neerl* 25 117-240

- 73 Deffner M 1942 *Ann Chem Liebigs* 522 191-202
- 74 DeMoss J A H E Swim and L O Krampitz 1955 *Bacteriol Proc (Soc Am Bacteriologists)* 1955 114
- 75 DeMoss R D 1953 *J Cellular Comp Physiol* 41 Suppl 1 207-204
- 76 Dickens F and H McIlwain 1938 *Biochem J (London)* 32 1615-1625
- 77 Dixon M 1951 *Manometric Methods* 3rd ed Cambridge Cambridge University Press pp 165
- 78 Edwards G A C B Buell and W H Weston 1947 *Am J Botany* 34 551-555
- 79 Elsdon S R 1952 In J B Sumner and K Myrback (eds) *The Enzymes Vol 2 (2)* p 791-843 New York Academic Press
- 80 deFiebre C W and S G Knight 1953 *J Bacteriol* 66 170-172
- 81 Fincham J R S 1951 *J Gen Microbiol* 5 793-806
- 82 Fincham J R S and J B Boylen 1957 *J Gen Microbiol* 16 438-448
- 83 Finn R K 1954 *Bacteriol Revs* 18 254-274
- 84 Foster J W 1949 *Chemical Activities of Fungi* New York Academic Press pp 648
- 85 Foster J W and S F Carson 1950 *Proc Natl Acad Sci U S* 36 219-229
- 86 Foster J W and J B Davis 1948 *J Bacteriol* 56 329-338
- 87 Foster J W and J B Davis 1949 *Arch Biochem* 21 135-149
- 88 Foster J W and F W Denison Jr 1950 *Nature (London)* 166 833-834
- 89 Foster J W et al 1949 *Proc Natl Acad Sci U S* 35 663-672
- 90 Foulkes E C 1954 *J Bacteriol* 68 505
- 91 Franke W 1944 *Ann Chemie Liebigs* 555 111-132
- 92 Franke W and M Deffner 1939 *Ann Chem Liebigs* 541 117-150
- 93 Ganguly S and S C Roy 1955 *Arch Biochem Biophys* 59 45-51
- 94 Ganguly S and S C Roy 1956 *Arch Biochem Biophys* 63 26-31
- 95 Garner H H et al 1953 *Am J Botany* 40 289-296
- 96 Gentile A C 1954 *Plant Physiol* 29 257-261
- 97 Gest H and J L Stokes 1952 *Antonie van Leeuwenhoek J Microbiol Serol* 18 55-69
- 98 Gibbs M V W Cochrane L M Paegle and H Wolin 1954 *Arch Biochem Biophys* 50 237-242
- 99 Gibbs M J M Earl and J L Ritchie 1955 *J Biol Chem* 217 161-168
- 100 Gibbs M and R Gastel 1953 *Arch Biochem Biophys* 43 33-38
- 101 Gibbs M and W A Wood 1952 *Bacteriol Proc (Soc Am Bacteriologists)* 1952 11
- 102 Giese A C and E L Tatum 1946 *Arch Biochem* 9 1-13
- 103 Gillespie R J G A Maw and C A Vernon 1953 *Nature (London)* 171 1147-1149
- 104 Gilmour C M E M Butterworth E P Noble and C H Wang 1955 *J Bacteriol* 69 719-724
- 105 Gilvarg C 1952 *J Biol Chem* 199 57-64
- 106 Gitterman C O and S G Knight 1952 *J Bacteriol* 64 223-231
- 107 Goddard D R 1945 *Science* 101 352-353
- 108 Goddard D R and P E Smith 1938 *Plant Physiol* 13 241-264
- 109 Godin P 1953 *Biochim et Biophys Acta* 11 119-121
- 110 Godzieski C and R W Stone 1955 *Arch Biochem Biophys* 59 135-144
- 111 Goepfert G J 1941 *J Biol Chem* 140 525-534
- 112 Goepfert G J and F F Nord 1942 *Arch Biochem* 1 289-301
- 113 Goldschmidt E P I Yall and H Koffler 1956 *J Bacteriol* 72 436-446
- 114 Golueke C G 1957 *J Bacteriol* 74 337-344
- 115 Gottlieb D 1953 *Intern Cong Microbiol 6th Congr Rome* 5 122-136
- 116 Gregg J H 1950 *J Exp Zool* 114 173-196
- 117 Grumm P W and P J Allen 1954 *Plant Physiol* 49 369-377

- 118 Gunsalus I C B L Horecker and W A Wood 1955 *Bacteriol Revs* 19 9-178
- 119 Harris J O 1953 *Bacteriol Proc (Soc Am Bacteriologists)* 1953 90
- 120 Harrison D C 1931 *Biochem J (London)* 25 1016-1077
- 121 Harter L L and J L Weimer 1921 *J Agr Research* 21 11-226
- 122 Haskins F A A Tissieres H K Mitchell and M B Mitchell 1953 *J Biol Chem* 200 819-826
- 123 Heath E C J Hurwitz and B L Horecker 1956 *J Am Chem Soc* 78 5449
- 124 Heath E C and H Koffler 1956 *J Bacteriol* 71 174-181
- 125 Heath E C D Nasser and H Koffler 1956 *Arch Biochem Biophys* 64 80-87
- 126 Heberling R L and R W Stone 1957 *Bacteriol Proc (Soc Am Bacteriologists)* 1957 127
- 127 Henderson M K 1956 *J Gen Microbiol* 14 681-691
- 128 Hepler J Q and E L Tatum 1954 *J Biol Chem* 208 489-491
- 129 Herzog R O and A Meier 1908 *Z physiol Chem Hoppe Seyler's* 57 35-49
- 130 Hes J W 1938 *Nature (London)* 141 647
- 131 Hida T 1935 *J Shanghai Sci Inst Sect IV* 1 01-14
- 132 Hill R L and R C Mills 1954 *Arch Biochem Biophys* 53 174-183
- 133 Hochster R M and J H Quastel 1954 *Arch Biochem Biophys* 56 132-146
- 134 Hockenhull D J D A H Fantes M Herbert and B Whitehead 1954 *J Gen Microbiol* 10 353-370
- 135 Hockenhull D J D A D Walker G D Wilkin and F G Winder 1955 *Biochem J (London)* 50 605-609
- 136 Hockenhull D J D G D Wilkin and F C Winder 1951 *Nature (London)* 168 1013
- 137 Holter H and B M Pollock 1955 *Compt rend trav lab Carlsberg Ser chim* 3 221-243
- 138 Horecker B L 1953 *Bewers Dig* 28 214-219
- 139 Horecker B L and A H Mehler 1955 *Ann Rev Biochem* 24 207-274
- 140 Horowitz N H 1911 *J Biol Chem* 154 141-149
- 141 Huennekens G M 1956 *Experientia* 12 1-6
- 142 Hughes D E 1951 *Brit J Exp Pathol* 3 91-109
- 143 Hugo W B 1954 *Bacteriol Rev* 18 81-105
- 144 Hunter G D M Herbert and D J D Hockenhull 1954 *Biochem J (London)* 58 219-251
- 145 Ingraham J L and R Emerson 1954 *Am J Botany* 41 146-159
- 146 Jacquot R and R Raoux 1913 *Compt rend (Paris)* 216 318-319
- 147 Jagannathan V P N Rangachari and M Damodaran 1956 *Biochem J (London)* 64 177-181
- 148 Jagannathan V and K Singh 1953 *Enzymologia* 16 150-160
- 149 Jagannathan V and K Singh 1954 *Biochim et Biophys Acta* 15 134
- 150 Jagannathan V K Singh and M Damodaran 1956 *Biochem J (London)* 63 91-103
- 151 Jefferson W E and J W Foster 1953 *J Bacteriol* 6 587-59
- 152 Jefferson W E J W Foster L F Phares and S F Carson 1955 *J Am Chem Soc* 77 1477-1488
- 153 Jermy M A 1953 *Australian J Biol Sci* 6 77-97
- 154 Johnson E M L C Knight and T K Walker 1951 *Biochem J (London)* 51 903-908
- 155 Karrer P and F Haab 1918 *Helv Chim Acta* 31 9-198
- 156 Katz J S Abraham R Hill and I L Chalkoff 1955 *J Biol Chem* 214 857-868

- 157 Keilin D and E F Hartree 194₁ *Biochem J (London)* 39 293-301
- 158 Keilin D and E F Hartree 1948a *Biochem J (London)* 4^o 291-299
- 159 Keilin D and E F Hartree 1948b *Biochem J (London)* 42 230-238
- 160 Keilin D and E F Hartree 194₂a *Biochem J (London)* 50 331-341
- 161 Keilin D and E F Hartree 1942b *Biochem J (London)* 50 311-318
- 162 Kira D A and W H Peterson 1943 *J Biol Chem* 203 861-868
- 163 Koffler H R L Emerson D Perlman and R H Burris 1943 *J Bacteriol* 50 517-548
- 164 Kornberg H L and N B Madsen 1947 *Biochim et Biophys Acta* 94 641-643
- 165 Krebs H A 1940 *The Harvey Lectures 1948-49* p 163-199 Springfield Ill C C Thomas
- 166 Krebs H A 1941 *Biochem J (London)* 48 349-359
- 167 Krebs H A S Gurin and L V Eggleston 1942 *Biochem J (London)* 51 614-628
- 168 Kusnetsov S J 1932 *Microbiologiya* 1 3-18 83
- 169 Lampen J O 1953 *J Cellular Comp Physiol* 41 Suppl 1 183-20
- 170 Lattes G G 1949 *Arch Biochem* 22 8-15
- 171 Leitch H and J J Willaman 1926 *Phytopathology* 16 911-919
- 172 Lewis K H J Blumenthal C E Wenner and S Weinhouse 1944 *Federation Proc* 13 2,2
- 173 Lewis K F and S Weinhouse 1951a *J Am Chem Soc* 73 2500-2503
- 174 Lewis K F and S Weinhouse 1951b *J Am Chem Soc* 73 2906-2909
- 175 Lockwood L B and G E N Nelson 1946 *J Bacteriol* 52 581-586
- 176 Lockwood L B J J Stubbs and C J Senseman 1933 *Zentr Bakteriol Parasitenk Abt II* 98 167-171
- 177 Loewy A G 1942 *J Cellular Comp Physiol* 40 127-156
- 178 Lopez Ramos B and W J Schubert 1945 *Arch Biochem Biophys* 5 566-577
- 179 Lueders H R Kuhles and H Fink 1930 *Biochem Z* 917 243-248
- 180 Lundegårdh H 1923 *Botan Notiser* 1923 25-32
- 181 Lynch V H and M Calvin 1952 *J Bacteriol* 63 524-531
- 182 Lynen F and H P Hoffmann Walbeck 1948 *Ann Chem Liebigs* 549 143-168
- 183 Lynen F and H Scherer 1948 *Ann Chem Liebigs* 560 183-190
- 184 Lyr H 1944 *Planta* 46 408-413
- 185 Lyr H 1956 *Planta* 48 239-264
- 186 MacMillan A 1946 *Physiol Plantarum* 9 533-543
- 187 Mandels G R and A B Norton 1948 *Research Rept Quartermaster Gen Labs Microbiol Ser II* 1-50 (mimeographed)
- 188 Mandels G R and R G H Siu 1940 *J Bacteriol* 60 249-26
- 189 Mann T 1944 *Biochem J (London)* 38 339-345
- 190 Martin S M 1944 *Can J Microbiol* 1 6-11
- 191 Martin S M and P W Wilson 1941 *Arch Biochem Biophys* 32 150-15
- 192 Martin S M P W Wilson and R H Burris 1940 *Arch Biochem* 26 103-111
- 193 Maximov N A 1904 *Ber deut botan Ges* 22 225-235
- 194 Mickelson M N 1940 *J Bacteriol* 59 649-666
- 195 Mickelson M N and M N Schuler 1953 *J Bacteriol* 65 297-301
- 196 Mitchell M B and H K Mitchell 1946 *J Gen Microbiol* 14 84-89
- 197 Mitchell M B H K Mitchell and A Tissieres 1943 *Proc Natl Acad Sci U S* 39 606-613
- 198 Mosbach E H E F Phares and S F Carson 1942 *Arch Biochem Biophys* 34 435-442
- 199 Moses V 1944 *Biochem J (London)* 57 547-556

- 900 Moses V 195 a *Ann Botany* 19 211-223
- 901 Moses V 195b *J Gen Microbiol* 13 235-251
- 902 Moses V 1957 *J Gen Microbiol* 16 534-549
- 903 Moses V and P J Syrett 1955 *J Bacteriol* 70 201-204
- 904 Mukherjee S 1951 *Arch Biochem Biophys* 33 364-366
- 905 Muller D 1956 *Chem Ztg* 50 101
- 906 Muller D 1958 *Biochem Z* 199 136-170
- 907 Muller D 1941 *Enzymologia* 10 40-47
- 908 Neilands J B 1952 *J Biol Chem* 197 701-708
- 909 Neilson N E 1955 *Biochim et Biophys Acta* 17 139-140
- 910 Neilson N E 1956 *J Bacteriol* 71 356-361
- 911 Neuberg C and C Cohen 1951 *Biochem Z* 125 204-224
- 912 Newburgh R W C A Claridge and V H Cheldelin 1955 *J Biol Chem* 214 27-35
- 913 Nicholas D J D A Nason and W D McElroy 1951 *J Biol Chem* 207 341-351
- 914 Nickerson W J and J B Chadwick 1946 *Arch Biochem* 10 81-100
- 915 Nickerson W J and G A Edwards 1949 *J Gen Physiol* 33 41-55
- 916 Nickerson W J and R R Motan 1953 *Intern Congr Microbiol 6th Congr Rome* 5 137-146
- 917 Nielsen N and D Dresden 1939 *Compt rend trav lab Carlsberg Sér physiol* 22 287-300
- 918 Nord F F E Dammann and H Hofstetter 1936 *Biochem Z* 285 241-269
- 919 Nord F F H Hofstetter and E Dammann 1937 *Biochem Z* 293 231-55
- 920 Nord F F and R P Mull 1945 *Advances in Enzymol* 5 165-0
- 921 Nord F F and L J Sciarini 1946 *Arch Biochem* 9 419-437
- 922 Ochoa S 1955 *The Harvey Lectures 1950-51* p 153-180 Springfield Ill C C Thomas
- 923 Ogston A G 1948 *Nature (London)* 16 963
- 924 Ogura Y 1939 *Acta Phytochim (Japan)* 11 197-143
- 925 Ogura Y 1951 *J Biochem (Tokyo)* 38 75-81
- 926 Ogura Y and M Nagahisa 1937 *Botan Mag (Tokyo)* 54 597-614
- 927 Ohta J 1954 *J Biochem (Tokyo)* 41 489-497
- 928 Oliver I T and J L Peel 1956 *Biochim et Biophys Acta* 20 390-399
- 929 Olson J A 1954 *Nature (London)* 174 695-696
- 930 Owens R G 1955 *Contribs Boyce Thompson Inst* 18 1 13-144 145-159
- 931 Pardee A B 1949 *J Biol Chem* 179 1083-1091
- 932 Perlman D 1949 *Am J Botany* 36 180-181
- 933 Perlman D 1950 *Am J Botany* 37 937-941
- 934 Pickett M J and C E Clifton 1943 *J Cellular Comp Physiol* 21 77-91
- 935 Quastel J H and A H M Wheatley 1938 *Biochem J (London)* 3 936-945
- 936 Racker E 1957 *The Harvey Lectures Series* 51 143-174 New York Academic Press
- 937 Ramachandran K and T K Walker 1951 *Arch Biochem Biophys* 31 224-233
- 938 Ramachandran K and T K Walker 1952 *Arch Biochem Biophys* 3 195-203
- 939 Ramakrishnan C V 1954 *Enzymologia* 17 169-174
- 940 Ramakrishnan C V and S M Martin 1954 *Can J Biochem and Physiol* 3 431-439
- 941 Ramakrishnan C V and S M Martin 1955 *Arch Biochem Biophys* 55 403-407

- 212 Ramakrishnan C V R Steel and C P Lentz 1955 *Arch Biochem Biophys* 55 270-273
- 213 Raveux R 1950 *Compt rend (Paris)* 238 729-731 1006-1008 1107-1108
- 214 Reiner J M H Gest and M D Kamen 1949 *Arch Biochem* 20 175-177
- 215 Rippel A and H Bortels 1927 *Biochem Z* 184 237-241
- 216 Rippel A and F Heilmann 1930 *Arch Mikrobiol* 1 119-136
- 217 Rippel A and H Wiangke 1941 *Arch Mikrobiol* 10 124-127
- 218 Roberts R B et al 1955 *Carnegie Inst Wash Publ* 607 1-91
- 219 Robinson R 1926 *Trans Brit Mycol Soc* 10 307-314
- 220 Rockwell G E and J H Highberger 1927 *J Infectious Diseases* 40 438-446
- 221 Rothstein A and H Berke 1952 *Arch Biochem Biophys* 36 195-201
- 222 Sakaguchi K H Takahashi and H Morino 1953 *J Agr Chem Soc Japan* 27 591-593
- 223 Schade A L and K V Thimann 1940 *Am J Botany* 27 659-670
- 224 Schatz A G S Trelawney V Schatz and R R Mohan 1956 *Mycologia* 48 883-885
- 225 Schmidt Lorenz W 1956 *Arch Mikrobiol* 25 137-165
- 226 Seaman G R and M D Navchke 1955 *J Biol Chem* 217 1-19
- 227 Semeniuk G 1944 *Iowa State Coll J Sci* 18 325-358
- 228 Shepherd C J 1951 *Biochem J (London)* 48 483-486
- 229 Shibata K and H Tamiya 1930 *Acta Phytochim (Japan)* 5 23-97
- 230 Shu P 1953 *J Agr Food Chem* 1 1119-1123
- 231 Shu P A Funk and A C Neish 1954 *Can J Biochem and Physiol* 30 68-80
- 232 Siegel B V 1950 *Proc Natl Acad Sci U S A* 36 515-517
- 233 Sih C J P B Hamilton and S G Knight 1957 *J Bacteriol* 73 447-451
- 234 Sih C J and S G Knight 1956 *J Bacteriol* 72 694-699
- 235 Sisler H D and C E Cox 1954 *Am J Botany* 41 338-345
- 236 Smith L 1954 *Bacteriol Revs* 18 106-130
- 237 Solé A and G de la Fuente 1957 *Biochim et Biophys Acta* 24 206-207
- 238 Sowden J C S Frankel B H Moore and J E McClary 1954 *J Biol Chem* 206 547-552
- 239 Steinberg R A 1942 *Plant Physiol* 17 129-132
- 240 Stuckland H 1956 *Biochem J (London)* 64 498-503
- 241 Stokes J L 1950 *Bacteriol Proc (Soc Am Bacteriologists)* 1950 119
- 242 Stout H A and H Koffler 1951 *J Bacteriol* 60 253-268
- 243 Strauss B S 1952 *Arch Biochem Biophys* 36 33-47
- 244 Strauss B S 1953 *Arch Biochem Biophys* 44 200-210
- 245 Strauss B S 1956 *J Gen Microbiol* 14 494-511
- 246 Strauss B S 1957 *J Biol Chem* 225 535-544
- 247 Strauss B S and S Pierog 1954 *J Gen Microbiol* 10 221-235
- 248 Sumiki Y 1929 *Bull Agr Chem Soc Japan* 5 14-16
- 249 Sumiki Y 1931 *Bull Agr Chem Soc Japan* 7 62-63
- 250 Sussman A S and C L Markert 1953 *Arch Biochem Biophys* 45 31-40
- 251 Suthers A J and T K Walker 1932 *Biochem J (London)* 26 317-319
- 252 Swaby R J and B I Passey 1953 *Australian J Agr Research* 4 331-339
- 253 Swim H E and L O Krampitz 1954 *J Bacteriol* 67 419-425
- 254 Syrett P J 1953 *Ann Botany* 17 1-36
- 255 Takahashi T and T Asai 1933a *Zentr Bakteriell Parasitenk Abt II* 88 316-384
- 256 Takahashi T and T Asai 1933b *Zentr Bakteriell Parasitenk Abt II* 89 81-84
- 257 Takahashi T and T Asai 1933c *J Agr Chem Soc Japan* 9 443-448
(*Chem Abstr* 27 4273 1933)

- 288 Takahashi T K Sakaguchi and T Asai 1957 *Bull Agr Chem Soc Japan* 3 87-97
- 289 Tamiya H 1957 *Acta Phytotchim (Japan)* 3 51-173
- 290 Tamiya H 1958 *Acta Phytotchim (Japan)* 4 77-113
- 291 Tamiya H 1959 *Advances in Enzymol* 2 183-238
- 292 Tamiya H and Y Miwa 1959 *Z Botan* 21 417-437
- 293 Tatum E L and S R Gross 1956 *J Biol Chem* 219 797-807
- 294 Tauber H 1954 *Enzymologia* 16 311-316
- 295 Theorell H 1951a In J B Sumner and K Myrback (eds) *The Enzymes* Vol 2 (1) p 333-356 New York Academic Press
- 296 Theorell H 1951b In J B Sumner and K Myrback (eds) *The Enzymes* Vol 2 (1) p 97-127 New York Academic Press
- 297 Thumann K V 1951 *The Life of Bacteria* New York The Macmillan Co pp 175
- 298 Thunberg T 1950 *Skand Arch Physiol* 40 1-91
- 299 Tissieres A 1954 *Biochim J (London)* 58 147-146
- 300 Tissieres A and H K Mitchell 1954 *J Biol Chem* 208 941-949
- 301 Tissieres A H K Mitchell and F A Haskins 1953 *J Biol Chem* 205 423-433
- 302 Tomlinson T C 1957 *New Phytologist* 56 418-434
- 303 Tso P O P J Bonner L Eggman and J Vinograd 1956 *J Gen Physiol* 39 325-347
- 304 Tso P O P L Eggman and J Vinograd 1956a *J Gen Physiol* 39 801-812
- 305 Tso P O P L Eggman and J Vinograd 1956b *Arch Biochem Biophys* 66 64-70
- 306 Tytell A A and B S Gould 1941 *J Bacteriol* 47 513-516
- 307 Umbreit W W 1957 *J Cellular Comp Physiol* 41 Suppl. 1 39-66
- 308 Umbreit W W R H Burris and J F Stauffer 1949 *Manometric Techniques and Tissue Metabolism* 2nd ed Minneapolis Burgess Publishing Co pp 2-7
- 309 Utter M F and H G Wood 1951 *Advances in Enzymol* 12 41-151
- 310 Waksman S A and J W Foster 1938 *J Agr Research* 57 873-899
- 311 Walker T K A N Hall and J W Hopton 1951 *Nature (London)* 168 1042-1043
- 312 Warburg O 1938 *Ergeb Enzymforsch* 7 210-211
- 313 Warburg O and F Kubowitz 1959 *Biochem Z* 214 24-25
- 314 Ward G E L B Lockwood B Tabenkin and P A Wells 1938 *Ind Eng Chem* 30 1233-1235
- 315 Webley D M 1954 *J Gen Microbiol* 11 114-121
- 316 Webley D M and P C DeKock 1952 *Biochem J (London)* 51 371-375
- 317 Wehmer C 1907 In *Lafar's Handbuch der technischen Mykologie* Vol 4 p 506-518
- 318 Wenner C E and S Weinhouse 1956 *J Biol Chem* 219 691-704
- 319 Whitaker D R and P E George 1951 *Can J Botany* 29 16-181
- 320 Wikén T and H Sommar 1957 *Experientia* 8 140-147
- 321 Wilson T C C and P W Wilson 1955 *J Bacteriol* 70 30-31
- 322 Winkler R J 1940 *J Cellular Comp Physiol* 15 313-354
- 323 Wirth J C and F F Nord 1951 *Arch Biochem* 1 143-163
- 324 Wolf F T 1947 *Arch Biochem* 13 83-97
- 325 Wong D F O and S J Ayl 1956 *J Am Chem Soc* 78 330-331
- 326 Wood H G 1946 *Physiol Revs* 26 193-241
- 327 Wood H G 1955 *Physiol Revs* 35 841-859
- 328 Wood H G C R Brewer M N Mickelson and C H Werkman 1940 *Enzymologia* 8 314-317

- 329 Wood W A 1955 *Bacteriol Revs* 19 223-233
330 Wood W A and R F Schwerdt 1953 *J Biol Chem* 201 501-511
331 Yanofsky C and J L Reissig 1953 *J Biol Chem* 202 567-577
332 Yemm E W and B F Folkes 1951 *Biochem J (London)* 57 49-508
333 Yull J L 1928 *Biochem J (London)* 2^o 1504-1507

8. Nitrogen Nutrition and Metabolism

With the exception of certain problems in amino acid biosynthesis the nitrogen metabolism of the fungi has been relatively neglected. Numerous surveys of nitrogen nutrition have been made unfortunately too often without a clear conception of the physiological problems involved. As will be seen little is known of the mechanism of nitrogen assimilation into amino acids and proteins the synthesis of specific peptides and the metabolism of purines and pyrimidines.

There is no optimum amount of nitrogen for a culture the demand depends in the first instance upon the carbon supply but in principle at least any factor may change the apparent optimum concentration of the nitrogen source. The dependence upon carbon is illustrated in Figure 1 in this organism *Aspergillus niger* enough is known of other cultural factors so that a strict proportionality between carbohydrate supply and nitrogen demand can be demonstrated (496).

No single pattern of nitrogen assimilation can be described to apply to all fungi. In general however inorganic or organic nitrogen is taken up rapidly during the phase of growth. Even in this phase however there may be a back movement of ammonia nitrogen from the cells to the medium (211) reflecting the permeability of the cell to ammonia. Certain enzymes also begin to appear in the medium during this period (115-278) and in at least some organisms amino acids and other soluble nitrogen compounds are liberated (139-181-374-413). The major liberation of proteins and other nitrogenous




Figure 1 The relation between growth of *Aspergillus niger* and nitrogen supply (ammonium nitrate) in media containing sucrose at 50 gm per liter (curve 1) and 2.5 gm per liter (curve 2). Drawn from data of Steinberg and Bowling (496)

compounds occurs of course in the phase of autolysis (Chapter 1)

Studies of *Scopulariopsis brevicaulis* indicate that upon exhaustion of the utilizable nitrogen of the medium there is a mobilization and reutilization of at least part of the nitrogen from older hyphae to support development of new mycelium (374)

1 INORGANIC SOURCES OF NITROGEN

Primary attention must be devoted here to three inorganic forms of nitrogen: ammonium salts, nitrites, and nitrates. Other inorganic nitrogen compounds have been little studied but appear in general to be utilized only if easily or spontaneously converted to one of the types above named (278-491). Cyanide as a nutrient and metabolite is considered separately.

The Utilization of Nitrate Nitrogen Nitrates are excellent sources of nitrogen for many fungi (30, 221, 326, 459). Inability to utilize nitrate is especially common in some groups, e.g., the higher basidio mycetes (221, 234, 275, 327, 397, 546), the Saprolegniaceae (55, 434, 562), and the Blastocladales so far as known (90). Nitrate utilization or at least nitrate reduction is common in the actinomycetes although

one could ask for more quantitative data (15 306 419 534 569 570) but some strains definitely cannot grow with nitrate as the sole source of nitrogen (138 144 393 541)


It is unlikely in the extreme that inability to use nitrate can serve as a phylogenetic criterion. Within a genus individual species differ in nitrate utilization (150 327 397 610). The capacity to use nitrate can be lost by mutation (250 255 420 428 455) and survival in nature of strains carrying this mutation is presumably governed by their ecological situation rather than their taxonomic position.

Failure to utilize nitrate is usually presumed to be absolute but there has been little exploration of the possibility that nitrate can be metabolized once growth is initiated on another more readily available nitrogen source. When spores of *Streptomyces griseus* are inoculated into a synthetic nitrate medium no growth occurs but a pregrown mycelium uses nitrate readily (105).

As mentioned below nitrite may be toxic in an acid medium. It follows that the kind and the concentration of the carbon source insofar as these affect the pH will decisively influence the response of some organisms e.g. *Streptomyces coelicolor* (107) to nitrate.

Ammonium nitrate is often used in media and it is of interest to ask whether there is preferential utilization of either ion. The frequent experience that utilization of ammonium nitrate is accompanied by a pH drop (227 256 409 415 529) is presumptive evidence for preferential utilization of the ammonium ion. More direct evidence is provided however by analytical studies. In *Scopulariopsis brevicaulis* and other fungi nitrate utilization does not begin until virtually all the ammonium has disappeared from the medium (Figure 2) this is

Fig 2 The utilization of ammonium and nitrate by *Scopulariopsis brevicaulis* in a medium containing glucose sucrose and ammonium nitrate. Curve 1 nitrate concentration in the medium curve 2 ammonium concentration in the medium. From tabular data of Morton and MacMillan (375)



also true of *Neurospora crassa* (511). Detailed study of the phenomenon (315, 373, 375) brings out that nitrate utilization is completely stopped by the addition of ammonium salts provided the ammonia is assimilated i.e. that a carbon source is available. Since nitrite utilization is not blocked by ammonium it must be assumed that in these fungi ammonium prevents the reduction of nitrate to nitrite; this is borne out by enzymatic studies which show that the nitratase system declines in quantity soon after ammonia assimilation begins.


Analytical data on utilization of ammonium nitrate by *Helminthosporium gramineum* show a different relation: ammonium ion is used much more rapidly than nitrate but there is a slight assimilation of nitrate even in the early stages of growth (110). This relation is stated to prevail in some other fungi (375).

The preferential utilization of ammonia from ammonium nitrate is influenced by pH. Foster (163) reviews several studies in *Aspergillus niger* all of which agree that at very low pH ammonium utilization is reduced and at least some nitrate is assimilated. Attempts have been made to explain this phenomenon as one of permeability or as based on the adsorption capacity of cell colloids (86, 264); neither of these explanations is convincing and it would appear desirable to reinvestigate the phenomenon itself before speculating on its theoretical implications. As shown later (Figure 5) the optimum pH for nitrate uptake in *Scopulariopsis brevicaulis* is about 6.0, far above the critical pH at which *Aspergillus niger* is claimed to assimilate nitrate (439).

Nitrate reduction in the fungi is believed to be strictly assimilatory in nature—that is to occur only in the incorporation of nitrate nitrogen into the cell. This process is in contrast to that of the bacteria, many of which reduce nitrate without assimilation of the nitrogen; nitrate if it has an essential function is acting as a hydrogen receptor for substrate oxidations (556). This nitrate respiration appears to be mediated by the cytochrome system in *Micrococcus denitrificans* and other bacteria (460, 557) whereas as will be seen nitrate reduction in the fungi is mediated so far as we know by flavin enzymes. It should be obvious of course that utilization of nitrate in fungi does depend upon substrate oxidations and affects therefore the respiratory quotient (521).

The nitrate reductase of *Neurospora crassa* has been studied intensively; this work is reviewed by Nason (384). The enzyme is formed only on nitrate or nitrite media i.e. it is inducible. Flavin adenine dinucleotide is the prosthetic group; sulphhydryl groups are essential to activity and molybdenum is an essential factor in electron transport.

The passage of electrons is visualized



That is nitrate is reduced with concurrent oxidation of some other presumably carbon compound by a triphosphopyridine linked dehydrogenase. The enzyme of *Hansenula anomala* differs in that either di- or triphosphopyridine nucleotide is functional (483).

Two other flavin enzymes of *Neurospora crassa* nitrite reductase and hydroxylamine reductase also inducible metal dependent enzymes have not as yet been separated. Together and possibly with the aid of a third enzyme they catalyze the reduction of nitrite to ammonia (384 385 451).

From these enzymatic studies and from the somewhat uncertain occurrence of hydroxylamine in nitrate cultures (322 396 494) an inorganic pathway of nitrogen reduction has been proposed (384)



The unknown intermediate possibly hyponitrite is included on the assumption that the four electron change from nitrite to hydroxylamine is unlikely to be a single reaction.

Whether this inorganic pathway is the sole and obligatory sequence of nitrate reduction in fungi is not yet certain. Since molybdenum deficiency reduces both nitrate reductase content and the ability to grow on nitrate in *N. crassa* and *Aspergillus niger* it seems probable that the first proposed step is essential. Genetic studies bear out this conclusion (484). However we cannot be sure that succeeding steps are necessarily inorganic in nature as implied in Equation 2 incorporation into organic combination may occur at an oxidation level above that of ammonia (220 338 484).

A nitroaryl reductase system in *Neurospora crassa* catalyzes the reduction of *m* dinitrobenzene by reduced pyridine nucleotides to *m* nitroaniline (384). This system although possibly not specific is of interest as a model of the reduction of organic nitro compounds as an alternative to the inorganic pathway of Equation 2.

The increased toxicity of thiourea to *Aspergillus niger* in nitrate media suggests that the poison interferes with assimilation (158). Chlorate toxicity is also more evident in nitrate media (203). Fluoride causes the accumulation of nitrite in fungi presumably acting specifically on the nitrite reductase system (373).

The Utilization of Nitrite Nitrogen Nitrite serves more or less well as the sole source of nitrogen for a number of fungi e.g. *Fusarium niveum* (622) *Coprinus* spp (165) *Phymatotrichum omnivorum* (520) *Scopulariopsis brevicaulis* (375) and *Rhizophilicystis rosea* (424). Different species of *Aspergillus* use it well poorly or not at all (305 457 494). Failure to utilize nitrite which is the more common situation in fungi is explained on the basis of its known toxicity (233 402 693). The fact that growth in the presence of nitrite is best in alkaline media (107 402 454) indicates that it is the free unionized acid rather than the nitrite ion which is the toxic species. Nitrite toxicity to *Fusarium* *lim* is evidenced by accumulation of pyruvic acid (396). Induction by nitrite of morphological variants in *Aspergillus* spp also suggests toxicity (495 497 498).

The Utilization of Ammonium Nitrogen Studies on *Scopulariopsis brevicaulis* (336 338 375) provide the best general view of the physiology of ammonium utilization. Both oxygen and an exogenous carbon source are required i.e. assimilation depends on respiratory energy or glucose breakdown products or both. Ammonium utilization increases with pH and unlike nitrate absorption has no definite pH optimum (Figure 3). Ammonia enters and leaves the cell by passive diffusion of the undissociated NH_3 molecule; the requirement for respiration is not therefore an indication of accumulation against a gradient.

In organisms generally (123 558) in *Torulopsis utilis* in particular (620) it is believed that the major and primary reaction of ammonium assimilation is the formation of glutamic acid. Other




Figure 3 The rate of uptake of ammonia and nitrate by *Scopulariopsis brevicaulis* as a function of pH. Curve 1 ammonia (in buffers); curve 2 nitrate. From Morton and MacMillan (375) by permission of the Oxford University Press.




Fig. 4 The effect of nitrogen assimilation on pH in *Karlingia* (*Rhizophyctis*) *rosea*. Redrawn from Hallins and Weston (297) by permission of the *American Journal of Botany*.

amino acids can be formed from glutamate by transamination (p. 267). Secondary roles in ammonium assimilation may be filled by reactions which incorporate free ammonia into alanine, aspartic acid, or glutamine. Respiratory breakdown of glucose is obviously necessary to supply the carbon skeleton.

One of the best established physiological correlates of ammonium assimilation from such salts as the sulfate, nitrate, or chloride is the rapid and often quantitatively large drop in pH consequent upon preferential utilization of the cation. Typical data are shown in Figure 4. The generality of this reaction may be emphasized by listing a few of the fungi in which careful observation demonstrates the phenomenon beyond question.

Alternaria solani (391) *Aspergillus niger* (496) *Chaetomium funicola* (477) *Choanephora cucurbitarum* (43) *Chytridium* sp. (113) *Diplocarpon rosae* (482) *Entomophthora* spp. (604) *Myrothecium verrucaria* (122) *Mycosphaerella pinodes* (45) *Penicillium* spp. (119, 254) *Phymatotrichum omnivorum* (520) *Piricularia oryzae* (406) *Psalliota bispora* (546) *Pythium* spp. (459) *Polyporus* spp. (221) *Ferticillium* spp. (221, 256) *Nylaria mali* (221).

In many, if not most, of the instances cited, the decrease in pH was sufficient to reduce growth. Practically, several remedies exist; use

of ammonium carbonate or phosphate (177) increase in buffer concentration (122) use of mixed nitrogen sources (254) or inclusion of utilizable organic acids in the medium. This last phenomenon has given rise to much confusion in the literature the improvement of growth by use of organic acids in media relying on ammonium nitrogen being taken for a specific nutritional or metabolic effect of the acid. Morton and MacMillan (375) correctly and with adequate evidence dispose of this argument the organic acid effect is a pH effect. Illustrative evidence for the effect itself is abundant—work on the Mucorales (29) higher basidiomycetes (275) and miscellaneous fungi (221) may be considered typical.

This point has been treated at some length because a requirement for organic nitrogen i.e. inability to grow with ammonium nitrogen may reflect only failure of the organism to grow at low pH. *Allomyces javanicus* var *macrognus* (340) and *Chytridium* sp. (113) illustrate this phenomenon. No claim of failure to utilize ammonium nitrogen can stand unless the pH effect is excluded experimentally.

Other possibilities of error in claims for a non specific organic nitrogen requirement may be mentioned. The amino acids which appear to be essential may be acting to reduce by complex formation the inhibitory effect of heavy metals present in the basal medium. copper is especially likely to be removed by amino acids (16, 17) and histidine specifically antidotes cobalt toxicity (411, 462). Apparently essential sources of organic nitrogen may be effective by virtue of their content of vitamins (234, 591).

The apparent glutamic acid requirement of *Allomyces javanicus* var *macrognus* disappears when the concentrations of glucose, potassium and minor elements are changed and sulfate is eliminated from the medium (339).

It seems therefore that many reports of failure to utilize ammonium nitrogen (318, 343, 359, 506) must remain *sub judice*. There does appear however to be in some of the lower phycomycetes—*Blastocladiella emersonii* (42), *Sapromyces elongatus* (204) and probably *Leptomitius lacteus* (463)—a genuine inability to grow with ammonium as the sole nitrogen source but no specific amino acid requirement. Contradictory and only semi quantitative data on *Apo-dachlya brachynema* (200, 461) do not permit of a decision. Finally the relatively few naturally occurring fungi with specific amino acid requirements (p. 259) and the many known deficient mutants are of course exceptions to the rule that fungi as a group can at least under specified conditions grow with ammonium ion as the sole source of nitrogen.

Ammonia in high concentrations is toxic (332) and may even find limited use as a fungicide (218 388). Abnormalities in growth (458) spore germination (169) and development (108) have been traced to ammonia toxicity especially in media with large amounts of organic nitrogen ammonia may accumulate to an inhibitory level and contribute to the so called staling effect (421). The toxicity of high ammonium ion concentration to *Allomyces javanicus* var *macrognus* is alleviated by phosphate (340).

Nitrogen Fixation From the earliest days of microbiology claims have been made that fungi are able like certain bacteria to convert gaseous nitrogen to combined nitrogen. These studies have been reviewed several times in the last generation (18 137 163 202 478). The evidence is overwhelmingly against the occurrence of nitrogen fixation in saprophytic fungi. Results with mycorrhizal fungi particularly *Phoma* spp. have been more often positive but isotope results fail to corroborate earlier claims (601). There is therefore at present no good evidence for fixation of nitrogen by the fungi. Nitrogen fixation occurs in root nodules of *Alnus glutinosa* (426) but it is not yet known what microorganism is involved the causal agent has been variously assigned to the bacteria the actinomycetes the filamentous fungi and most recently the Plasmodiophorales (228). Until the organism can be cultivated both its identity and its role in nitrogen fixation will remain uncertain.

Oxidation of Inorganic Nitrogen Compounds Heterotrophic oxidation of inorganic nitrogen compounds is effected by a few bacteria (126) this is to be distinguished from the better known autotrophic oxidation of ammonium and nitrite. Among the true fungi *Aspergillus aureus* and *A. batatae* are reported to form nitrate from nitrite during growth (457). *Aspergillus flavus* growing on a glucose peptone yeast extract medium produces small amounts of nitrite and rather larger amounts of nitrate (468). This finding if confirmed has importance to soil microbiology. Color tests suggest that *A. niger* oxidizes ammonia to hydroxylamine (494).

Among the actinomycetes *Nocardia corallina* oxidizes pyruvic oxime and hydroxylamine to nitrite (277 321). An unusual species of *Streptomyces* *S. nitrificans* has been studied by Schatz and coworkers (257 258 464 465). This organism utilizes urethan as the sole source of both carbon and nitrogen and forms nitrite during growth on carbamates and during metabolism of urea or ammonium carbonate. Activity of cell free preparations suggests a pyridine nucleotide linked ammonium dehydrogenase. Nitrate is not formed.

Cyanide as a Metabolite Cyanide is of course an effective respiratory poison acting on iron enzymes. It has also however been reported to provide assimilable nitrogen to preformed mycelium of *Aspergillus niger* (271) and to cause an apparent increase in growth of *Fusarium lini* (395-437). Data so far available however are not adequate for a final conclusion on the utilizability of cyanide as a nitrogen source. It is to be noted that Chughtai and Walker (104) found that the growth promoting effect of cyanide is not duplicated by nitrate; they suggest a metabolic disturbance which diverts more carbon into dry matter. An unidentified soil organism, probably an aerobic actinomycete, makes visible growth with cyanide as the sole source of both carbon and nitrogen (578).

The liberation of hydrogen cyanide from the fruit bodies of basidiomycetes has been noted repeatedly; reports are reviewed by Bach (38), Muller (379), Robbins et al. (444) and Zellner (629). An unidentified *Mucor* sp. was reported in 1916 to form both cyanide and benzaldehyde (219). Cyanide is produced in culture but apparently only during autolysis, not during growth (444). Fruit bodies of *Pholiota aurea* do not contain free cyanide when freshly collected, and circumstantial evidence indicates that it is released from another compound by an enzymatic reaction (37).

Production of cyanide by an unidentified basidiomycete growing under snow cover is apparently responsible for crown rot in alfalfa; the fungus in culture produces cyanide at low temperature (319-320).

2. PROTEINS AND PEPTIDES

Nutritional Aspects The fungi and actinomycetes as a group are active in the decomposition of proteins in soil and other materials (1568-571, 572-573). In pure culture it appears that proteins such as gelatin, casein, and egg albumin can serve as sources of nitrogen for at least the common saprophytic fungi and actinomycetes (278-574). Much more data are available on enzymatic destruction of common proteins than on their use for growth. Utilization of keratin is common only among dermatophytic fungi and possibly chytrids (Chapter 3). Whether fungi can attack highly purified proteins in the absence of any other nitrogen source is in doubt.

Commercial peptone—a mixture of peptides of varying chain length—is generally utilizable but in recent years has been replaced in culture media meant for physiological studies by materials of known composition. Dipeptides and tripeptides support the growth of *Aspergillus* spp. (2-531) and *Ustilago* spp. (62-470); the histidine

requirement of *Trichophyton rosaceum* is met by carnosine and histidylhistidine (136). The widespread occurrence of enzymes hydrolyzing peptides to amino acids indicates that in general at least the simple linear peptides will be found to be adequate sources of nitrogen for most fungi. *Tricholoma gambosum* is unusual in that growth is tripled by glycylglycine unaffected by glycine (398).

The requirement of some Myxomycetes for protein has been mentioned elsewhere (Chapter 10).

Enzymatic Breakdown of Proteins and Peptides Following current usage (487) we distinguish two general types of enzyme which attack peptide bonds.

1 Endopeptidases (proteinases) acting on peptide bonds either in simple peptides or in the interior of a protein chain.

2 Exopeptidases acting only on peptide bonds which are adjacent to a free α amino or a free carboxyl group and therefore restricted generally to hydrolysis of small peptides.

Both these types are subsumed under the general term protease.

Surveys of the activity of culture filtrates indicate that an enzyme or enzymes able to hydrolyze gelatin is extremely common among the Fungi Imperfecti (129, 130, 142, 351) and *Streptomyces* spp. (130, 276). Generalization as to other fungi is difficult; gelatin hydrolysis is however effected by some ascomycetes (130) and basidiomycetes (53, 629). Among the Mucorales *Rhizopus* sp. grows upon and hydrolyzes several proteins (571) and *Mortierella renispora* forms active proteases (587, 588).

However, surveys limited to studies of the culture filtrate cannot be taken as final evidence of lack of a protease. The distribution of proteolytic enzymes between mycelium and culture fluid is conditioned both by age (185) and by the medium (208). This may explain some apparent disagreements (130, 142).

The most extensive study of the utilization of proteins by fungi and actinomycetes is that of Waksman and Starkey (571). *Streptomyces violaceus ruber* and *Rhizopus* sp. grow upon and decompose casein, zein, gliadin, fibrin, gelatin, and egg albumin. *Trichoderma koningi* utilizes edestin, zein, gliadin, and casein. In culture, proteins are used more rapidly if there is no other carbon source, but the liberation of ammonia under these conditions is large.

Other less comprehensive studies show that fungi and actinomycetes generally hydrolyze fibrin (19, 185, 169, 567) and casein (262, 333, 485, 489, 524, 567). Keratin hydrolyzing preparations have been ob-

tained from *Streptomyces* sp (399), and it may be assumed that the destruction of keratin by dermatophytic fungi is carried out by an extracellular enzyme (491 492 555). A protease of *Penicillium* sp converts trypsinogen to trypsin but is distinct from the enterokinase which carries out this reaction in animal tissues (312).

Fungal endopeptidases cannot be identified with known exocellular animal enzymes. However it is reasonable to believe that studies of purified enzymes acting on synthetic substrates will eventually support a rational classification. For the present a few general observations must suffice.

1 Crude and purified preparations from a fungus usually have their pH optimum somewhat on the alkaline side of neutrality (142 185 283 335 485) but particular activities may have an optimum as low as 4.5 (115).

2 Substrate and inhibition studies suggest that endopeptidases of the trypsin type occur in *Streptomyces* sp (377) and in *Trichophyton gypseum* (185) and other fungi (335). A purified enzyme from *Streptomyces proteolyticus* acts on pepsin substrates but at a pH very different from that of the animal enzyme (549).

3 Crude and highly purified preparations alike consist of mixtures of different enzymes (116 279 335 388 589 621), enzymes of even closely related strains show differences in rate of attack on different proteins (142).

Endopeptidases of *Aspergillus oryzae* have been studied thoroughly (115 116 198). Three different enzymes acting on proteins have been separated. The major component probably acts on most proteins including those which lack aromatic acids and has been obtained in a crystalline although not completely pure state (114 198).

Culture filtrates of *Streptomyces* spp lyse bacterial cells. Lysis of heat killed Gram negative bacteria has been attributed to a protease (67 285 376, 519). Lysis of Gram positive bacterial cells by different species of *Streptomyces* is effected however by an enzyme similar to lysozyme i.e. a mucopolysaccharase (333 334) or by the combined action of a ribonuclease and a protease (376 378).

The formation of exocellular protein hydrolyzing enzymes by some fungi and actinomycetes proceeds in synthetic media with inorganic nitrogen sources (131 185 352) for other species a more complex medium may be required (131). There is as yet no good evidence of enzyme induction by substrate although use of protein media may increase yields (53 530). A study of cultural conditions as they affect

endopeptidase formation by *Aspergillus oryzae* shows as expected optima for carbon source salts iron and temperature (352)

Culture filtrates or mycelial extracts of fungi commonly perhaps universally hydrolyze peptides. However it is hazardous at present to identify this activity with particular exopeptidases known from other sources or even to state categorically that endopeptidase activity has been excluded in many of the reports. *Aspergillus parasiticus* forms four exopeptidases including an aminopeptidase which is activated by zinc and by reducing agents (284) these same activators affect the aminopeptidase of *Penicillium* spp (52). A number of fungi—*Aspergillus* spp *Penicillium* spp *Cunninghamella* sp *Coprinus radians* and *Agaricus campestris*—form a dipeptidase an aminopeptidase and a carboxypeptidase many of these fungi also elaborate an aminotripeptidase assayed with triglycine and a glycylglycine dipeptidase (53). A similar range of activities is reported for *Mortierella renispora* *Chaetomium* sp and other fungi (336).

Glycylglycine dipeptidase of *Aspergillus oryzae* is activated by cobalt (115) as is the same enzyme from other sources (487). Dipeptidases also are produced by *Physarum polycephalum* (244) *Piricularia oryzae* (259) *Penicillium notatum* (263) and other fungi (336 403).

Hydrolysis of some amides may be effected by enzymes which also act on peptides or proteins. hippuricase for example may be identical with a carboxypeptidase (182).


It appears in summary that fungi produce a battery of enzymes attacking peptides of low molecular weight. Information is at present too fragmentary for a detailed classification of the fungal exopeptidases the availability of synthetic test substrates (487) makes the development of such a classification a possible enterprise.

Peptides of Fungi and Actinomycetes Following Bricas and Fromageot (76) we may define peptides as compounds of low molecular weight which contain two or more amino acids and peptide or peptidoid bonds. This excludes certain compounds such as pantothenic acid and biocytin (Chapter 10) in which one amino acid and a non amino acid moiety are joined by a peptide linkage. The naturally occurring peptides may be classified as homeomeric—consisting only of amino acids—and heteromeric—consisting of amino acids and a non amino acid moiety.

Glutathione perhaps deserves first place in this discussion because of its probable ubiquity in living cells its vital role in triose phosphate dehydrogenase and its probable role in peptide synthesis (76). Using

an iodometric method of possibly doubtful specificity Miller and Stone (366) report the presence of glutathione in mycelium of *Neurospora sitophila* and other fungi. The common occurrence in fungi of triose phosphate dehydrogenase (Chapter 7) of which glutathione is the prosthetic group (308) indicates that the peptide is at least wide spread among fungi and actinomycetes. A mutant of *Glomerella* sp requires glutathione and nutritional evidence suggests that it is deficient in the ability to form the component dipeptide γ glutamylcysteine (348).

The actinomycins form a group of related antibiotics first discovered by Waksman and Woodruff (575) and produced by *Streptomyces* spp (79) and by *Micromonospora* sp (157). Chemically they are heteromeric peptides all contain the same chromophore or despeptido actinomycin of the structure (81)




The peptide chain distinguishes the various actinomycins very many of which have been described (78-79) the identity of the individual actinomycins is much in doubt since it appears that many of the isolated fractions are in fact different mixtures of the same components (452). A given strain may produce one type or mixture early and other later in the culture cycle (210).

Several other antibiotics of *Streptomyces* spp are or contain peptide chains examples include viomycin streptothricin and streptoligin (76) and the less completely characterized amphomycin (230) and levomycin (95). Polypeptide antibiotics are very common also among the true bacteria. Such compounds are not usual among the filamentous fungi although penicillin (Chapter 6) may be considered as a heteromeric cyclic peptide and the toxin of *Helminthosporium victoriae* is probably a polypeptide (421a).

Lycomarasmin reviewed by Gaumann (190) is a peptide produced by *Fusarium oxysporum* f. *lycopersici* although it is able to cause wilting of plants its role in natural pathogenesis in the wilt diseases is in dispute. It appears in quantity in the medium fairly late in the culture cycle after autolysis has begun (Figure 5). A heteromeric peptide its structure has been determined (609) to be N (α [α hydroxy propionic acid]) glycyl asparagine.

Figure 5 Growth and lycomarasmin formation (in the culture medium) in *Fusarium oxysporum* f. *lycopersici*. Redrawn from Dimon¹ and Waggoner (127) by permission of Phytopathology.



Lycomarasmin forms a complex with iron and it is the complex which is responsible for all or most of its toxicity to higher plants (190 564). Both free lycomarasmin and the iron complex are relatively unstable in solution (128).

Two heteromeric peptides ennatiin A and ennatiin B are also produced by *Fusarium* spp. (190 417 418) and like lycomarasmin cause wilting of plants (193).

Aspergillus fumigatus and *Streptomyces* spp. release into the culture medium a diketopiperazine, the anhydride of L-leucyl-L-proline, previously known only from animal sources (280).

Lastly, the toxins of *Amanita* spp. may be mentioned; they have been reviewed most recently by Block, Stephens and Murrill (61). Originally the hemolytic nitrogenous glucoside phallin was separated from amanitin (8 466); later it was discovered that the second material is in fact three closely related peptides now known as α -amanitin, β -amanitin, and phalloidin. The European form of *Amanita phalloides* contains all three of these compounds amounting in total to about 0.005 per cent of its fresh weight (596). Other species of *Amanita* have only one or two of the peptides and most species have none (61). A complete structure has been proposed only for phalloidin (597) although the three toxins have been separated and purified (60 594 596). The suggested structure that of a bicyclic hexapeptide is unusual in that two amino acids are joined by a sulfur bridge.

The excretion of unidentified nitrogenous compounds, primarily peptides, into the medium during growth has been studied by Morton and Broadbent (374). This excretion occurs in *Scopulariopsis brevicaulis* during the period of rapid growth, i.e. it is not an autolytic phe-

nomenon in the usual sense, about 25 per cent of the nitrogen supplied appears in the medium and it is not reutilized

The enzymatic basis of peptide synthesis in fungi is at present unknown. Extracts of *Neurospora crassa* mycelium show γ -glutamyl transferase activity conceivably this system could function in the synthesis of peptides of glutamic acid (563). In *Bacillus subtilis* glutamyl di- and tripeptides are synthesized by transfer reactions of this general type (600).

3 UTILIZATION OF AMINO ACIDS AND AMIDES

Amino Acids and Amides in Nutrition The specificity of response to different amino acids indicates that in fungi as in other organisms amino acids are assimilated primarily as such and that conversion to ammonia before assimilation of the nitrogen is at best a secondary pathway (44-542). It can be shown by analysis that amino acids are assimilated at varying rates from a complex medium and that they exist in part as mentioned earlier (Chapter 2) as free acids in the mycelium. Typically the amino acid content of a medium drops as the acids are taken up then rises again as they are liberated during autolysis (181-413-423).

The value of individual amino acids for fungi is customarily asayed by growth on media containing the amino acid of interest as the sole nitrogen source. Studies of this type are numerous but generalization from them is difficult. We find that a given amino acid allows very good growth of one organism and only very little growth of another but we are not able to say whether this reflects permeability enzymatic capacities or merely such secondary problems as acidity changes consequent upon utilization. In addition amino acids and amides from natural sources may be contaminated with vitamins and there may also be metabolic consequences of the chelation of inorganic ions by amino acids.

Most investigations more or less agree that glycine, asparagine, glutamic acid and aspartic acid are the most likely to support good growth. However it is relatively easy to find fungi which grow only poorly with asparagine (164-252-440-625), aspartate (110-278) or glutamate (275-278). Conversely leucine is generally not an adequate source of nitrogen (409) but for *Trichophyton persicolor* it is one of the most easily utilizable of the amino acids (506). Tryptophan usually not well utilized supports normal growth of *Streptomyces coelicolor* (106). Examples of this type could be multiplied to establish the point that although generalizations may be made which fit most




Figure 6 The growth of *Venturia inaequalis* in media with amino acids as the source of nitrogen. Each value is the mean of 97 observations (3 concentrations, 3 incubation periods and 3 replicates). Growth on casein hydrolyzate has been given a relative value of 100. Drawn from tabular data of Pelleuer and Keitt (409).

fungi, they will not necessarily apply to any one fungus which happens to be of interest.

The data of Figure 6 show the range of utilizability of the common amino acids for *Venturia inaequalis*. Statistical analysis of these and other data shows that the apparent relative value of different amino acids is affected by the period of incubation, the nature of the basal medium, and the concentrations of amino acids used (409). Figure 6 also brings out the point that growth is usually better with a natural

or artificial mixture of different amino acids than with any single amino acid (442)

Asparagine deserves special attention since it is so often one of the best sources of nitrogen for fungi and actinomycetes. The question is does the amide have some particular value or is asparagine merely a source of aspartic acid and extra nitrogen? Asparagine and aspartate are about equivalent for many fungi (25-46, 275-604) but the amide is superior to the acid for *Tricholoma imbricatum* (398) *Piricularia oryzae* (406) and *Leptographium* sp. (317) and at least is sole nitrogen source for *Phycomyces blakesleeana* (323). No decision can be reached at this time nor can it be until possible pH effects of utilization of the two compounds are excluded: the role of pH is emphasized by the response of *Coprinus hirsutus* to asparagine and aspartate (165). It is significant in this connection that aspartic acid does not replace the asparagine requirement of a mutant of *Neurospora crassa* (523). It is also known that asparagine is in some systems more active as an amino group donor to transamination reactions than is aspartate (354-355). *Piricularia oryzae* utilizes the amide nitrogen of asparagine more rapidly than the amino nitrogen (260).

Glutamine has been little studied in fungi: it is superior to glutamic acid as a nitrogen source for *Tricholoma gambosum* (398) and is generally utilized by other basidiomycetes (275). Like asparagine it is active in transamination (354-355) and it may also have a role in transpeptidation. One specific and important function of glutamine has been pointed out (Chapter 5): the provision by transamidation of the amino group of glucosamine.

As mentioned later (Chapter 12) spore germination of *Neurospora crassa* is inhibited by amino acids: e.g. glycine at 50 μ g per milliliter (453). Hydroxyproline at about this level is markedly inhibitory to *Trichophyton* spp. and other dermatophytic fungi (443-471). Hydroxyproline is not toxic to other fungi: its action on *Trichophyton mentagrophytes* is partially reversed by L-proline (443). Methionine and phenylalanine are also inhibitory to *Trichophyton* spp. (506) methionine to other fungi (169). The possibility that toxicity is exerted at the cell surface deserves exploration in view of studies on histidine mentioned below.

The aliphatic amides although possibly acted upon by fungal enzymes (p. 265) are not generally utilizable: acetamide may be an exception (39-63, 120).

Utilization of the D-isomers of the amino acids is specific: thus *Apodachlya brachynema* utilizes D-leucine but *Leptomitia lacteus* does not (463) and *Ustilago scabiosae* can grow with D-valine, D-isoleucine

or D-alanine but not with D-leucine or D-phenylalanine as nitrogen source (62)

It seems likely that utilization of the D-amino acids depends upon oxidative deamination by D-amino acid oxidase (p. 260) ammonia liberated by the oxidase system is then presumed to be incorporated into L-acids (245). Specific racemases effecting direct interconversion of the D and L forms are known in bacteria (22, 28, 286, 383, 608) not as yet in fungi. Presumably a second factor in utilization of the D-amino acids is their occasional toxicity (149).

Specific amino acid requirements appear with reasonable frequency as induced mutations in fungi. They are however uncommon in nature and when found are more apt to be conditioned or partial deficiencies than absolute requirements. Thus *Mycena rubromarginata* responds in growth to tyrosine or phenylalanine but grows without either (169) and *Cenococcum graniforme* has a partial requirement for histidine (358). *Eremothecium ashbyi* requires several amino acids only in acid media (619). An absolute requirement for histidine has been reported in *Trichophyton* spp. (135, 136, 196) as has a requirement for arginine in *T. tonsurans* (515); these reports are based on responses in agar media. *Labrynthula minuta* var. *atlantica* requires leucine (559). *Lentinus omphalodes* tryptophan (171).

Requirements for methionine or cysteine are assumed to be related to sulfur rather than nitrogen metabolism and are considered in Chapter 9. Cysteine may also promote the growth of an organism requiring a low oxidation-reduction potential (275).

Transport of Amino Acids into the Cell In the present context the main purpose of this section must be to call attention to a problem which has been only tentatively explored in the fungi and for the solution of which promising experimental techniques are available. We know that a free amino acid pool exists in fungal cells that transport across the membrane occurs in both directions and that proteins and peptides are synthesized. In bacteria, yeast and animal cells there is good evidence that some or all amino acids enter the cell by an active transport mechanism requiring an expenditure of metabolic energy, but different organisms cannot at present be fitted into any single pattern (100, 183, 188). Lysine accumulation for example appears to be independent of an energy source in Gram positive bacteria but glutamic acid enters only if glucose is being respired. Histidine enters the cells of *Neurospora crassa* against a concentration gradient; the uptake is inhibited by arginine and methionine which are consequently toxic to a histidine-requiring mutant (350). An analogous


interaction accounts for the toxicity to *Lactobacillus arabinosus* of D-alanine (293) Threonine uptake is possibly inhibited in *N. crassa* by other amino acids but this is not yet certain (566)

Catabolism of Amino Acids The breakdown of amino acids and amides may be reviewed under two general headings the processes which lead to removal of nitrogen and the processes which alter the carbon skeleton These two categories of biochemical change are reviewed by Cohen (109) and Greenberg (214) respectively We must restrict this discussion largely to the fungi and only a very incomplete picture of amino acid metabolism will emerge inasmuch as many important reactions known in other organisms have not yet been investigated in the fungi


One approach to amino acid catabolism is to determine what compounds are either oxidized or deaminated by resting cells using manometric data or ammonia determinations as criteria *Penicillium chrysogenum* for example oxidizes glutamic acid proline and alanine rapidly other amino acids more slowly or not at all as judged by oxygen uptake (603) Washed cells of *Streptomyces venetiae* liberate ammonia from glutamate arginine proline and other amino acids not from tyrosine, tryptophan leucine or cysteine only those amino acids which are attacked can serve as sole sources of nitrogen for growth (212) *Leptomitia lacteus* oxidizes and deaminates L alanine glycine and L leucine (463)

A second avenue of investigation of amino acid catabolism is the study of particular enzymatic activities in cell free preparations or as reflected in activities of intact cells Most efforts of this type have been focused on amino acid oxidase (dehydrogenase) systems scattered information on other reactions is available but difficult to systematize

The D and L amino acid oxidases are flavoproteins known and studied in animal systems their general properties are reviewed by Cohen (109) and Meister (355) The usual manometric system employing crude extracts which contain catalase may be formulated as follows



In the absence of catalase the keto acid is decarboxylated and the over all reaction then proceeds



The D-amino acid oxidase of *Neurospora crassa* acts at varying rates on a number of amino acids (Figure 7) detailed comparison with the




Figure 7 The oxidation of some amino acids by the D-amino acid oxidase system of *Neurospora crassa*. Curves 1 to 6 represent in sequence the rate of oxidation of methionine, leucine, isoleucine, valine, lysine, and ornithine. The broken horizontal line is drawn at the theoretical value for the complete oxidation of the D-isomer. Redrawn from Horowitz (245) by permission of the *Journal of Biological Chemistry*.

corresponding enzyme of sheep kidney reveals that the rates of attack on amino acids and the response to inhibitors of the two systems are somewhat different (47). The enzyme exhibits maximal activity at pH 8.0–8.5 (245). The same or a very similar enzyme is found in mycelial extracts of *Aspergillus niger* and three *Penicillium* spp. not in *P. sanguineum* (143). The different fungal preparations differ somewhat in their substrate preferences (47, 143). *Penicillium chrysogenum* forms much more of the enzyme in a synthetic than in a complex medium (143).


An oxidase system acting on L-amino acids has been shown in *Neurospora* spp. (47, 87, 536) both in cells and in the medium (48). The enzyme has a somewhat broader substrate specificity than the L-amino acid oxidase of cobra venom (47). Indirect evidence supports the assumption that the prosthetic group is flavin adenine dinucleotide (87). Enzyme yield is increased by cultivation with casein hydrolyzate but this presumed induction occurs only in a low biotin medium (536). Endogenous amino acids interfere with detection of the enzyme (536).

The L-amino acid oxidase of *Neurospora crassa* converts lysine to its

Catabolism of the carbon chain of amino acids by fungi can be considered at present only on the basis of fragments of information many of which however are suggestive of breakdown pathways known in other organisms

Tyrosine in at least some fungi and actinomycetes is attacked by tyrosinase a polyphenol oxidase (Chapter 6) The sequence of reactions in *Neurospora crassa* (248) appears to be at least in general the same as in mammalian systems (324) tyrosine is oxidized via 2,4 dihydroxyphenylalanine (dopa) through a red hallochrome pigment to an insoluble black pigment a melanin The pigment of *Streptomyces scabies* probably—although evidence is incomplete—is formed by this same pathway (15 183 242 243) and it is usually assumed that black insoluble pigments as a group are products of tyrosinase action

In mammalian liver tyrosine is oxidized to acetoacetic acid and fumaric acid by the following sequence of reactions (214)




The last reaction—actually three reactions (300)—is not known in fungi Homogentisic acid can be isolated as a product of the tyrosine metabolism of *Aspergillus niger* and *Penicillium* spp (261 298 554) inhibition data implicate *p*-hydroxyphenylpyruvate in the tyrosine metabolism of *Neurospora crassa* (125)

The isolation of tyramine from *Aspergillus niger* cultures (552) suggests another type of tyrosine breakdown by decarboxylation as in bacterial systems

Tryptophan catabolism in fungi has been studied principally in relation to nicotinic acid biogenesis (Chapter 10) Other pathways are known in bacteria (14 121 214 493) involving indole catechol kynurenic acid or 5-hydroxytryptophan A preparation from *Neurospora crassa* converts 3-hydroxykynurenine an intermediate in nicotinic acid biosynthesis to xanthurenic acid (353)

Indoleacetic acid (auxin) a growth regulator for higher plants is produced by several fungi including *Aspergillus niger* *Phycomyces nitens* and *Rhizopus stolonatus* and by soil actinomycetes and bacteria (72 73 236 302 394 446 448) Its obvious structural relation to tryptophan and the effect of the amino acid in stimulating its formation by fungi indicate that the auxin is formed from tryptophan (537 605) Production by *Rhizopus stolonatus* is increased by aeration (538) Detailed enzymatic studies of the synthesis have not appeared but it

has been suggested that oxidative deamination decarboxylation and oxidation reactions occur as follows (353 584)



Although from time to time it is reported that indoleacetic acid increases the growth of fungi most studies have not shown this (367 438 584) or show it only under special circumstances (235). However acceleration of the maturation of the mesosporangium of *Allomyces arbuscula* on agar media perhaps by an effect on the sporangium wall indubitably occurs (341). Growth inhibition by indoleacetate is common (71 235 438 449) in *Nectria galligena* inhibition is reduced by biotin (400 401). It has been argued on the basis of inhibition studies alone that indoleacetate is an essential metabolite of *Diplodia natalensis* (195) the evidence however is subject to other interpretations.

Destruction of indoleacetic acid occurs in the cultures in which it is produced (584) and cell free enzyme systems of several basidiomycetes inactivate the auxin (77 148 479 545). The enzyme of *Omphalia flavida* is highly specific oxidizing indoleacetate with equimolar amounts of oxygen at pH 3.5 the first product of the reaction is not known (430 431).

Threonine is cleaved by *Neurospora crassa* to glycine and acetaldehyde (565) this non-oxidative reaction occurs also in animal systems (214).

The breakdown of histidine in bacterial and animal preparations proceeds via urocanic and glutamic acids (339 516) this system has not been studied in fungi although glutamate does accumulate during histidine breakdown by *Streptomyces venezuelae* (212). Histidine decarboxylase could not be detected in *Claviceps purpurea* or *Aspergillus* spp (585).

The oxidation of cysteine sulfur is mentioned in Chapter 9 and the role of methionine as a methyl donor in Chapter 6. No other significant information on the breakdown of the carbon chains of these compounds by fungi is available. Cysteine desulhydrase is not detectable in *Penicillium chrysogenum* its occurrence in bacteria is reviewed by Fromageot (180).

Arginine is broken down in the urea cycle discussed later. *Streptomyces griseus* extracts contain an inducible enzyme which converts arginine to guanidinobutyramide (539 540).

Glutamic acid is probably the principal donor of amino groups in transaminations (p 267) in the process it is converted to α -ketoglutaric acid. Hence its carbon chain may be assumed to be oxidized by

the tricarboxylic acid cycle Glutamic decarboxylase presumably yielding γ aminobutyric acid has been detected in *Penicillium chrysogenum* (201) and *Endomycopsis vernalis* (290)

It might be at first glance surprising that proline a ring compound is generally an excellent source of nitrogen for fungi (275 278 409 412) The answer conceivably lies in a ready conversion of proline to glutamic acid i.e. reversal of the steps in the synthesis of proline from glutamate (p 268)

Hydrolysis of Acyl Amino Acids and Amides Nitrogen acylated amino acids are hydrolyzed liberating the free amino acid by preparations from *Aspergillus* spp and *Penicillium* spp (263 364 389 405) Although these reactions may be referred to as acylase activity it must be realized that there is in fungi no clear separation of acylases amidases and peptidases all of which act on the $\text{—C—O—N}<$ bond

Hippuricase (histozyme) splits hippuric acid (N benzoylglycine) to benzoic acid and glycine Its identity and relation to other acylases and to peptidases is still obscure certain other N benzoylated L α -amino acids are attacked (325) It has most recently been studied in *Penicillium* spp (263 456) reports of its occurrence in other fungi are reviewed by Leuthardt (325)

Asparaginase hydrolyzing asparagine to aspartic acid and ammonia has received some attention although it has not been purified Activity is found in *Aspergillus niger* (467) *Penicillium* spp (134 456 535) *Microsporum* spp (51) and *Piricularia oryzae* (260) The optimum pH is about 8 (32 51 467) The enzyme of *A. niger* is liberated into the medium only on autolysis (33) is not easily solubilized in sand ground preparations (35) is unstable (36 630) and is formed in cells grown on a synthetic medium (467)

Amidase activity against glutamine is found in yeast bacteria and higher animals (630) it is probably common in the fungi but has been reported so far only in *Neurospora crassa* (523)

Several early reports reviewed by Foster (163) suggest that fungal enzymes also act on aliphatic or aromatic amides e.g. acetamide and benzamide In view of the negative results of Schmalfuss and Mothes (467) it is perhaps best to suspend judgment although aliphatic amides are hydrolyzed by *Torula utilis* (630) Penicillinase and penicillin amidase both acting on amide linkages have been mentioned briefly in Chapter 6

4 BIOSYNTHESIS OF AMINO ACIDS


The formation of amino acids by fungi may be considered for our purposes from three aspects (1) the entrance of nitrogen into the

amino acid pool (2) transfer of the nitrogen from one amino acid to another and (3) biosynthesis of the carbon chains of known amino acids either before or after incorporation of nitrogen

Most attention has been directed toward the synthesis of the protein amino acids. Even a superficial survey reveals that many other amino acids are formed by fungi either free or in combined form. Gamma aminobutyric acid probably arising from glutamate by decarboxylation is rather common (374 380 486 612). Djenkolic acid is found free in extracts and culture fluids of *Tilletia caries* (380). Beta lysine occurs in several actinomycete antibiotics (56 76 382). Antibiotics of fungi and actinomycetes yield on hydrolysis numerous N methyl amino acids for example sarcosine N methylvaline and N methylisoleucine (78 417). Alpha hydroxy amino acids occur combined in lycomarasin (609) and in ergot compounds (76) while D α methylserine is a constituent of the antibiotic amucetin (160).


Entrance of Nitrogen into the Amino Acid Pool We have observed that most fungi utilize ammonium nitrogen as the major or sole nitrogen source other inorganic forms are we may assume with certain reservations converted to ammonia before assimilation. The best established reaction is the amination of α ketoglutaric acid derived from carbohydrate via the tricarboxylic acid cycle to form glutamic acid i.e. by the reversal of Equation 5. As mentioned the only fungus in which this enzyme has been unequivocally demonstrated is *Neurospora crassa*. Distribution of carbon 14 in glutamate is consistent with the operation of this reaction in *Streptomyces griseus* (89 199).

Reversal of aspartase action occurs in bacteria (145 183) and the enzyme is known in fungi (p 262). It is not yet however known whether the reverse reaction




makes a significant contribution to amino acid synthesis in fungi. This leaves us for the moment with only one mechanism for the entrance of nitrogen into amino acids. On grounds of comparative biochemistry however it seems likely that future work will uncover other mechanisms especially in the synthesis of amides and of tryptophan histidine serine and other amino acids. *Blastocladiella emersonii* which will not grow with ammonia as sole nitrogen source forms glutamate from labeled glucose but isotopic equilibrium with α -ketoglutarate is not observed. This suggests that glutamic acid arises in some other way (91).

Transamination In organisms generally the amino group of glutamic acid can be transferred to various keto acids yielding α -ketoglutaric acid and a new α -amino acid thus



Investigation of these enzymes in filamentous fungi has only begun with the identification in *Neurospora crassa* of the transaminase reactions in which aspartate and alanine participate as the $\text{R}-\text{CH}(\text{NH})\text{COOH}$ of Equation 9 (151). Mutants which lack glutamic dehydrogenase exhibit a requirement for any of several amino acids; the fact that so many different amino acids support growth argues for an extensive transaminase system in *N. crassa* (151-152, 154).

Aspergillus fumigatus homogenates carry out the reaction (445)



This represents the transfer of an ω -amino group and is similar to the formation of glutamate from ornithine and α -ketoglutarate by extracts of *Neurospora crassa* (153-155). *Endomycopsis ternalis* also forms glutamic acid by the reaction of Equation 10 (290). As in other organisms transaminations in *N. crassa* are activated by pyridoxal phosphate (151).

A kynurenine transaminase catalyzing the transfer of the α -amino group of kynurenine or 3-hydroxykynurenine to keto acids has been partially purified from extracts of *Neurospora crassa*; again pyridoxal phosphate is required (274).


Asparagine and glutamine participate also in transaminations; the amino nitrogen is transferred to any of several keto acids (355). This reaction has not been explored in the fungi.

Biosynthesis of Amino Acid Carbon Skeletons Perhaps in no field of biochemistry have the techniques of biochemical genetics been more fruitfully employed than in that of amino acid synthesis. *Neurospora crassa* has been especially useful. In recent years the application of isotope and enzymatic methods has provided additional valuable in-

formation. It is not always realized that there are serious pitfalls in all methods which seek to state unequivocally that this or that compound is an obligatory precursor of a given metabolite. Adelberg (11) has critically reviewed these problems with the conclusion that present day techniques are unable to establish metabolic intermediates with finality. In practice one approaches certainty or conviction as more and more evidence from different types of experiment is accumulated (123).


It seems appropriate to include here only a sketch of the more important biosynthetic pathways which are known or are likely to be functional in one or more fungi. As will be seen there is little evidence from fungi other than *Neurospora crassa*. Following Davis (123) we shall consider groups of amino acids which are known to be biosynthetically related.

A family of amino acids is derived more or less directly from glutamic acid. Omitting details the following over all formulation may be suggested (123, 500)



Evidence from fungi is however limited at present to the glutamate-proline interconversion (9, 64, 561, 624) and the appearance of Δ -pyrroline 5-carboxylic acid, the immediate precursor of proline in Equation 11, in glutamate cultures of *Aspergillus oryzae* (486).

Ornithine biosynthesis in *Neurospora crassa* starts from glutamic acid also; the pathways however are not at all clear. Vogel (560) suggests the following sequence in *N. crassa* and *Torulopsis utilis*:



This is rather different from the *Escherichia coli* system (123) which depends upon acetylated intermediates. There may be in *N. crassa* some other pathway of ornithine biosynthesis than that of Equation 12 (153, 560).

Ornithine participates in the ornithine cycle first worked out in animal systems (307) and later found to function in *Neurospora crassa* (156, 490). The basic elements of the cycle, leaving aside biochemical details, are schematized in Figure 8. *N. crassa*, although forming urease, does not liberate urea; we may suppose that the cycle here serves primarily to supply arginine for protein synthesis. Nutritional data suggest the same general pattern in *Penicillium chrysogenum* (61). Urea is a common product of fungi, but direct evidence implicating





Figure 8 The ornithine cycle (499)

the ornithine cycle in net formation of urea has not yet appeared. Ornithine appears in the medium during breakdown of arginine by *Streptomyces venezuelae* (212).

Canavanine, a hydroxyguanidine derivative structurally related to arginine, inhibits the growth of some normal strains of *Neurospora crassa* (251). The inhibition has been traced (576) to the formation of a canavanino succinate which blocks the splitting of arginino succinate.

Of the short chain amino acids—glycine, alanine, serine, and cysteine—only the first named has received much attention in fungi. A glycine requiring mutant of *Neurospora crassa* can grow normally if given either glycolic or glyoxylic acids (613). This is in agreement with animal studies (356–582) and suggests the following sequence:



Drawing on experience with other organisms, we may suggest that the glycolate arises from pentose phosphate (583) and that the amino group is introduced by a transamination from glutamate or other amino donor.

Glycine and serine are closely related in *Penicillium notatum* (26) and *Eremothecium ashbyi* (206–207). Both participate in a variety of biochemical reactions and are clearly related in their biosynthesis. It cannot yet be said which is formed first.




Figure 9 The probable interrelations of aspartic acid, threonine and the sulfur amino acids in *Neurospora crassa*. Several of the steps involve more than one reaction. All amino acids are of the *L* configuration.

Nutritional and metabolic studies on methionine requiring mutants of *Neurospora crassa* established an overall sequence leading from cysteine to methionine (159, 247, 533). A relation of threonine and methionine biosynthesis is suggested by both inhibition studies (13^o) and nutritional requirements of a single gene mutant (533). These relations are schematized in Figure 9 to which has been added the postulate that aspartic acid is the precursor of both threonine and methionine as it is in yeast and bacteria (59). The scheme is by no means complete: there is probably an intermediate between homocysteine and methionine (92) and the source of the methyl group of methionine is not yet clear. An enzyme converting homoserine to threonine occurs in fungi (578a).

Sulfur methylcysteine is formed by *Neurospora crassa* and the isolated compound is utilizable as a source of sulfur (427).

Canavanine has been mentioned as an inhibitor of arginine synthesis. It has also been suggested that canavanine resistant strains convert canavanine to homoserine (329). *Streptococcus faecalis* carries out this reaction (292). Such a conversion would explain why canavanine replaces threonine for some *N. crassa* mutants (532).


Isoleucine and valine are synthesized by *Neurospora crassa* from respectively α, β dihydroxy β ethylbutyric acid and α, β dihydroxy β methylbutyric acid (13). An observed double requirement for both isoleucine and valine (65) has been traced to a deficiency in the dehydrase—possibly dehydrases—which convert these dihydroxy acids to keto acids (381). There is some evidence that the precursors of valine and isoleucine ultimately arise from respectively pyruvic acid and threonine (12, 13, 565).

The keto analogue of leucine, α ketoisocaproic acid, replaces leucine for a mutant of *Neurospora crassa* and is considered therefore to be a precursor of it (432). Earlier steps in leucine biosynthesis by fungi are not known.

Studies on *Neurospora crassa* have shown that both α aminodipic

acid and α -amino- ϵ -hydroxycaproic acid serve as precursors of lysine probably both being converted to a semialdehyde (475-611). Since *Ophiostoma multiannulatum* lysineless mutants utilize α -ketoadipic acid it is probable that this compound is the precursor of α -aminoadipic acid (54).

A series of investigations of histidineless mutants of *Neurospora crassa* is reviewed by Ames (20) the following over all pathway is suggested




Conventional nutritional techniques were of limited use in these studies since the phosphorylated intermediates do not penetrate the cell. More reliance was placed on isolation of intermediates from blocked mutants and on direct enzymatic studies (20-21).

The known early steps of the conversion of glucose to aromatic compounds have been diagrammed in Chapter 6. Figure 10 takes up the story from shikimic acid on, although this presentation is based in large part on studies of *Escherichia coli*; all of the available evidence from *Neurospora crassa* is in agreement with it. In particular a multiple requirement for *p*-aminobenzoic acid, phenylalanine, tyrosine and tryptophan is met by shikimic acid (526). Additional evidence is found in the accumulation of prephenic acid by a strain deficient for the three aromatic amino acids (361) and in the replacement of tyrosine by phenylpyruvic acid (125).

The suggestion that α -phenylglycine is a precursor of tyrosine and phenylalanine (222) has not been borne out by later work (124).

Tryptophan synthesis from anthranilic acid and indole by *Neurospora crassa* is indicated by nutritional responses of deficient mutants (66, 171, 525) and by studies with isotopic nitrogen (408). The conversion of indole to tryptophan is catalyzed by tryptophan synthetase (indole serine ligase); this enzyme has been isolated and partially purified from *N. crassa* (616). The over all reaction which requires pyridoxal phosphate may be written



The mechanism of this reaction is considered by Tatum and Shemin (528). Tryptophan synthesis in *Claviceps purpurea* is probably effected by the same system (548).

Presence or absence of the enzyme is controlled by a single gene in




Figure 10 The biosynthesis of tryptophan phenylalanine tyrosine and p -aminobenzoic acid. This is a general scheme based on data from several organisms but consistent with knowledge of *Neurospora crassa*. Many of the reactions shown are multiple.

Neurospora crassa (241). However, an antigenically related protein, possibly an enzyme precursor or possibly a complex of altered enzyme with an inhibitor, has been isolated from tryptophan requiring mutants (514a). Cultivation of *N. crassa* in a medium deficient in zinc reduces the concentration of the synthesizing enzyme (386, 512).

This summary of amino acid synthesis may give a false impression of finality. There are in fact many complexities and uncertainties which cannot be considered here, and it is to be expected that our picture of these reactions will change as time goes on and more information becomes available.

5 AMINES

Although only one specific decarboxylase activity—glutamic decarboxylation yielding γ aminobutyric acid—is known in fungi (p 265) the number of amines which occur and which presumably originate by decarboxylation of known amino acids is large. Sclerotia of *Claviceps purpurea* yield histamine tyramine isoamylamine agmatine putrescine and cadaverine (503 508 586). Occasional reports—not all certain—suggest that putrescine phenylethylamine isoamylamine and cadaverine occur in fruit bodies of basidiomycetes (291 310 499 580).

A mutant of *Aspergillus nidulans* requires putrescine for growth and the requirement is not met by L ornithine or by other amines (488). Several fungi act on histamine (585). Amines are in general poor or unavailable sources of nitrogen (278).

Muscarine the toxin of *Amanita muscaria* is an amine. Eugster (146) reviews earlier studies a new structure has been confirmed by synthesis (301).


Choline has been found as a constituent of several fungi (508 581) whether however it occurs in the free state or only in phospholipids (Chapter 2) has not been determined. A single report (147) suggests that acetyl choline is a constituent of ergot. Cyclic choline sulfate has been mentioned (Chapter 2).

Barley and other grains scabbed by *Fusarium graminearum* or other related *Fusarium* spp. contain a water soluble material toxic to animals (101) identification of this substance with choline has been shown (473) to be in error.

Choline is required by mutants of *Neurospora crassa* (246) and a mutant can be used for the bioassay of choline (249 527). The synthesis of choline by successive methylations of aminoethanol has been described (Chapter 6).

Trimethylamine has been isolated from numerous basidiomycetes e.g. *Tilletia levis* *Phallus impudicus* and *Russula aurata* (225 499 580) and from *Claviceps purpurea* (508). In plants and in bacteria trimethylamine arises from choline (98) and the same origin may be proposed in fungi.

Betaine $(\text{CH}_3)_3\text{N}^+\text{---CH}_2\text{---COO}^-$ has been isolated from fungi (581) but as with all the amines found in gross studies there is some doubt as to whether it occurs as such or combined. One such combined form of betaine ergothioneine has the structure of the trimethyl betaine of thiohistidine.



Ergothioneine occurs in mammalian blood and seminal fluid (182) in sclerotia and in cultivated mycelium of *Claviceps purpurea* (40 508), and in the mycelium of several cotton fungi (360). Experiments with carbon 14 indicate that ergothioneine and histidine are synthesized by the same pathway (229) and that histidine is not a precursor of either (599).

Amanita mappa fruit bodies contain an amine first called mappin but now known to be identical with bufotenine; its structure 5 hydroxy N dimethyl tryptamine (595) suggests that it is biosynthetically derived from 5 hydroxytryptophan, an amino acid known in other organisms (553 550) but not to date in fungi.

6 METABOLISM OF NUCLEIC ACIDS AND THEIR DERIVATIVES

Nucleic acids, nucleotides, and the purine bases can probably all be used more or less well as sources of nitrogen by fungi (606 607) but the important aspects of nucleic acid metabolism are more narrowly metabolic: the breakdown of nucleic acids and their biosynthesis. Information on nucleic acid breakdown is rather scanty and bears almost exclusively on the purine ribonucleotides. Biosynthetic studies are even less definitive and are limited primarily to nutritional studies of deficient mutants.

Catabolism of Nucleic Acids and Their Derivatives. That fungal enzymes attack nucleic acids has been known for some time. In early experiments liberation of phosphorus or of nitrogen bases or utilization of nitrogen for growth were the criteria, so that it is not possible to specify what enzymes were involved (134 265 291 357 404).

On comparative grounds we may suppose that fungi like bacteria attack nucleic acids first by nucleodepolymerase action, yielding primarily mononucleotides. A few such activities have been shown in fungi (Table I).

The second step in the classical pathway of nucleic acid degradation is the dephosphorylation of nucleotides, yielding inorganic phosphate

Table 1 Enzymes Acting on Nucleic Acids Nucleotides and Nucleosides *

Substrate	Type of Action	Occurrence
Ribonucleic acid	Depolymerization	<i>Streptomyces</i> spp (67 378) <i>Aspergillus oryzae</i> (311)
Deoxyribonucleic acid	Depolymerization	<i>Neurospora crassa</i> (627) <i>Streptomyces</i> sp (378)
Nucleotides	Deamination	<i>Aspergillus</i> spp (68 69 288 311)
Nucleosides	Deamination	<i>Aspergillus oryzae</i> (288 311 370) <i>Neurospora crassa</i> (310)
Nucleosides	Hydrolysis	<i>Aspergillus oryzae</i> (311)

* Dephosphorylation of adenosine phosphates is summarized in Table 1 of Chapter 9

This action is often perhaps usually carried out by non specific phosphatases (316) Hydrolysis of adenosine phosphates summarized in Chapter 9 and the dephosphorylation of riboflavin monophosphate by extracts of *Penicillium chrysogenum* (96) are presumably although not necessarily effected by such phosphatases A phosphatase of *Streptomyces* sp acts both on glycerophosphate and on nucleotide (378)

Diphosphopyridine nucleotide is attacked by a specific enzyme diphosphopyridine nucleotidase which splits off nicotinamide from the dinucleotide leaving adenosine diphosphate ribose The enzyme is formed by *Neurospora crassa* (289) its high concentration in the conidia suggests that it is formed only during sporulation (626) The concentration of enzyme in the mycelium is increased by deficiencies of zinc nitrogen or usually growth factors (386 387) The enzyme of *N. crassa* unlike the corresponding animal enzyme is not inhibited by nicotinamide except at very high concentration and does not catalyze an exchange reaction between the dinucleotide and free nicotinamide (628)

Deamination of nucleotides and nucleosides is usually effected by specific enzymes but an adenyi deaminase of *Aspergillus oryzae* acts on both types of compound deaminating adenosine adenosine 5 phosphate adenosine 3 phosphate adenosine diphosphate and diphosphopyridine nucleotide (288)

These scattered data do not of course give a complete or coherent picture of nucleic acid breakdown they do indicate that catabolism of the purine ribonucleotides in fungi is similar to that in other organisms It seems likely therefore that the free purines adenine hypoxanthine and guanine are converted to xanthine which is then oxidized by xanthine oxidase to uric acid (518) the further metabolism of which is described below


In *Nocardia corallina* the pyrimidines thymine and uracil are metabolized by way of barbituric acid (315) the over all process presumably

also responsible for the catabolism of cytosine and 5-methylcytosine leads eventually in bacteria to urea and malonic acid (474)


Purines and Pyrimidines as Nutritional Factors Naturally occurring fungi with a partial requirement for adenine have been reported (172 174 275 625) among the basidiomycetes: exogenous adenine accelerates growth but has no effect on final yield (174). The effect of guanine and hypoxanthine on spore germination and early growth of *Phycomyces blakesleeanus* is mentioned in Chapter 12. Purine and pyrimidine requiring mutants have been obtained by conventional procedures in several fungi (166 167 170 197 331 410 544) and have been adapted for bioassay (178 330).

For both *Neurospora sitophila* and *Fusarium oxysporum* f. *nicotianae* the purines are generally adequate sources of nitrogen: the pyrimidines poor sources: the methyl purines—caffeine and theophylline—unutilizable (606 607). Xanthine as sole nitrogen source supports good growth of several unrelated fungi (517) and an isolate of *Nocardia corallina* can obtain both carbon and nitrogen from uracil (341).

Biosynthesis of Components of Nucleic Acids It seems certain that purines as such are not synthesized: rather a nucleoside or more complex molecule is probably the first stable compound to accumulate (215). Study of eight adenineless mutants of *Neurospora crassa* led Mitchell and Houlahan (368) to propose the following sequence of reactions:



Hypoxanthine replaces adenine for some mutants of *N. crassa* (414). *Ophiostoma multiannulatum* (166 168 176) and *Ustilago zeae* (410). These data on fungi are consistent with our knowledge of animal systems in which the general pathway appears to be (433):



In *O. multiannulatum* there is some evidence for a role of 4-amino-5-imidazole-carboxamide in purine biogenesis (176): purine reversal of sulfonamide toxicity to *Fremothecium ashbyi* (472) is consistent with this.

Guanosine (or guanylic acid) synthesis in *Ophiostoma multiannulatum* has been suggested to proceed from adenosine (or adenylic acid) i.e. by a further step or steps added to Equation 16 (173). 2,6-diamino-

purine replaces guanine for some mutants of this organism the corresponding riboside does not (175)

Adenylosuccinic acid has been proposed as a precursor of adenylic acid (93) and an adenine requiring mutant of *Neurospora crassa* accumulates both this compound and its riboside (593)

Caffeine is inhibitory at low concentrations to *Ophiostoma multannulatum* since the inhibition is reversed not only by adenine but by structurally unrelated compounds the status of caffeine as a purine antimetabolite is doubtful (253) Inhibition of *Diplodia natalensis* by 8 azaguanine is reversed by adenine suggesting an interconversion in this species of adenine and guanine or their derivatives (295) Other structural analogues of purines cause inhibition of growth in fungi (179 184) but whether they act specifically on normal purine metabolism is uncertain

Biosynthesis of pyrimidines in fungi is even less well understood than that of purines Pyrimidineless mutants respond better to nucleosides than to the free pyrimidine bases (166 170 331) suggesting that ribosidation of a precursor precedes formation of the pyrimidine ring Orotic acid replaces a pyrimidine requirement in *Neurospora crassa* and the importance of this compound is underlined by the finding that pyrimidine requiring mutants accumulate orotic acid or its nucleoside orotidine (362 369) These findings and the demonstrated incorporation of the carbamyl group of citrulline into pyrimidines of *N. crassa* (231) suggest that the following over all pathway of uridylic acid biosynthesis proposed largely on the basis of animal studies (93) is applicable to fungi

Citrulline \rightarrow L ureidosuccinic acid \rightarrow dihydroorotic acid \rightarrow

orotic acid \rightarrow orotidine 5 phosphate \rightarrow uridine 5 phosphate (18)

Cytidylic acid may then be formed by reactions at the triphosphate level although there is some doubt about the occurrence of this interconversion in fungi (447)

7 UREA

Urea is generally recognized as a utilizable nitrogen source it should be realized that only experiments with cold sterilized urea are valid since urea breaks down to ammonia on autoclaving Such experiments show urea utilization by for example several of the Mucorales (31) *Coccidioides immitis* (39) and *Coprinus* spp (165) Although urease is fairly common in actinomycetes utilization of urea has been considered infrequent among *Streptomyces* spp (211)

Urease the enzyme hydrolyzing urea is well known from higher plants and bacteria (514). It is evidently of very general occurrence among fungi including *Aspergillus* spp. (34, 35, 36, 84, 115, 161, 399, 181, 522), *Penicillium* spp. (134, 371, 547), *Pythiactystis citrophthora* (296), several dermatophytic fungi (524) and most of the fleshy basidiomycetes which have been studied in this connection (70, 189, 209, 357). Both direct and indirect evidence indicate that urease is formed by at least some actinomycetes (10, 15, 240, 485).

The urease of *Aspergillus niger* is most active at pH 7.6 (34) although its production is favored to a marked degree by low pH (371). It is formed by *A. niger* on various media and appears in the medium after autolysis (34). In basidiomycete sporophores the enzyme is most abundant in the hymenium (209).


Urea is formed in the fruit bodies of the higher basidiomycetes (209, 268, 269, 579, 615) and by many saprophytic fungi and actinomycetes in pure culture (102, 194, 217, 266, 267, 579). Oxygen and exogenous ammonia increase urea formation by sporophores (272, 273). The function of urea—in some fungi very large amounts are reported (266, 268)—is presumably as a nitrogen reserve.

The biochemical origin of urea in fungi has not been definitely proved and there are several possible processes. First among these is the breakdown of arginine in the ornithine cycle (p. 269). Urea biogenesis by this route is indicated by the finding that exogenous arginine increases urea accumulation in fungi (267, 522) and by the demonstration of urea formation from arginine by cell free preparations from basidiomycete sporophores (273, 615).

Apart from the specific action of arginase on arginine it appears that urea is also formed by one or more enzymes which act on guanidine and derivatives of it to form urea and ammonia; this activity is found in *Streptomyces griseus* (510) and in several saprophytic fungi (103, 270).

Formation of urea from calcium cyanamide by cell free preparations of *Aspergillus niger* has been reported (314).

An origin of urea from purines is also possible in the fungi by the general route



Evidence for this pathway is still incomplete but may be summarized briefly

1. Xanthine oxidase and uricase occur in several fungi (518). Uricase is an inducible enzyme in *Torulopsis utilis* (450).

2 Uric acid has been isolated although possibly as a degradation product from spores of *Aspergillus oryzae* (513)

3 Allantoinase converting allantoin to allantoinic acid is found in cell free extracts of basidiomycete sporophores (83) and *Aspergillus niger* (84) Allantoin serves as a nitrogen source for *Fusarium oxysporum* f. *nicotianae* (607)


4 Sporophores of basidiomycetes may contain significant amounts of allantoinic acid (162)

Proof of the role of this sequence in a single organism and demonstration of allantoinase and its products in a cell free system are required for acceptance of the pathway as written in Equation 19

8 ERGOT ALKALOIDS

At least since the Middle Ages it has been known that the black sclerotia of the ergot fungus cause disease and often death if eaten (41) The alkaloids of the fungus now also occupy a place in pharmacology and have been intensively investigated as drugs (20, 436) Although the classical organism is *Claviceps purpurea* there is reason to believe that sclerotia of other species of the genus especially *C. paspali* also contain the active materials (82)

Chemically the known ergot alkaloids comprise three basic types all amides of lysergic acid which has the structure



In the ergotamine and ergotovine alkaloids lysergic acid is linked to a peptide ergonovine (ergometrine ergobasine) is made up of lysergic acid and L 2 amino 1 propanol (508) A number of compounds closely related to the pharmacologically active materials are produced in culture by specific strains of *Claviceps* spp. (3 509) the structure and possible biosynthetic relationships of these are reviewed by Stickings and Rastick (504) Agroclavine production is greater in still than in

shake culture (5, 6 7) The ergot alkaloids whether produced by the fungus or supplied exogenously disappear in time from a culture (4 363)

9 MISCELLANEOUS NITROGEN COMPOUNDS

Although the antibiotics of actinomycetes form a large share of the known nitrogenous metabolites a number of fungal products are also known Table 2 lists some representative compounds in a classification based on chemical similarity Almost nothing is known of biochemical relationships

Group I of Table 2 includes compounds more or less similar to amino acids Azaserine is of some general interest by reason of its apparently specific blocking of a single step in the biogenesis of purine nucleotides (49 543) Other serine derivatives include the antibiotic cycloserine and the non antibiotic O carbamyl D serine (293) both actinomycete products Mycelianamide produced by *Penicillium griseofulvum* is structurally related as an acylhydroxylamine to cycloserine (57)

The structure of the related antibiotics thiolutin and aureothricin has been determined (97) and their resemblance to cystine has been noted (14)

Compounds more closely and obviously related to the amino acids include fumaryl DL alanine of *Penicillium resticulosum* (58) an unidentified ester of glutamine in *Leptota naucina* (502) and the lichen product picroroccelin a diketopiperazine (27)





The organic nitro group occurs only rarely in biological material chloramphenicol (Chapter 6) is one example β nitropropionic acid another (Table 2) An antibiotic of *Streptomyces thiolutens* has the nitro group (238) *Neurospora crassa* forms an inducible enzyme acting on β nitropropionic acid and related compounds (328)

Indigo is shown in Table 2 as a representative of Group III compounds related to indole There is good evidence that indole is the starting point for indigo formation by both soil bacteria (213) and in the mammalian gut (121) indole in turn we may assume arises from tryptophan

The indole nucleus is found also in lysergic acid (p 279) and in echinulin a product of *Aspergillus glaucus* (425) Gliotoxin produced by *Aspergillus fumigatus* *Penicillium* spp, and *Trichoderma viride* (74) is believed to have a reduced indole nucleus (281) *Penicillium terlikowski* earlier identified as *P. obscurum* forms both gliotoxin and its monoacetate (282)

Fusaric acid is the sole representative of Group IV pyridine deriva





Table 2 Representative Nitrogenous Compounds of Fungi and Actinomycetes

Group*	Compound	Organisms
I	Azaserine	<i>Streptomyces</i> sp (186 187 372 505)
	$\text{N}^+=\text{N}^-=\text{CH}-\text{CO}-\text{O}-\text{CH}_2-\underset{\text{NH}_2}{\text{CH}}-\text{COOH}$	
	Cycloserine (oxamycin)	<i>Streptomyces</i> spp (226 237 309)
		
II	Beta nitropropionic acid	<i>Aspergillus flavus</i> † (88)
	$\text{O}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{COOH}$	
III	Indigo (indigotin)	<i>Schizophyllum commune</i> (365)
		
IV	Fusaric acid	<i>Fusarium</i> spp <i>Gibberella</i> <i>fujikuroi</i> <i>Nectria cinnabarina</i> (191 192 416 507 614)
		
V	Viridicatin	<i>Penicillium</i> spp (75 117)
		

* See p 280

† Also known in higher plants (94)

Table 2 (continued)

Group*	Compound	Organisms
VI	Aspergillic acid	<i>Aspergillus flavus</i> (141 390 591)
		
VII	Cordycepin	<i>Cordyceps militaris</i> (50 118)
		
VIII	Cycloheximide	<i>Streptomyces</i> spp (112 304 313 590)
		
IX	Actithiazic acid	<i>Streptomyces</i> spp (352)
		

tives although lysergic acid also has a reduced N methyl pyridine structure Fusaric acid is a potential wilt toxin in diseases caused by *Fusarium* spp and related fungi Its production by *F. vasinfectum* is narrowly controlled by the zinc content of the medium (287) *Gibberella fujikuroi* also forms dehydrofusaric acid (510)

Viridicatin (Group V Table 2) is a derivative of quinoline quino linic acid is noted later (Chapter 10) as a possible intermediate in the biosynthesis of nicotinic acid from tryptophan Some strains of *Penicillium cyclopium* form viridicatin but in one it is replaced by the related cyclopenin (75)

Aspergillic acid one tautomeric form of which is displayed in Table 2 represents a sixth group derivatives of pyrazine Gliotoxin mentioned above as an indole derivative also contains this grouping as does pulcherriminic acid the iron salt of which pulcherrimin is the red pigment of *Candida pulcherrima* (111 297) *Aspergillus flavus* forms in addition to aspergillic acid the compound flavocol a diiso butyl derivative of deoxyaspergillic acid (140)

The structure of the pyrazines strongly suggests a biosynthetic origin from leucine and isoleucine (74)

Cordycepin yields adenine on hydrolysis Achromycin an antibiotic of *Streptomyces albo niger* also is an adenine derivative (577) Another actinomycete antibiotic amicetin incorporates the pyrimidine cytosine (501)

Nitrogenous quinones include the azanthraquinone chromophore of the actinomycins (p 254) cycloheximide (Table 2) and the pigment phomazarin of *Phoma terrestris* (239 303)

Thiazolidine derivatives are represented only by the antibiotic acti thiazic acid Evidence has been presented indicating that this may act as an antimetabolite of biotin (553)

Polyacetylenic compounds of fungi have been considered briefly (Chapter 6) Of these two—agrocycin (85) and diatetryne 1 (23)—are amides and a third diatetryne 2 (24) is a nitrile

Most pigments so far investigated have proved not to contain nitrogen but several do To those already mentioned may be added as examples the pigments of *Trametes cinnabarina* (216) *Streptomyces coelicolor* (79a) and *S. limosus* (80)

BIBLIOGRAPHY

- 1 Abbott E V 1923 *Soil Sci* 16 207 216
Abderhalden A and Y Teruuchi 1906 *Z Physiol Chem Hoppe Seyler's*
47 394-396

- 3 Abe M 1949a *J Agr Chem Soc Japan* 23 135-139
- 4 Abe M 1949b *J Agr Chem Soc Japan* 23 212-219
- 5 Abe M T Yamano Y Kozu and M Kusumoto 1951 *J Agr Chem Soc Japan* 24 416-422
- 6 Abe M T Yamano Y Kozu and M Kusumoto 1953 *J Agr Chem Soc Japan* 27 18-23
- 7 Abe M T Yamano and M Kusumoto 1953 *J Agr Chem Soc Japan* 2 617-620
- 8 Abel J J and W W Ford 1907 *J Biol Chem* 2 273-288
- 9 Abelson P H and H J Vogel 1955 *J Biol Chem* 213 355-364
- 10 Adachi M and T Imamura 1933 *J Soc Trop Agr (Taiwan)* 5 191 130 (*Biol Abstr* 8 16055 1934)
- 11 Adelberg E A 1953 *Bacteriol Revs* 17 253-267
- 12 Adelberg E A 1955a *J Biol Chem* 216 431-437
- 13 Adelberg E A 1955b In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 419-430 Baltimore The Johns Hopkins Press
- 14 Adelberg E A and M Rabinovitz 1956 *Ann Rev Biochem* 25 349-396
- 15 Afanasiev M M 1937 *Nebraska Univ Agr Exp Sta Research Bull* 9 1-13
- 16 Albert A 1950 *Biochem J (London)* 47 531-538
- 17 Albert A 1952 *Biochem J (London)* 50 690-697
- 18 Allison F E S R Hoover and H J Morris 1934 *J Agr Research* 49 1115-1123
- 19 Amatayakul T 1955 *Ohio J Sci* 55 343-353
- 20 Ames B N 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 357-372 Baltimore The Johns Hopkins Press
- 21 Ames B N and B L Horecker 1956 *J Biol Chem* 220 113-128
- 22 Amos H 1954 *J Am Chem Soc* 76 3858
- 23 Anchel M 1953 *J Am Chem Soc* 75 4621-4622
- 24 Anchel M 1955 *Science* 121 607-608
- 25 Anderson A K 1924 *Studies Biol Sci Univ Minn* 5 237-280
- 26 Arnstein H R V and P T Grant 1954 *Biochem J (London)* 57 353-359
- 27 Asahina Y and S Shibata 1953 *Chemistry of Lichen Substances* Tokyo Japan Society for the Promotion of Science pp 240
- 28 Ayengar P and E Roberts 1952 *J Biol Chem* 197 453-460
- 29 Bach D 1927a *Compt rend (Paris)* 184 766-768
- 30 Bach D 1927b *Compt rend (Paris)* 184 1578-1590
- 31 Bach D 1927c *Compt rend (Paris)* 185 1309-1310
- 32 Bach D 1929a *Bull soc chim biol* 11 119-145
- 33 Bach D 1929b *Bull soc chim biol* 11 995-1006
- 34 Bach D 1929c *Bull soc chim biol* 11 1007-1015
- 35 Bach D 1929d *Bull soc chim biol* 11 1016-1024
- 36 Bach D 1929e *Compt rend soc biol* 100 831-833
- 37 Bach E 1948 *Physiol Plantarum* 1 387-389
- 38 Bach E 1949 *Friesia* 3 377-378
- 39 Baker E E and C E Smith 1942 *J Infectious Diseases* 70 51-53
- 40 Ban R W and E C Stowell Jr 1956 *Arch Biochem Biophys* 63 1-9-260
- 41 Barger G 1931 *Ergot and Ergotism* London Gurney and Jackson pp 2,9
- 42 Barner H D and E C Cantino 1952 *Am J Botany* 39 746-751
- 43 Barnett H L and V G Lilly 1956 *Mycologia* 48 617-627
- 44 Barton Wright E C 1952 *Wallerstein Labs Commun* 1, 117-131

- 4) Baumann G 1933 *Kuhn Archiv* 67 303-393
- 46 Beckman C H J E Kuntz and A J Riker 1933 *Phytopathology* 43 441-447
- 47 Bender A E and H A Krebs 1950 *Biochem J (London)* 46 210-219
- 48 Bender A E H A Krebs and N H Horowitz 1949 *Biochem J (London)* 43 xxi-xxii
- 49 Bennett L L Jr F M Schabel Jr and H E Skipper 1956 *Arch Biochem Biophys* 61 473-486
- 50 Bentley H R K G Cunningham and F S Spring 1951 *J Chem Soc (London)* 1951 2301-2303
- 51 Bentley M L 1953 *J Gen Microbiol* 8 363-377
- 52 Berger J and M J Johnson 1939 *J Biol Chem* 130 641-654
- 53 Berger J M J Johnson and W H Peterson 1937 *J Biol Chem* 117 409-438
- 54 Bergstrom S and R. Rasmussen 1951 *Physiol Plantarum* 4 4.1-4.3
- 55 Bhargava K S 1945 *J Indian Botan Soc* 21 67-72
- 56 Binkley S B 1955 *Ann Rev Biochem* 24 596-670
- 57 Birch A J R A Massy Westropp and R. W Rickards 1956 *J Chem Soc (London)* 1956 3717-3721
- 58 Birkshaw J H H Raistrick and G Smith 1942 *Biochem J (London)* 36 879-883
- 59 Black S and N C Wright 1951 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 591-600 Baltimore The Johns Hopkins Press
- 60 Block S S R L Stephens A Ba reto and W A Murrill 1955 *Science* 121 505-506
- 61 Block S S R L Stephens and W A Murrill 1955 *J Ag Food Chem* 3 584-587
- 62 Blumer S and W H Schopfer 1940 *Ber schweiz botan Ges* 50 248-252
- 63 Boas F 1919 *Ber deut botan Ges* 37 57-6
- 64 Bonner D M 1916 *Am J Botany* 33 788-791
- 65 Bonner D M E L Tatum and G W Beadle 1943 *Act Biochem* 3 1-91
- 66 Bonner D M and C Yanofsky 1951 *J Nutrition* 44 603-616
- 67 Born G V R 1952 *J Gen Microbiol* 6 344-351
- 68 Borsook H and J W Dubnoff 1939a *Enzymologia* 7 256
- 69 Borsook H and J W Dubnoff 1939b *J Biol Chem* 131 163-166
- 70 Bose S R 1939 *Ergeb Enzymforsch* 8 266-276
- 71 Boullenné R and M Boullenné Walrand 1951 *Bull classe sci Acad roy Belg* 3 5 566-567-587
- 72 Boysen Jensen P 1931 *Biochem Z* 239 243-249
- 73 Boysen Jensen P 1937 *Biochem Z* 250 210-280
- 74 Bracken A 1951 *The Chemistry of Microorganisms* London Sir Isaac Pitman and Sons Ltd pp 313
- 75 Bracken A A Pocker and H Raistrick 1951 *Biochem J (London)* 57 83-89
- 76 Brucas F and C Fromageot 1955 *Advances in Protein Chem* 8 1-195
- 77 Briggs W R and P M Ray 1956 *Plant Physiol* 31 165-167
- 78 Brockmann H 1954 *Angew Chem* 66 1-10
- 79 Brockmann H and H Grune 1954 *Chem Ber* 87 1036-1051
- 80 Brockmann H and V Loeschcke 1955 *Chem Ber* 88 778-788
- 81 Brockmann H and H U May 1956 *Chem Ber* 88 419-423
- 82 Brockmann H and H Muxfeldt 1956 *Angew Chem* 67 617-618
- 83 Brown H B 1916 *J Ag Research* 7 401-406
- 84 Brunel A 1931 *Compt rend (Paris)* 190 442-444

- 84 Brunel A 1939 *Bull soc chim biol* 21 380-87
- 85 Bu Lock J D et al 1954 *Chem & Ind* 1954 990-991
- 86 Bünning E 1936 *Flora* (V F) 31 87-112
- 87 Burton K 1951 *Biochem J* (London) 50 258-268
- 88 Bush M T O Touster and J E Brockman 1951 *J Biol Chem* 188 683-693
- 89 Butterworth E M C M Gilmour and C H Wang 1955 *J Bacteriol* 69 725-727
- 90 Cantino E C 1955 *Quart Rev Biol* 30 138-149
- 91 Cantino E C and E A Horestein 1956 *Mycologia* 48 777-799
- 92 Carlson G L and R A McRorie 1955 *Biochim et Biophys Acta* 18 454-455
- 93 Carter C E 1956 *Ann Rev Biochem* 25 123-146
- 94 Carter C L and W J McChesney 1949 *Nature* (London) 164 515-516
- 95 Carter H E C P Schaffner and D Gottlieb 1954 *Arch Biochem Biophys* 53 282-293
- 96 Casida L E Jr and S G Knight 1954 *J Bacteriol* 67 658-661
- 97 Celmer W D and I A Solomons 1955 *J Am Chem Soc* 77 2861-865
- 98 Challenger F 1951 *Advances in Enzymol* 12 429-491
- 99 Chattaway F W C C Thompson and A G E Barlow 1954 *Biochim et Biophys Acta* 14 585-584
- 100 Christensen H N 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 63-106 Baltimore The Johns Hopkins Press
- 101 Christensen J J and H C H Kernkamp 1936 *Minn Univ Agr Exp Sta Tech Bull* 113 1-28
- 102 Chrzyszcz T and M Zakomorny 1934 *Biochem Z* 273 31-42
- 103 Chrzyszcz T and M Zakomorny 1935 *Biochem Z* 275 97-105
- 104 Chughtai I D and T K Walker 1954 *Biochem J* (London) 56 484-487
- 105 Cochrane V W 1950 *Bull Torrey Botan Club* 77 16-180
- 106 Cochrane V W and J F Conn 1947 *J Bacteriol* 54 213-218
- 107 Cochrane V W and J E Conn 1950 *Bull Torrey Botan Club* 77 10-18
- 108 Cohen A L 1953 *Proc Natl Acad Sci U S* 39 68-74
- 109 Cohen P P 1954 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 2 p 1-46 New York Academic Press
- 110 Converse R H 1953 *Mycologia* 45 335-344
- 111 Cook A H and C A Slater 1956 *J Chem Soc* (London) 1956 4130-4133 4133-4135
- 112 Corbar R L Ettlinger W Keller Schuerlein and H Zahner 1957 *Arch Mikrobiol* 26 192-208
- 113 Crasemann J M 1954 *Am J Botany* 41 302-310
- 114 Crewther W G and F C Lennox 1950 *Nature* (London) 165 680
- 115 Crewther W G and F G Lennox 1953a *Australian J Biol Sci* 6 410-427
- 116 Crewther W G and F G Lennox 1953b *Australian J Biol Sci* 6 428-446
- 117 Cunningham K G and G G Freeman 1953 *Biochem J* (London) 53 528-532
- 118 Cunningham K G S A Hutchinson W Man on and T S Spring 1951 *J Chem Soc* (London) 1951 2299-2300
- 119 Curtiss P J H G Hemming and C H Unwin 1951 *Trans Brit Mycol Soc* 34 332-339
- 120 Czapek F 1902 *Beitr Chem Physiol Pathol* 1 538-60
- 121 Dalgliesh C F 1955 *Advances in Protein Chem* 10 33-150
- 122 Darby R T and G R Mandels 1954 *Mycologia* 46 26-28

- 123 Davis B D 1955 *Advances in Enzymol* 16 247-312
- 124 De Busk A G 1956 *Advances in Enzymol* 17 393-416
- 125 De Busk A G and R P Wagner 1953 *J Am Chem Soc* 75 3151
- 126 Delwiche C C 1956 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 218-237 Baltimore The Johns Hopkins Press
- 127 Dimond A E and P E Waggoner 1950a *Phytopathology* 43 193-199
- 128 Dimond A E and P E Waggoner 1953b *Phytopathology* 43 319-321
- 129 Dingle J and G L Solomons 1950 *J Appl Chem (London)* 2 395-399
- 130 Dion W M 1950a *Can J Research C* 28 571-585
- 131 Dion W M 1950b *Can J Research C* 28 586-599
- 132 Doudney C O and R P Wagner 1950 *Proc Natl Acad Sci U S* 38 196-205
- 133 Douglas R J and C L San Clemente 1956 *Can J Microbiol* 2 407-415
- 134 Dox A W 1909 *J Biol Chem* 6 461-467
- 135 Drouhet E 1952 *Ann inst Pasteur* 82 343-355
- 136 Drouhet E 1954 *Congr intern botan Sc Cong Paris Rapports et Commun* 8 Sect 18-20 112-113
- 137 Duggar B M and A R Davis 1916 *Ann Missouri Botanical Garden* 3 413-437
- 138 Dulaney E L 1948 *J Bacteriol* 56 303-313
- 139 Dulaney E L E Bujinski and W B McConnell 1956 *Can J Biochem and Physiol* 34 1193-1198
- 140 Dunn G G T Newbold and P S Spring 1949 *J Chem Soc (London)* 1949 2586-587
- 141 Dutcher J D 1947 *J Biol Chem* 171 341-353
- 142 Dworschack R G H J Koepsell and A A Lagoda 1950 *Arch Biochem Biophys* 41 48-60
- 143 Emerson R L M Putiss and S G Knight 1950 *Arch Biochem* 25 299-308
- 144 Erskson D 1950 *J Gen Microbiol* 6 86-94
- 145 Erkama J and I Virtanen 1951 In J B Sumner and K Myrback (eds) *The Enzymes Vol 1* (2) p 1244-1249 New York Academic Press
- 146 Eugster C H 1956 *Helv Chim Acta* 39 1007-1023 103-1037
- 147 Ewins A J 1914 *Biochem J (London)* 8 44-49
- 148 Fähræus G and V Tullander 1956 *Physiol Plantarum* 9 494-501
- 149 Fairley J L 1954 *Federation Proc* 13 205-206
- 150 Fergus C L 1950 *Mycologia* 44 183-189
- 151 Fincham J R S 1950 *J Biol Chem* 182 61-75
- 152 Fincham J R S 1951 *J Gen Microbiol* 5 793-806
- 153 Fincham J R S 1953 *Biochem J (London)* 53 313-320
- 154 Fincham J R S 1954 *J Gen Microbiol* 11 236-246
- 155 Fincham J R S and A B Boulter 1956 *Biochem J (London)* 62 72-77
- 156 Fincham J R S and J B Boylen 1957 *J Gen Microbiol* 16 438-448
- 157 Fisher W P J Charney and W A Bolhofer 1953 *Antibiotics & Chemotherapy* 1 571-572
- 158 Fleury C 1948 *Ber Schweiz botan Ges* 58 467-477
- 159 Fling M and N J Horowitz 1951 *J Biol Chem* 190 217-225
- 160 Flynn E H J W Hinman E L Caron and D D Woolf Jr 1953 *J Am Chem Soc* 75 5867-5871
- 161 Fosse R 1916 *Ann inst Pasteur* 30 39-755
- 162 Fosse R and A Brunel 1933 *Compt rend (Paris)* 197 288-290
- 163 Foster J W 1949 *Chemical Activities of the Fungi* New York Academic Press pp 618
- 164 Fothgill P G and R Ashcroft 1955 *J Gen Microbiol* 12 387-393

- 165 Fries L 1905 *Svensk Botan Tidskr* 49 475-530
- 166 Fries N 1917 *Arkiv Botan* 33A (7) 1-7
- 167 Fries N 1918 *Hereditas* 34 338-350
- 168 Fries N 1919a *Physiol Plantarum* 2 78-102
- 169 Fries N 1919b *Svensk Botan Tidskr* 43 316-312
- 170 Fries N 1920a *Arkiv Botan Ser 2* 1 271-287
- 171 Fries N 1920b *Physiol Plantarum* 3 180-196
- 172 Fries N 1920c *Svensk Botan Tidskr* 44 319-386
- 173 Fries N 1923 *J Biol Chem* 200 320-333
- 174 Fries N 1924 *Svensk Botan Tidskr* 48 559-578
- 175 Fries N 1925 *Acta Chem Scand* 9 1020
- 176 Fries N S Bergstrom and M Rottenberg 1949 *Physiol Plantarum* 2 210-211
- 177 Fries N and U Bjorkman 1949 *Physiol Plantarum* 2 212-215
- 178 Fries N and A Panders 1920 *Arkiv Botan Ser 2* 1 437-444
- 179 Fromageot C 1921 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 1237-1243 New York Academic Press
- 180 Fromageot C M Jutisz and I Tessier 1919 *Bull soc chim biol* 31 689-695
- 181 Fruton J S and S Simmonds 1923 *General Biochemistry* New York John Wiley and Sons pp 940
- 182 Fry B A 1955 *The Nitrogen Metabolism of Microorganisms* London Methuen and Co pp 166
- 183 Fuerst R C E Somers and T C Hsu 1926 *J Bacteriol* 72 381-393
- 184 Fujii T 1955 *J Biochem (Tokyo)* 42 257-265
- 185 Fusari S A et al 1954a *J Am Chem Soc* 76 2878-2881
- 186 Fusari S A et al 1954b *J Am Chem Soc* 76 2881-2883
- 187 Gale E F 1923 *Advances in Protein Chem* 8 280-301
- 188 Garren K H 1938 *Phytopathology* 28 839-840
- 189 Gaumann E 1921 *Advances in Enzymol* 11 401-431
- 190 Gaumann E 1954 *Endeavour* 13 198-204
- 191 Gaumann E S Naef Roth and H Kobel 1922 *Phytopathol Z* 20 1-33
- 192 Gaumann E S Naef Roth P Reusser and A Ammann 1922 *Phytopathol Z* 19 160-220
- 193 Gentile A C 1921 *Physiol Plantarum* 4 310-386
- 194 Gentile A C and R M Klein 1925 *Physiol Plantarum* 8 291-293
- 195 Georg L K 1922 *Mycologia* 41 470-492
- 196 Giles N H 1926 *J Bacteriol* 57 504
- 197 Gillespie J M and E F Woods 1923 *Australian J Biol Sci* 6 417-46
- 198 Gilmour C M E M Butterworth E I Noble and C H Wang 1923 *J Bacteriol* 69 719-724
- 199 Gilpin R H 1924 *Mycologia* 46 702-707
- 200 Gitterman C O and S G Knight 1922 *J Bacteriol* 61 223-231
- 201 Goddard H N 1913 *Botan Ga* 56 249-30
- 202 Gokhsper J 1921 *Physiol Plantarum* 4 494-513
- 203 Golueke C G 1927 *J Bacteriol* 74 337-344
- 204 Goodman L S and A Gilman 1926 *The Pharmacological Basis of Therapeutics* New York The Macmillan Co pp 1831
- 205 Goodwin T W and O T G Jones 1926 *Biochem J (London)* 61 9-13
- 206 Goodwin T W and S Pendlington 1924 *Biochem J (London)* 57 631-641
- 207 Gorbach G and O G Koch 1925 *Arch Mikrobiol* 23 260-283
- 208 Goss A and P Costy 1922 *Compt Rend (Paris)* 175 539-541 998-999
- 209 Goss W A E Katz and S A Waksman 1926 *Proc Natl Acad Sci U S A* 12 10-12

- 211 Gottlieb D 1953 *Int rn Congr Microbiol 6th Congr Rome* 5 122-136
- 212 Gottlieb D and D Ciferri 1956 *Mycologia* 48 253-263
- 213 Cray P H H 1957 *Proc Roy Soc (London) B* 107 263-280
- 214 Greenberg D M 1951 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 2 p 47-112 New York Academic Press
- 215 Greenberg C R 1951 *J Biol Chem* 190 611-631
- 216 Gripenberg J 1951 *Acta Chem Scand* 5 590-595
- 217 Guillemeau G 1954 *Compt rend (Paris)* 178 1383-138
- 218 Gunther F A et al 1956 *Phytopathology* 46 637-633
- 219 Guyot H 1916 *Bull soc botan Genève Sér 2* 8 80-8
- 220 de la Haba G 1950 *Science* 112 703-704
- 221 Haddock J V C Lilly and H L Barnett 1954 *Mycologia* 46 691-701
- 222 Haddock C H 1952 *Proc Natl Acad Sci U S* 38 48-480
- 223 Hagemann C L Penas e and J Teillon 1955 *Biochim et Biophys Acta* 17 240-243
- 224 Hakamura Y and T Shimomura 1953 *J Agr Chem Soc Japan* 27 694-699
- 225 Hanna W F H B Vickery and G W Pucker 1937 *J Biol Chem* 97 351-358
- 226 Harris D A et al 1955 *Antibiotics & Chemotherapy* 5 183-190
- 227 Haskins R H and W H Weston Jr 1950 *Am J Botany* 37 739-750
- 228 Hawker L E and J Fraymouth 1951 *J Gen Microbiol* 5 369-386
- 229 Heath H and J Wildy 1956 *Biochem J (London)* 64 619-620
- 230 Heinemann B M A Kaplan R D Muir and I R Hooper 1953 *Antibiotics Chemotherapy* 3 1239-1242
- 231 Heinrich M R V C Dewey and C W Kildner 1954 *J Am Chem Soc* 76 3107-3103
- 232 Heppel L A 1951 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 2 p 763-786 New York Academic Press
- 233 Herrick J A 1910 *Ohio J Sci* 40 123-179
- 234 Herrick J A and C J Alexopoulos 1942 *Ohio J Sci* 42 109-111
- 235 Hessayon D G 1952 *Nature (London)* 169 803 804
- 236 Heyn A V J 1935 *Proc Koninkl Acad Wetenschap Amsterdam* 38 1074-1081
- 237 Hidy P H et al 1955 *J Am Chem Soc* 77 2345-2346
- 238 Hirata Y K Okuhara and T Naito 1954 *Nature (London)* 173 1101
- 239 Hoffman Ostenhoff O 1950 *Fortschr Chemie org Naturstoffe* 6 151-241
- 240 Hofmann E and E Latzko 1950 *Biochem Z* 350 264-272
- 241 Hogness D S and H K Mitchell 1954 *J Gen Microbiol* 11 401-411
- 242 Hollis J P 1952 *Phytopathology* 42 273-276
- 243 Hollis J P 1954 *Bull Torrey Botan Club* 81 98-103
- 244 Holter H and B M Pollock 1950 *Compt rend trav lab Carlsberg Sé chim* 28 221-245
- 245 Horowitz N H 1944 *J Biol Chem* 154 141-149
- 246 Horowitz N H 1946 *J Biol Chem* 167 413-419
- 247 Horowitz N H 1947 *J Biol Chem* 171 255-264
- 248 Horowitz N H 1951 *Growth* 15 (Suppl) 47-6
- 249 Horowitz N H and G W Beadle 1943 *J Biol Chem* 150 325-335
- 250 Horowitz N H D M Bonner H K Mitchell F L Tatum and G W Beadle 1941 *Am Naturalist* 79 304-317
- 251 Horowitz N H and A M Srb 1918 *J Biol Chem* 174 371-378
- 252 How J E 1910 *Ann Botany* 4 135-150
- 253 Hultgren B B Kihlman and N Fries 1955 *Physiol Plantarum* 8 493-500
- 254 Hultén E and L Nordström 1949 *Acta Chem Scand* 3 1402-1417

- 255 Iguchi N 1950 *J Agr Chem Soc Japan* 23 357-360
- 256 Isaac I 1919 *Trans Brit Mycol Soc* 32 137-157
- 257 Isenberg H D A Schatz A A Angrist V Schatz and G S Trelawny 1954 *J Bacteriol* 68 5-9
- 258 Isenberg H D G S Trelawny and A Schatz 1953 *Bacteriol Proc (Soc Am Bacteriologists)* 1953 17
- 259 Ishizu K 1953 *Yamaguchi J Sci (Japan)* 4 41-44 (*Chem Abstr* 48 10830 1954)
- 260 Ishizu K 1954 *Yamaguchi J Sci (Japan)* 5 49-55 (*Chem Abstr* 48 12873 1954)
- 261 Isono M 1954 *J Agr Chem Soc Japan* 28 196-200 475-479
- 262 Ito Y 1950a *J Biochem (Tokyo)* 37 51-63
- 263 Ito Y 1950b *J Biochem (Tokyo)* 37 231-247
- 264 Itzerott D 1936 *Flora (N F)* 31 60-86
- 265 Iwanoff L 1903 *Z physiol Chem Hoppe Seyler's* 39 31-43
- 266 Iwanoff N N 1925a *Biochem Z* 157 231-242
- 267 Iwanoff N N 1925b *Biochem Z* 162 425-440
- 268 Iwanoff N N 1927 *Z physiol Chem Hoppe Seyler's* 170 274-288
- 269 Iwanoff N N 1928 *Biochem Z* 192 36-40
- 270 Iwanoff N N and A N Awetissowa 1931 *Biochem Z* 231 67-78
- 271 Iwanoff N N and L K Osnizkaja 1934 *Biochem Z* 271 22-31
- 272 Iwanoff N N and M I Smirnowa 1928 *Biochem Z* 201 1-12
- 273 Iwanoff N N and A Toschewikowa 1927 *Biochem Z* 181 1-7
- 274 Jakoby W B and D M Bonner 1956 *J Biol Chem* 221 689-690
- 275 Jennison M W M D Newcomb and R Henderson 1955 *Mycologia* 47 275-304
- 276 Jensen H L 1930 *Soil Sci* 30 59-77
- 277 Jensen H L 1951 *J Gen Microbiol* 5 360-368
- 278 Jermyn M A 1953a *Australian J Biol Sci* 6 48-69
- 279 Jermyn M A 1953b *Australian J Biol Sci* 6 77-97
- 280 Johnson J L W G Jackson and T E Eble 1951 *J Am Chem Soc* 73 2947-2948
- 281 Johnson J R and J B Buchanan 1953 *J Am Chem Soc* 75 2103-2109
- 282 Johnson J R A R Kidwai and J S Warner 1953 *J Am Chem Soc* 75 2110-2112
- 283 Johnson M J 1934 *Z physiol Chem Hoppe Seyler's* 204 163-175
- 284 Johnson M J and J Berger 1942 *Advances in Enzymol* 2 69-97
- 285 Jones A S A J Swallow and M Webb 1948 *Biochim et Biophys Acta* 167-183
- 286 Kallio R E and A D Larsen 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 616-631 Baltimore The Johns Hopkins Press
- 287 Kalyanasundaram R and L Saraswathi Devi 1955 *Nature (London)* 175 945
- 288 Kaplan N O S P Colowick and M M Ciotta 1952 *J Biol Chem* 194 579-591
- 289 Kaplan N O S P Colowick and A Nason 1951 *J Biol Chem* 191 473-483
- 290 Kating H 1955 *Arch Mikrobiol* 22 396-407
- 291 Keil W and H Bartmann 1935 *Biochem Z* 280 58-60
- 292 Kihara H J M Prescott and E E Snell 1955 *J Biol Chem* 217 497-503
- 293 Kihara H and E E Snell 1952 *J Biol Chem* 197 791-800
- 294 Kikkoji T 1907 *Z physiol Chem Hoppe Seyler's* 51 201-206

- 295 Klein D T and R M Klein 1955 *Plant Physiol* 30 410-413
- 296 Klotz J 1927 *Hilgardia* 3 97-40
- 297 Kluyver A J J P van der Walt and A J van Tret 1933 *Proc Natl Acad Sci U S* 39 583-593
- 298 Kluyver A J and J C M van Zijp 1931 *Antonie van Leeuwenhoek J Microbiol Serol* 17 315-324
- 299 Knight S G 1948 *J Bacteriol* 55 401-407
- 300 Knox W E 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 836-866 Baltimore The Johns Hopkins Press
- 301 Kogl F H C Cox and C A Salemink 1957 *Experientia* 13 137-138
- 302 Kogl F and D G F R Kostermans 1934 *Z physiol Chem Hoppe Seyler's* 208 113-121
- 303 Kogl F and J Sparenburg 1940 *Rec trav chim* 59 1180-1197
- 304 Kornfeld E C R G Jones and T V Parke 1949 *J Am Chem Soc* 71 150-159
- 305 Kostytschew S and E Tswetkova 1930 *Z physiol Chem Hoppe Seyler's* 111 171-200
- 306 Krainsky A 1914 *Centr B kteriolo Pa asienk Abt II* 41 649-688
- 307 Krebs H A and K Henseleit 1932 *Z physiol Chem Hoppe Seyler's* 210 33-66
- 308 Krimsky I and E Racker 1950 *J Biol Chem* 198 721-729
- 309 Kuehl F A et al 1955 *J Am Chem Soc* 77 2344-2345
- 310 Kung A 1914 *Z physiol Chem Hoppe Seyler's* 91 241-250
- 311 Kuninaka A 1955 *J Agr Chem Soc Japan* 29 52-57 797-800 801-805
- 312 Kuntz M 1938 *J Gen Physiol* 21 601-610
- 313 Kupferberg A B H Styles H O Singher and S A Waksman 1950 *J Bacteriol* 59 523-526
- 314 Lemaire Y and A Brunel 1931 *Compt rend (Paris)* 232 872-873
- 315 Lara F J S 1952 *J Bacteriol* 64 271-277 219-85
- 316 Laskowski M 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 956-985 New York Academic Press
- 317 Leaphart C D 1956 *Myologia* 48 2-40
- 318 Leaver F W J Leal and C R Brewer 1947 *J Bacteriol* 54 401-408
- 319 Lebeau J B and J G Dickson 1953 *Phytopathology* 43 581-582
- 320 Lebeau J B and J G Dickson 1955 *Phytopathology* 45 667-673
- 321 Lees H J R Simpson H L Jensen and H Sørensen 1954 *Nature (London)* 173 358
- 322 Lemoigne M and R Descaux 193 *Compt rend (Paris)* 201 239-241
- 323 Leonian L H and V G Lilly 1940 *Am J Botany* 27 18-6
- 324 Lerner A B 1953 *Advances in Enzymol* 14 73-18
- 325 Leuthardt F 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 951-955 New York Academic Press
- 326 Lilly V G and H L Barnett 1931 *Physiology of the Fung* New York McGraw Hill Book Co pp 464
- 327 Lindeberg G 1944 *Symbol e Botan Upsalienses* 8 (2) 1-183
- 328 Little H N 1951 *J Biol Chem* 193 347-358
- 329 Lockhart W R and H R Garner 1955 *Genetics* 40 71-725
- 330 Loring H S G L Ordway and J G Perce 1948 *J Biol Chem* 176 113-1130
- 331 Loring H S and J G Perce 1944 *J Biol Chem* 153 61-69
- 332 McCallan S E A and F R Weedon 1940 *Contribs Boyce Thompson Inst* 11 331-349
- 333 McCarty M 1955 *J Exp Med* 96 555-568
- 334 McCarty M 1956 *J Exp Med* 96 569-580

- 335 McConnell W B 1950 *Can J Research C* 28 600-612
- 336 McConnell W B E Y Spencer and J A Trew 1953 *Can J Chem* 31 697-704
- 337 McElroy W D and H K Mitchell 1946 *Federation Proc* 5 376-379
- 338 McElroy W D and D Spencer 1956 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 131-152 Baltimore The Johns Hopkins Press
- 339 Machlis L 1953a *Am J Botany* 40 189-190
- 340 Machlis L 1953b *Am J Botany* 40 450-459
- 341 Machlis L and E Ossia 1953 *Am J Botany* 40 465-468
- 342 McLamore W M et al 1953 *J Am Chem Soc* 75 105-108
- 343 MacLeod D M 1954 *Ann N Y Acad Sci* 60 (1) 58-70
- 344 MacMillan A 1956a *J Exp Botany* 7 113-116
- 345 MacMillan A 1956b *Physiol Plantarum* 9 470-481
- 346 MacMillan A 1956c *Physiol Plantarum* 9 533-545
- 347 Magasanik B and H R Bowser 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 398-406 Baltimore The Johns Hopkins Press
- 348 Markert C L 1952 *Genetics* 37 603-604
- 349 Martin J K and R D Batt 1955 *Proc Univ Otago Med School (New Zealand)* 33 6-7
- 350 Mathieson M J and D C Catchside 1955 *J Gen Microbiol* 13 83
- 351 Maxwell M E 1950 *Australian J Appl Sci* 1 348-362
- 352 Maxwell M E 1952 *Australian J Sci Research Ser B* 5 42-55
- 353 Mehler A H 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 882-908 Baltimore The Johns Hopkins Press
- 354 Meister A 1955a *Advances in Enzymol* 16 185-246
- 355 Meister A 1955b In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 3-32 Baltimore The Johns Hopkins Press
- 356 Meister A H A Sober S V Tice and P E Fraser 1952 *J Biol Chem* 197 319-330
- 357 Melin E and K Helleberg 1925 *Biochem Z* 151 146-155
- 358 Melin E and P Mikola 1948 *Physiol Plantarum* 1 109-119
- 359 Melin E and B Norkrans 1948 *Physiol Plantarum* 1 176-184
- 360 Melville D P D S Genghof E Inamine and V Kovalenko 1956 *J Biol Chem* 223 9-17
- 361 Metzenberg R L and H K Mitchell 1956 *Arch Biochem Biophys* 64 51-56
- 362 Michelson A M W Drell and H K Mitchell 1951 *Proc Natl Acad Sci U S* 37 396-399
- 363 Michener H D and N Snell 1950 *Am J Botany* 37 52-59
- 364 Michi K and H Nonaka 1954 *J Agr Chem Soc Japan* 28 343-346 346-349
- 365 Miles P G H Lund and J R Raper 1956 *Arch Biochem Biophys* 6 1-5
- 366 Miller T E and R W Stone 1938 *J Bacteriol* 36 248-249
- 367 Mishra J N 1953-54 *Proc Bihar Acad Agr Sci (India)* 1 81-90 (*Chem Abstr* 48 7109 1954)
- 368 Mitchell H K and M B Houlahan 1946 *Federation Proc* 5 370-375
- 369 Mitchell H K M B Houlahan and J F Nyc 1948 *J Biol Chem* 172 525-529
- 370 Mitchell H K and W D McElroy 1946 *Arch Biochem* 10 313-319 311-318
- 371 Miwa T and S Yoshii 1934 *Sci Repts Tokyo Bunrika Daigaku Sect B* 1 243-270 (*Japan J Botany* 7 183 1934)

- 372 Moore J A et al 1954 *J Am Chem Soc* 76 2884-2887
- 373 Morton A G 1956 *J Exp Botany* 7 97-112
- 374 Morton A G and D Broadbent 1955 *J Gen Microbiol* 12 248-258
- 375 Morton A G and A MacMillan 1954 *J Exp Botany* 5 232-252
- 376 Muggleton P W and M Webb 1952a *Biochim et Biophys Acta* 8 431-441
- 377 Muggleton P W and M Webb 1952b *Biochim et Biophys Acta* 8 506-536
- 378 Muggleton P W and M Webb 1952c *Biochim et Biophys Acta* 9 313-355
- 379 Muller D 1944 *Friesia* 3 50-57
- 380 Murray H C and F I Ischeile 1956 *Phytopathology* 46 363-366
- 381 Myers J W and E A Adelberg 1954 *Proc Natl Acad Sci U S A* 40 493-499
- 382 Nakanishi K et al 1954 *Bull Chem Soc Japan* 27 539-543
- 383 Narrod S A and W A Wood 1950 *Arch Biochem Biophys* 3 460-463
- 384 Nason A 1956 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 109-136 Baltimore The Johns Hopkins Press
- 385 Nason A R G Abraham and B C Averbach 1954 *Biochim et Biophys Acta* 15 159-161
- 386 Nason A N O Kaplan and S P Colowick 1951 *J Biol Chem* 188 397-406
- 387 Nason A N O Kaplan and H A Oldewurtel 1953 *J Biol Chem* 201 435-444
- 388 Neal D C R E Wester and K C Gunn 1933 *J Agr Research* 4 107-118
- 389 Neuberg C and I Mendi 1950 *Enzymologia* 14 128-133
- 390 Newbold G T W Sharp and F S Spring 1951 *J Chem Soc (London)* 1951 2679-2682
- 391 Newton W 1916 *Sci Ag* 56 303-304
- 392 Nicholson W N M Nierenstein J C Pool and N V Rice 1931 *Biochem J (London)* 25 75-75
- 393 Nickell I G and P R Burkholder 1947 *J Am Soc Agron* 39 771-779
- 394 Nielsen N 1930 *Jahrb wiss Botan* 73 1-121
- 395 Nord F F 1929 *Ergeb Enzforsch* 8 149-184
- 396 Nord F F and R P Mull 1945 *Advances in Enzymol* 5 165-200
- 397 Norkrans B 1950 *Symbolae Botan Upsalienses* 11 (1) 1-176
- 398 Norkrans B 1953 *Physiol Plantarum* 6 584-593
- 399 Noval J J and W J Nickerson 1956 *Bacteriol Proc (Soc Am Bacteriologists)* 19 6 100-126
- 400 Nyström F 1942a *Compt rend (Paris)* 238 143-145
- 401 Nyström F 1942b *Rev gén botan* 61 285-299 337-338, 416-451
- 402 Ohtsuki T 1936 *Japan J Botany* 8 269-293
- 403 Otani H 1934 *Acta Sci et Med Univ Imp Kyoto (Japan)* 17 42-48 49-59 60-208 69-87 (*Chem Abstr* 29 2005 1935)
- 404 Otani H 1935a *Acta Sci et Med Univ Imp Kyoto (Japan)* 17 303-329 (*Chem Abstr* 9 403 1935)
- 405 Otani H 1935b *Acta Sci et Med Univ Imp Kyoto (Japan)* 17 330-333 (*Chem Abstr* 9 403 1935)
- 406 Otani Y 1937 *Ani Phytothol Soc Japan* 7 9-15
- 407 Page R M 1937 *Ann J Botany* 39 731-739
- 408 Partridge C W H D M Bonner and C Yanofsky 1950 *J Biol Chem* 191 963-968
- 409 Pelletier R L and G W Keitt 1954 *Am J Botany* 41 360-361
- 410 Perkins D D 1949 *Genetics* 34 607-616

- 411 Perlman D 1949 *Am J Botany* 36 180-184
- 412 Peterson E A and H Katznelson 1954 *Can J Microbiol* 1 190-197
- 413 Pfenning N 1956 *Arch Mikrobiol* 25 109-136
- 414 Pierce J G and H S Loring 1945 *J Biol Chem* 160 409-415
- 415 Pistor R 1930 *Zentr Bakteriell Parasitenk Abt II* 80 169-200
- 416 Plattner P A W Keller and A Boller 1954 *Helv Chim Acta* 37 1319-1392
- 417 Plattner P A and U Nager 1948 *Helv Chim Acta* 31 665-671 1949 2203
- 418 Plattner P A U Nager and A Boller 1948 *Helv Chim Acta* 31 594-602
- 419 Plotho O von 1940 *Arch Mikrobiol* 11 34-72
- 420 Pontecorvo G et al 1953 *Advances in Genet* 5 141-238
- 421 Pratt C A 1924 *Ann Botany* 38 599-615
- 422a Pringle R B and A C Braun 1957 *Phytopathology* 47 369-371
- 422b Pringsheim H 1908 *Biochem Z* 8 128-131
- 423 Pyle A J H 1954 *J Gen Microbiol* 11 191-194
- 424 Quantz L 1943 *Jahrb wiss Botan* 91 120-168
- 425 Quilico A C Cardani and F Piozzi 1955 *Gazz chim ital* 85 3-33
- 426 Quispel A 1954 *Acta Botan Neerl* 3 495-511 512-532
- 427 Ragland J B and J L Liverman 1956 *Arch Biochem Biophys* 65 544-576
- 428 Raper K B R D Coghill and A Hollaender 1945 *Am J Botany* 32 165-176
- 429 Ratner S 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 231-256 Baltimore The Johns Hopkins Press
- 430 Ray P M 1956 *Arch Biochem Biophys* 64 193-216
- 431 Ray P M and K V Thimann 1956 *Arch Biochem Biophys* 64 115-190
- 432 Regnery D C 1944 *J Biol Chem* 154 151-160
- 433 Reichard P 1955 In E Chargaff and J N Davidson (eds) *The Nucleic Acids* Vol 2 p 277-308 New York Academic Press
- 434 Reischer H S 1951 *Mycologia* 43 319-328
- 435 Reissig J L 1952 *Arch Biochem Biophys* 36 234-235
- 436 Reynolds A K 1955 In R H F Manske (ed) *The Alkaloids* Vol 5 p 163-209 New York Academic Press
- 437 Reynolds E S 1924 *Am J Botany* 11 215-217
- 438 Richards R R 1949 *Botan Gaz* 110 523-550
- 439 Rippel K 1931 *Arch Mikrobiol* 2 72-135
- 440 Robbins W J and A Hervey 1955 *Mycologia* 47 155-162
- 441 Robbins W J and R Ma 1942 *Am J Botany* 29 835-842
- 442 Robbins W J and R Ma 1945 *Am J Botany* 32 509-523
- 443 Robbins W J and I McVeigh 1946 *Am J Botany* 33 638-647
- 444 Robbins W J A Rolnick and F Kavanagh 1950 *Mycologia* 42 161-166
- 445 Roberts E P Ayengar and I Posner 1953 *J Biol Chem* 203 191-204
- 446 Roberts J L and E Roberts 1959 *Soil Sci* 48 135-139
- 447 Roberts M and D W Visser 1952 *J Biol Chem* 194 695-701
- 448 Ronsdorf L 1935 *Arch Mikrobiol* 6 309-325
- 449 Rother W 1954 *Arch Mikrobiol* 20 89-108
- 450 Roush A H and A J Domnas 1956 *Science* 124 125-126
- 451 Roussos G G H Takahashi and A Nason 1957 *J Bacteriol* 73 594-595
- 452 Roussos G G and L C Vining 1956 *J Chem Soc (London)* 1956 2169-2174
- 453 Ryan F J 1946 *Federation Proc* 5 366-369
- 454 Sakaguchi K. and W Y Chang 1934 *J Agr Chem Soc Japan* 10 459-464 (Chem Abstr 28 58 2-5853 1934)

- 455 Sakaguchi K and C Ishitani 1949 *J Agr Chem Soc Japan* 26 219-285
- 456 Sakaguchi K and S Murao 1950 *J Agr Chem Soc Japan* 23 411
- 457 Sakaguchi K and Y Wang 1936 *Bull Agr Chem Soc Japan* 10 63-69
- 458 Saksena R K and K S Bhargava 1943 *Proc Indian Acad Sci* 18B 45-51
- 459 Saksena R K S K Jain and S M H Jafri 1953 *J Indian Botan Soc* 31 281-286
- 460 Sato R 1956 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 163-175 Baltimore The Johns Hopkins Press
- 461 Schade A L 1940 *Am J Botany* 27 316-384
- 462 Schade A L 1949 *J Bacteriol* 58 811-820
- 463 Schade A L and K V Thumann 1940 *Am J Botany* 27 659-670
- 464 Schatz A and R R Mohan 1955 *J Cellular Comp Physiol* 45 331-342
- 465 Schatz A G S Trelawney V Schatz and R R Mohan 1956 *Biochim et Biophys Acta* 21 391-399
- 466 Schlesinger H and W W Ford 1907 *J Biol Chem* 3 279-83
- 467 Schmalfuss K and K Mothes 1930 *Biochem Z* 221 134-153
- 468 Schmidt E L 1954 *Science* 119 187-189
- 469 Schmitz H and S M Zeller 1919 *Ann Missouri Botan Garden* 6 193-200
- 470 Schopfer W H and S Blumer 1938 *Arch Mikrobiol* 9 305-367
- 471 Schopfer W H and S Blumer 1943 *Beitr schweiz botan Ges* 53 409-456
- 472 Schopfer W H and M Guillaud 1946 *Helv Physiol Pharmacol Acta* 4 24-25
- 473 Schroeter G and L Strassberger 1931 *Biochem Z* 23 452-458
- 474 Schulman M P 1954 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 2 p 223-61 New York Academic Press
- 475 Schweet R S J T Holden and P H Lowy 1954 *J Biol Chem* 211 517-59
- 476 Schweet R S J T Holden and P H Lowy 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 496-505 Baltimore The Johns Hopkins Press
- 477 Semenuk G 1942 *Iowa State Coll J Sci* 16 337-348
- 478 Senn G 1908 *Bol Reus Cambridge Phil Soc* 3 1-91
- 479 Sequeira L and T A Steeves 1954 *Plant Physiol* 29 11-16
- 480 Shepherd C J 1951 *Biochem J (London)* 48 485-486
- 481 Shibata K 1904 *Beitr Chem Physiol Pathol* 5 384-394
- 482 Shirakawa H S 1955 *Am J Botany* 42 319-384
- 483 Silver W S 1957 *J Bacteriol* 73 241-246
- 484 Silver W S and W D McElroy 1954 *Arch Biochem Biophys* 51 379-394
- 485 Simon S 1955 *Acta Microbiol Acad Sci Hung* 3 53-65 (*Chem Abst* 50 7226 1956)
- 486 Simonart P and K Y Chow 1953 *Antonie van Leeuwenhoek J Microbiol Serol* 19 121-134
- 487 Smith E L 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 793-870 New York Academic Press
- 488 Sneath P H A 1955 *Nature (London)* 175 818
- 489 Spitzer G and M M Dehm 1931 *J Agr Research* 43 3-29
- 490 S b A M and N H Horowitz 1944 *J Biol Chem* 154 109-139
- 491 Stahl W H B McQue G R Mandels and R G H Su 1950 *Textile Reseach J* 20 510-519
- 492 Stahl W H B McQue and R G H Su 1950 *Arch Biochem* 27 211-220
- 493 Stanier R Y and O Hayaishi 1951 *Science* 101 316-330
- 494 Steinberg R A 1939 *J Agr Research* 9 731-748
- 495 Steinberg R A 1944 *Science* 100 10
- 496 Steinberg R A and J D Bowling 1939 *J Agr Research* 58 717-730
- 497 Steinberg R A and C Thom 1940 *J Heredity* 31 61-63

- 498 Steinberg R A and C Thom 1942 *J Agr Research* 64 645-659
- 499 Steiner M and E Stein von Kamienski 1953 *Naturwiss* 40 483
- 500 Stetten M R 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 277-290 Baltimore The Johns Hopkins Press
- 501 Stevens C L R L Gasser T K Mukherjee and T H Haskell 1956 *J Am Chem Soc* 78 6212
- 502 Steward F C and J K Pollard 1956 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 371-407 Baltimore The Johns Hopkins Press
- 503 Steward F C R M Zacharius and J K Pollard 1955 *Ann Acad Sci Fennicae Ser A II No 60* 321-366
- 504 Stickings C E and H Raistrick 1956 *Ann Rev Biochem* 25 223-256
- 505 Stock C C et al 1954 *Nature (London)* 173 71-72
- 506 Stockdale P M 1953 *J Gen Microbiol* 8 434-441
- 507 Stodola F H et al 1955 *Arch Biochem Biophys* 54 240-243
- 508 Stoll A 1952 *Fortschr der Chemie org Naturstoffe* 9 114-174
- 509 Stoll A et al 1954 *Helv Chim Acta* 37 1815-1825
- 510 Stoll C 1954 *Phytopathol Z* 22 233-274
- 511 Strauss B S 1951 *Arch Biochem Biophys* 30 292-303
- 512 Strauss B S 1953 *Arch Biochem Biophys* 44 200-210
- 513 Sumi M 1928 *Biochem Z* 193 161-174
- 514 Sumner J B 1951 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (2) p 873-892 New York Academic Press
- 514a Suskind S R 1957 *J Bacteriol* 74 308-318
- 515 Swartz H E and L K Georg 1955 *Mycologia* 47 415-493
- 516 Tabor H 1955 In W D McElroy and H B Glass (eds) *Symposium on Amino Acid Metabolism* p 373-390 Baltimore The Johns Hopkins Press
- 517 Taha E E M and M M Sharabash 1956 *Nature (London)* 177 646-693
- 518 Taha E E M L Storck Krieg and W Franke 1955 *Arch Mikrobiol* 93 67-78
- 519 Tai T Y and W E van Heyningen 1951 *J Gen Microbiol* 5 110-110
- 520 Talley P J and L M Blank 1942 *Plant Physiol* 17 52-68
- 521 Tamiya H 1949 *Advances in Enzymol* 2 183-238
- 522 Tamiya H and S Usami 1940 *Acta Phytochim (Japan)* 11 261-298
- 523 Tanenbaum S W L Garnjobst and E L Tatum 1954 *Am J Botany* 41 484-488
- 524 Tate P 1929 *Parasitology* 21 31-54
- 525 Tatum E L D M Bonner and G W Beadle 1943 *Arch Biochem* 3 477-478
- 526 Tatum E L S R Gross G Ehrensward and L Garnjobst 1954 *Proc Natl Acad Sci U S A* 40 271-276
- 527 Tatum E L M G Ritchey E V Cowdry and L F Wicks 1946 *J Biol Chem* 163 675-682
- 528 Tatum E L and D Shemin 1954 *J Biol Chem* 209 671-675
- 529 Tausson W O 1925 *Biochem Z* 155 356-368
- 530 Tazawa Y K Okunuki and N Urushima 1947 *J Penicillin (Japan)* 1 355-359
- 531 Tazawa Y and S Yamagata 1937 *Acta Phytochim (Japan)* 9 299-310
- 532 Teas H J 1951 *J Biol Chem* 190 369-375
- 533 Teas H J N H Horowitz and M Fling 1948 *J Biol Chem* 172 651-658
- 534 Tempel E 1931 *Arch Mikrobiol* 2 40-71
- 535 Thakur A K and R V Norris 1928 *J Indian Inst Sci 11A* 141-160 (*Chem Abstr* 23 2459 1929)
- 536 Thayer P S and N H Horowitz 1951 *J Biol Chem* 197 755-767

- 537 Thimann K V 1935 *J Biol Chem* 109 279-291
- 538 Thimann K V and H E Dolk 1933 *Biol Zentr* 53 49-66
- 539 Thoai N V J L Hatt and T F An 1956 *Biochim et Biophys Acta* 22 116-123
- 540 Thoai N V J L Hatt T F An and J Roche 1956 *Biochim et Biophys Acta* 22 337-341
- 541 Thornberry H H and H W Anderson 1948 *Arch Biochem* 16 389-397
- 542 Thorne R S W 1950 *Wallerstein Lab Commun* 13 319-340
- 543 Tomisek A J H J Kelly and H E Skipper 1956 *Arch Biochem Biophys* 64 437-455
- 544 Tomizawa C 1950 *Ann Phytopathol Soc Japan* 27 113-118
- 545 Tonhazy N E and M J Pelczar Jr 1954 *Science* 120 141-142
- 546 Treschow C 1944 *Dansk Botan Arkiv* 11 (6) 1-180
- 547 Tsuda N 1950 *Japan J Nutrition* 8 108-110
- 548 Tyler V E Jr and A E Schwartung 1953 *Science* 118 132-133
- 549 Tytell A A J Charney W A Bolhofer and C Curran 1954 *Federation Proc* 13 312
- 550 Udenfriend S and E Tius 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 945-949 Baltimore The Johns Hopkins Press
- 551 Uemura T 1937 *Bull Agr Chem Soc Japan* 13 107-108
- 552 Uemura T 1939 *J Ag Chem Soc Japan* 15 353-358 (*Chem Abstr* 33 8671-8672 1939)
- 553 Umezawa H K Oikawa Y Okami and K Maeda 1953 *J Bacteriol* 66 118-119
- 554 Utkin L M 1950 *Biokhimiya* 15 330-333 (*Chem Abstr* 45 216 1951)
- 555 Vanbreughem R 1950 *Mycologia* 41 176-182
- 556 Verhoeven W 1956 In W D McElroy and H B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 61-86 Baltimore The Johns Hopkins Press
- 557 Verhoeven W and Y Takeda 1956 In W D McElroy and H B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 159-162 Baltimore The Johns Hopkins Press
- 558 Virtanen A I 1950 *Ann Acad Sci Fennicae Ser A II No 39* 1-23
- 559 Vishniac H S 1955 *J Gen Microbiol* 12 455-463
- 560 Vogel H J 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 335-346 Baltimore The Johns Hopkins Press
- 561 Vogel H J and D M Bonner 1954 *Proc Natl Acad Sci U S A* 40 689-694
- 562 Volkonsky M 1934 *Ann Inst Pasteur* 50 76-101
- 563 Waelsch H 1950 *Advances in Enzymol* 13 337-319
- 564 Waggoner P E and A E Dimond 1953 *Phytopathology* 43 281-284
- 565 Wagner R P and A Berquist 1955 *J Biol Chem* 216 251-260
- 566 Wagner R P and P W Island 1956 *Compt rend trav Lab Carlsberg Ser Physiol* 26 381-400
- 567 Waksman S A 1918a *J Bacteriol* 3 509-530
- 568 Waksman S A 1918b *Soil Sci* 6 137-155
- 569 Waksman S A 1919 *Soil Sci* 8 71-715
- 570 Waksman S A 1920 *J Bacteriol* 5 1-30
- 571 Waksman S A 1927 *Principles of Soil Microbiology* Baltimore Williams and Wilkins Co pp 897
- 572 Waksman S A 1944 *Soil Sci* 58 89-113
- 573 Waksman S A 1950 *The Actinomycetes* Waltham Mass The Chronica Botanica Co pp 30
- 574 Waksman S A and R L Starkey 1930 *J Bacteriol* 23 405-408
- 575 Waksman S A and H B Woodruff 1940 *J Bacteriol* 40 581-600

- 576 Walker J B 1953 *J Biol Chem* 201 139-146
- 577 Waller C W P W Fryth B L Hutchings and J H Williams 1953 *J Am Chem Soc* 75 2025
- 578 Ware G C and H A Painter 1955 *Nature (London)* 175 900
- 578a Watanabe Y and K Shimura 1955 *J Biochem (Tokyo)* 42 181-190
- 579 Wehmer C and M Hadders 1933a In G Klein (ed) *Handbuch der Pflanzenanalyse* Vol 4 (1) p 222-228
- 580 Wehmer C and M Hadders 1933b In G Klein (ed) *Handbuch der Pflanzenanalyse* Vol 4 (1) p 247-253
- 581 Wehmer C and M Hadders 1933c In G Klein (ed) *Handbuch der Pflanzenanalyse* Vol 4 (1) p 291-298
- 582 Weinhouse S 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 637-657 Baltimore The Johns Hopkins Press
- 583 Weissbach A and B L Horecker 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 741-742 Baltimore The Johns Hopkins Press
- 584 Went F W and K V Thimann 1937 *Phytohormones* New York The Macmillan Co pp 294
- 585 Werle E 1941 *Biochem Z* 309 61-76
- 586 Werle E 1943 *Die Chemie (Angew Chem)* 56 141-148
- 587 Wetter L R 1952 *Can J Botany* 30 685-692
- 588 Wetter L R 1954a *Can J Biochem and Physiol* 32 20-26
- 589 Wetter L R 1954b *Can J Biochem and Physiol* 32 60-67
- 590 Whiffen A J 1948 *J Bacteriol* 56 283-291
- 591 White E C and J H Hill 1943 *J Bacteriol* 45 433-442
- 592 White N H 1941 *J Australian Council Sci Ind Research* 14 137-146
- 593 Whitfield P R 1956 *Arch Biochem Biophys* 65 585-586
- 594 Wieland T 1949 *Ann Chem Liebigs* 564 159-160
- 595 Wieland T W Motzel and H Merz 1953 *Ann Chem Liebigs* 581 10-16
- 596 Wieland T G Schmidt and L Wirth 1952 *Ann Chem Liebigs* 577 215 233
- 597 Wieland T and W Schon 1955 *Ann Chem Liebigs* 593 157-178
- 598 Wieland T and O Weiberg 1957 *Ann Chem Liebigs* 607 168-174
- 599 Wildy J and H Heath 1957 *Biochem J (London)* 65 220-223
- 600 Williams W J J Litwin and C B Thorne 1955 *J Biol Chem* 219 497 438
- 601 Wilson P W 1951 In C H Werkman and P W Wilson (eds) *Bacterial Physiology* p 467-499 New York Academic Press
- 602 Wirth J C and F F Nord 1943 *Arch Biochem* 2 463-468
- 603 Wolf F T 1948 *Arch Biochem* 16 143-149
- 604 Wolf F T 1951 *Bull Torrey Botan Club* 78 211-220
- 605 Wolf F T 1952 *Proc Natl Acad Sci U S* 38 106-111
- 606 Wolf F T 1953 *Mycologia* 45 825-835
- 607 Wolf F T 1955 *Bull Torrey Botan Club* 82 343-354
- 608 Wood W A and I C Gunsalus 1951 *J Biol Chem* 190 403-416
- 609 Woolley D W 1948 *J Biol Chem* 176 1291-1293 1299-1308
- 610 Wooster R C and V H Cheldelin 1945 *Arch Biochem* 8 311-320
- 611 Work E 1955 In W D McElroy and H B Glass (eds) *A Symposium on Amino Acid Metabolism* p 462-492 Baltimore The Johns Hopkins Press
- 612 Work E and D L Devay 1953 *J Gen Microbiol* 9 394-409
- 613 Wright B E 1951 *Arch Biochem Biophys* 31 332-333
- 614 Yabuta T K Kambe and T Hayashi 1934 *J Agr Chem Soc Japan* 10 1059-1068 (*Chem Abstr* 29 1132 1935)
- 615 Yamamoto S A Fritate and T Miwa 1953 *Botan Mag (Tokyo)* 66 231-238

- 616 Yanofsky C 1950a *J Biol Chem* 194 279-286
- 617 Yanofsky C 1950b *J Biol Chem* 198 343-350
- 618 Yanofsky C and J L Reissig 1955 *J Biol Chem* 209 567-577
- 619 Yaw K E. 1950 *Mycologia* 44 307-317
- 620 Yemm E W 1954 *Proc Symposium Colston Research Soc* 7 p 51-66
New York Academic Press
- 621 Yoshida F 1954 *J Agr Chem Soc Japan* 28 66-70
- 622 Yoshii H 1955 *Bull Sci Fakultato Terkultura Kyusu Imp Univ (Japan)*
6 310-330
- 623 Yoshii H 1956 *Ann Phytopathol Soc Japan* 6 199-204
- 624 Yura T and H J Vogel 1955 *Biochim et Biophys Acta* 17 580
- 625 Yusuf H 1953 *Bull Torrey Botan Club* 80 43-64
- 626 Zalokar M and V W Cochrane 1956 *Am J Botany* 43 107-110
- 627 Zamenhof S and E Chargaff 1949 *J Biol Chem* 180 727-740
- 628 Zatman L J N O Kaplan and S P Colowick 1953 *J Biol Chem* 200
197-210
- 629 Zellner J 1907 *Chemie der höheren Pflanze* Leipzig Wilhelm Engelmann
pp 257
- 630 Zittle C A 1951 In J B Sumner and K Myrback (eds) *The Enzymes*
Vol 1 (2) p 902-913 New York Academic Press

9. Inorganic Nutrition and Metabolism

Inorganic sources of nitrogen have been discussed in the preceding chapter here we are concerned with the utilization of other elements. Fungi have relatively large requirements for phosphorus, potassium, sulfur, and magnesium and much smaller but definite requirements for at least five micronutrients (trace elements, minor elements). The known essential micronutrients are iron, zinc, copper, manganese, and molybdenum. The calcium requirement of fungi, if specific, is of the same order of magnitude as the micronutrient requirement, whereas calcium is a major nutrient for higher plants. Boron, required by green plants, is not known to be needed by fungi. The possibility of other micronutrient requirements in fungi must also be considered. Effects of sodium chloride have been assumed here to be non-specific, although it has been shown that sodium is required by *Anabaena cylindrica* (4).

The history of investigations on the mineral nutrition of fungi is reviewed by Foster (57). Early studies were primarily nutritional in nature and were severely hampered by methodological difficulties. More recent nutritional studies owe much to the pioneering work of R. A. Steinberg on both the techniques and the basic concepts of the field. Interest at the present time is shifting toward the problem of the metabolic function of the inorganic nutrients, with special attention to enzymatic processes in which metals play a role. A return to the exploration of still unknown requirements should be possible as purer

chemicals and inert container materials come to the attention of the biologist

1 PHOSPHORUS

In culture media a level of potassium phosphate (monobasic or dibasic) of 0.004 M is usually provided—reported optimum concentrations—affected of course by the carbon and nitrogen level—are of the order of magnitude of 0.001–0.003 M (75, 100, 187, 241). These levels are however too low to provide adequate buffering against appreciable organic acid formation.

Orthophosphate may be replaced at least for some fungi by other forms of inorganic phosphate (49) by phytic acid (220) by adenosine phosphates (223) or by casein (22). *Aspergillus niger* cannot utilize phosphite or hypophosphite (49); phosphites appear to be generally inert in biological systems (203).

Phosphate deficiencies cause several metabolic disturbances. The most easily observed, as one would expect, is a lowered rate of glucose utilization (33, 123, 130); other metabolic symptoms of deficiency are presumably secondary to this direct effect.

In the typical culture cycle phosphorus is absorbed rapidly during




Fig. 1. Growth and phosphorus changes in *Aspergillus niger*. From data of Mann (135). Seven minute phosphorus is organic phosphorus mineralized by acid hydrolysis of 7 minutes duration.

the early stages of growth and is converted in the cells to organic compounds (69 134) Absorption is dependent upon oxygen and a respirable substrate and is inhibited if respiration is poisoned (79 134 205) The rate of absorption is retarded if available carbon or nitrogen is

Table 1 Phosphatases of Fungi and Actinomycetes

Substrate	Action	Organisms
Beta glycerophosphate sugar phosphates <i>p</i> nitrophenyl phosphate etc	Hydrolysis at acid pH	<i>Actinomyces israeli</i> (90) <i>Aspergillus flavus</i> * (251) <i>A niger</i> (132 134) <i>A oryzae</i> (39 41 42 179) <i>Candida albicans</i> † <i>Cryptococcus</i> spp † <i>Geotrichum candidum</i> † and <i>Torula cremoris</i> † (16) <i>Streptomyces coelicolor</i> (38) <i>Torulopsis utilis</i> (193)
Beta glycerophosphate	Hydrolysis at alkaline pH	<i>Allomyces javanicus</i> † (248) <i>Aspergillus flavus</i> * (251) <i>A oryzae</i> * (7) <i>Candida albicans</i> † (16) <i>Microsporium canis</i> (34) <i>Penicillium chrysogenum</i> (206 207) <i>Torula cremoris</i> † (16) <i>Trichophyton rubrum</i> † (172)
Glucosamine 6 phosphate	Hydrolysis	<i>Neurospora crassa</i> (24)
Adenosine triphosphate	Hydrolysis	<i>Aspergillus niger</i> (112 134) <i>Penicillium chrysogenum</i> (110 111)
Adenosine monophosphate	Hydrolysis	<i>Aspergillus niger</i> (132 134) <i>Penicillium chrysogenum</i> (206)
Monoallyl phosphate	Hydrolysis	<i>Aspergillus niger</i> (8)
Colloidal polymetaphosphate	Depolymerization	<i>Aspergillus</i> spp <i>Penicillium</i> spp † (95 96 97 98 111 132)
Low molecular weight metaphosphate	Hydrolysis	<i>Aspergillus</i> spp (105 106 111 112 135)
Pyrophosphate	Hydrolysis	<i>Aspergillus</i> spp (109 132 135 159 160 164 179) <i>Penicillium chrysogenum</i> (109 206)
Triphosphate	Hydrolysis	<i>Aspergillus niger</i> (132) <i>A oryzae</i> § (159 160 161)
Polyphosphates	Hydrolysis	<i>Aspergillus niger</i> (111 112 113)
Phytic acid	Hydrolysis	<i>Aspergillus niger</i> (50 134) <i>Penicillium chrysogenum</i> (206)
Thionophosphate	Hydrolysis	<i>Aspergillus</i> spp (56)

* Two pH optima

† Histochemical method

‡ No depolymerization by extracts from *Phycomyces blakesleeanae* *Rhizopus nigricans* *Collybia velutipes* *Marasmius* spp *Merulius domesticus* *Polyporus betulinus* or *Tricholoma equestre*

§ Two enzymes

made limiting (71-72) and of course if the external phosphate concentration is lowered (33-70). When conidia appear mycelial phosphorus enters the spores (10). Autolysis is marked by the appearance of phosphate primarily inorganic in the medium (33-42, 143-217) most of this appears to be derived from mycelial organic phosphorus inasmuch as the inorganic mycelial phosphorus remains relatively constant during autolysis. Most of our information is from studies on *Aspergillus niger* the data of Figure 1 are typical.

Combined forms of phosphate organic and inorganic are hydrolyzed by fungal enzymes the phosphatases the occurrence of which is summarized in Table 1. It may be that the acid and alkaline phosphatases of fungi are identical with phosphomonoesterases I and II of other organisms (201) but the specificity is very much in doubt. Indeed there is no certainty in most of the examples of Table 1 that separate specific enzymes are involved.

Probably most or all fungi and actinomycetes form a phosphatase active at low pH against organic phosphates although particular preparations have been reported to lack this activity (34-206-207).

2. POTASSIUM

Potassium is required by fungi at the usual levels of carbohydrate concentrations of 0.001-0.004 M are adequate (100-187-238). Surprisingly little information is available on the physiology of potassium in fungi the most striking effect of potassium deficiency in *Aspergillus niger* is an increase in oxalic acid accumulation (29-146-196). A very low supply of the element is accompanied by poor sugar utilization (195) and it seems likely from work with organisms other than fungi that potassium has an essential function in carbohydrate metabolism (154-176-202). Certain enzymes however are more active in preparations from potassium deficient cells (85).

Sodium can partially replace potassium in the nutrition of *Aspergillus niger* (73-238) but the degree of replacement is small. Under some conditions rubidium and cesium have slight effects (238) under others none (149). In certain of the bacteria rubidium and other alkali metals have a much more pronounced action either as replacements for or competitors of potassium (55-124-130-202-247). It seems likely that studies of this problem in fungi other than *Aspergillus niger* would be rewarding. Rubidium can replace potassium in certain cell free enzyme systems (126).

Figure 2 illustrates the typical growth response to the four major elements—potassium magnesium sulfur and phosphorus.




Figure 2 The response of *Cephalosporium salmosynnematum* to potassium magnesium phosphorus and sulfur Redrawn from Pisano et al (187) by permission of the Williams and Wilkins Company

3 SULFUR

Media in common use incorporate excess sulfur as magnesium sulfate at moderate carbohydrate concentrations the sulfur requirement is about 0.0001–0.0006 M (13, 91, 228)

Most fungi can supply all their needs for sulfur from inorganic sulfate i.e. they reduce the sulfate and incorporate it into organic molecules. The organic sulfur compounds known in fungi include amino acids (cysteine, cystine and methionine), vitamins (thiamine and biotin), cyclic choline sulfate (Chapter 2), sulfur-containing antibiotics e.g. penicillin and gliotoxin and such miscellaneous compounds as junipal (Chapter 6), thiourea (177) and methyl mercaptan (Chapter 6). Sulfur incorporation into organic combinations is often very large, whether or not sulfur is released on autolysis appears to be determined by the species studied (81, 192, 197).

The fungi which as isolated from nature are unable to utilize sulfate sulfur are all aquatic phycomycetes, most Saprolegniales and all known members of the Blastocladales (30, 31). Other phycomycetes—Chytridiales, Leptomitales, Peronosporales and the Zygomycetes—utilize sulfate. Whether dependence on reduced sulfur is biochemically or ecologically related to the lower oxygen requirement or to the fermentative capacity of the Blastocladales is still uncertain. Mutants requiring reduced sulfur have been obtained from *Neurospora crassa*.

(89) *Aspergillus nidulans* (83 188) *Penicillium chrysogenum* (82) *Ophiostoma multiannulatum* (64 65) and *Ustilago zeae* (180)

Some nutritional studies on sulfur are summarized in Table 2 from which it appears that most or all fungi utilize sulfide thiosulfate and

Table 2 The Sulfur Nutrition of Some Fungi*

Source of Sulfur	<i>Sapro- le-gnia delica</i> (22)	<i>Brevi- le-gnia gracilis</i> (22)	<i>Pythium</i> spp (208)	<i>Pestal- lotia malorum</i> (244)	<i>Asper- gillus niger</i> (234)	<i>Penicillium chrysogenum</i> (82)
Sulfate	0	+	+	+	+	+
Persulfate	0	+	+	+	+	
Sulfite	0	+	+			+
Bisulfite	0	+	+	0	+	
Dithionate	0	0	0	+	0	
Hyposulfite	+	+	+	0	+	+
Thiosulfate	+	+	+	+	+	+
Sulfide†	+	+	+	+	+	+
Disulfide†				+	+	+
Cysteine	+	+	+	+	+	+
Cystine					+	+
Methionine					+	
Thiourea	+	+	+	+	0	+
Thiocyanate						+


+ = growth 0 = no growth

† See text

the more oxidized forms of inorganic sulfur (except dithionate). However the data on some of these compounds must be questioned in view of their known instability; thus Steinberg (234) found that *Aspergillus niger* grows much better on old than on fresh samples of sodium sulfide and disulfide.

It should be noted that the difference between *Brevilegnia gracilis* and *Saprolegnia delica* in their sulfur nutrition parallels that seen elsewhere (Chapter 3) in their carbon nutrition—the soil inhabiting *B. gracilis* is more omnivorous than its aquatic relative.

The details of sulfate reduction are not fully known. The available data suggest the following pathway (44)



The obviously necessary intermediates are not known nor is it possible to say at what point in Equation 1 the sulfur becomes attached to an




Figure 2 The response of *Cephalosporium salmosynnematum* to potassium magnesium phosphorus and sulfur Redrawn from Pisano et al (187) by permission of the Williams and Wilkins Company

3 SULFUR

Media in common use incorporate excess sulfur as magnesium sulfate at moderate carbohydrate concentrations the sulfur requirement is about 0.0001–0.0006 M (13, 91, 228)

Most fungi can supply all their needs for sulfur from inorganic sulfate i.e. they reduce the sulfate and incorporate it into organic molecules. The organic sulfur compounds known in fungi include amino acids (cysteine, cystine and methionine), vitamins (thiamine and biotin), cyclic choline sulfate (Chapter 2), sulfur-containing antibiotics e.g. penicillin and gliotoxin and such miscellaneous compounds as junipal (Chapter 6), thiourea (177) and methyl mercaptan (Chapter 6). Sulfur incorporation into organic combinations is often very large, whether or not sulfur is released on autolysis appears to be determined by the species studied (81, 192, 197).

The fungi which as isolated from nature are unable to utilize sulfate sulfur are all aquatic phycomycetes, most Saprolegniales and all known members of the Blastocladales (30, 31). Other phycomycetes—Chytridiales, Leptomitales, Peronosporales and the Zygomycetes—utilize sulfate. Whether dependence on reduced sulfur is biochemically or ecologically related to the lower oxygen requirement or to the fermentative capacity of the Blastocladales is still uncertain. Mutants requiring reduced sulfur have been obtained from *Neurospora crassa*.

(89) *Aspergillus nidulans* (83 188) *Penicillium chrysogenum* (82) *Ophiostoma multiannulatum* (61 65) and *Ustilago zae* (180)

Some nutritional studies on sulfur are summarized in Table 2 from which it appears that most or all fungi utilize sulfide thiosulfate and

Table 2 The Sulfur Nutrition of Some Fungi*

Source of Sulfur	<i>Saprolegnia delica</i> (22)	<i>Brevilegnia gracilis</i> (22)	<i>Pythium</i> spp (208)	<i>Pestalotia molorum</i> (244)	<i>Aspergillus niger</i> (234)	<i>Penicillium chrysogenum</i> (82)
Sulfate	0	+	+	+	+	+
Persulfate	0	+	+	+	+	
Sulfite	0	+	+			+
Bisulfite	0	+	+	0	+	
Dithionate	0	0	0	+	0	
Hyposulfite	+	+	+	0	+	+
Thiosulfate	+	+	+	+	+	+
Sulfide†	+	+	+	+	+	+
Disulfide†				+	+	+
Cysteine	+	+	+	+	+	+
Cystine					+	+
Methionine					+	
Thiourea	+	+	+	+	0	+
Thiocyanate						+


* + = growth 0 = no growth.

† See text

the more oxidized forms of inorganic sulfur (except dithionate). However the data on some of these compounds must be questioned in view of their known instability thus Steinberg (234) found that *Aspergillus niger* grows much better on old than on fresh samples of sodium sulfide and disulfide.

It should be noted that the difference between *Brevilegnia gracilis* and *Saprolegnia delica* in their sulfur nutrition parallels that seen elsewhere (Chapter 3) in their carbon nutrition—the soil inhabiting *B. gracilis* is more omnivorous than its aquatic relative.

The details of sulfate reduction are not fully known. The available data suggest the following pathway (14)



The obviously necessary intermediates are not known nor is it possible to say at what point in Equation 1 the sulfur becomes attached to an

organic molecule but the scheme is in agreement with nutritional data on genetically different mutants of *Neurospora crassa* (89) *Aspergillus nidulans* (83 188) and *Venturia inaequalis* (104) Shepherd (219) proposes a dual pathway of cysteine biosynthesis in *A. nidulans*.

The naturally occurring fungi which require reduced sulfur are we may suppose unable to carry out one or more of the reactions of Equation 1. Thus *Blastocladiella emersonii* grows with either cysteine or methionine (13) and is presumably blocked prior to cysteine synthesis. *Allomyces arbuscula* on the other hand appears unable to use any source of sulfur other than methionine (99). The same considerations apply to other fungi which require cysteine or methionine (131 254).

Bulk reduction of sulfate with evolution of hydrogen sulfide is not effected by fungi as it is by some bacteria. *Candida* spp. reduce sulfite (173) many fungi enzymatically reduce elemental sulfur to H_2S (193 144 216) and *Neurospora crassa* reduces selenite to selenium (259).

Sulfur and sulfur compounds are oxidized by fungi: elemental sulfur by *Penicillium luteum* (1) sulfur-containing amino acids by *Aspergillus niger* (68 148) cysteine by *Microsporium gypseum* (224). Cysteine sulfinic acid has been proposed as an intermediate in cysteine oxidation (224).

A few minor aspects of sulfur metabolism may be mentioned in conclusion. Commercial enzyme preparations from *Aspergillus* spp. contain a phenolsulfatase hydrolyzing compounds of the type $R-OSO_3K$ (2 162 163 165) other sulfatases have not been reported from fungi (66). *Penicillium notatum* is able to split a carbon-sulfur bond (32). The toxicity of barium to *Aspergillus niger* results from the removal of sulfate and barium is not toxic if sulfur is supplied in some other form (228 234). Finally the reduced sulfur requirement of the yeast phase of *Histoplasma capsulatum* (186 209) is a requirement for reducing conditions not for reduced sulfur as such. Transfer of methyl groups to sulfur is summarized in Chapter 6. Reductive steps occur in the overall metabolism of sulfur, selenium and tellurium.

4 MAGNESIUM

Magnesium customarily provided as the sulfate at about 0.001 M probably has as its principal essential function the activation of enzymes necessary to normal metabolism and growth. The optimum is dependent on the concentration of the carbon source (211) the production by the organism of hydroxy acids or other chelate-forming molecules which tend to render magnesium unavailable and the concentrations of other ions to which magnesium is antagonistic. *Asper*

gillus terreus in surface culture requires much more magnesium than it does in shaken culture (119-120)

The enzymes activated by magnesium in *in vitro* systems are often also activated by manganese or other divalent ions but it is believed that magnesium is the physiologically active metal (115-126). None of the physiological effects of magnesium deficiency has been conclusively related to enzyme malfunction but it is reasonable to suppose that for example enhancement of riboflavin formation by magnesium deficiency in *Aspergillus niger* (114-116) reflects an impairment of the normal oxidative system. In bacteria magnesium accelerates the uptake of phosphate possibly by reason of the role of the metal in transphosphorylations (3).

The absorption of magnesium is somewhat slower at neutral than at acid reaction resulting in a slightly higher optimum at neutrality (236). Magnesium is precipitated and rendered virtually unavailable at high pH in the presence of ammonium ion (62).

As suggested above the apparent magnesium requirement is probably affected by other ions. This is deduced from the very great activity of magnesium in antagonizing the toxicity to fungi of such metals as aluminum, copper and mercury (120-122, 136); similar results with other ions have been reported in studies on bacteria (3, 130). The most attractive theory explaining this antagonism is that of MacLeod and Snell (130) that other metals compete with magnesium for enzyme surfaces and that the resulting metal-enzyme complex is inactive or less active than the natural magnesium complex. Competition however need not be limited to enzymatic reactions. Marsh (136) showed clearly that magnesium prevents the absorption of copper by conidia of *Sclerotinia fructicola*; this finding suggests a competition of the two elements at the cell membrane.

Beryllium is said to replace magnesium partially and under special circumstances (238).

5 IRON

Fungi and actinomycetes require approximately 0.1-0.3 ppm of iron more in concentrated media (36, 37, 100, 169, 241). *Aspergillus niger* utilizes either ferrous or ferric iron (168). Iron is lost from solution at neutral or alkaline pH; inclusion in the medium of a chelating agent e.g. citrate reduces the loss (62, 88).

The most obvious metabolic role of iron is in the formation of iron-containing metabolites. Those known in the fungi include catalase and the cytochromes (Chapter 7), coprogen and the ferrichromes

(Chapter 10) the heme pigment of *Neurospora crassa* and *Penicillium notatum* (103) and the pigment pulcherrimin of *Torulopsis pulcherrima* (40 107). Preparations of aspergillin the spore pigment of the black *Aspergillus* spp. contain 0.26 per cent iron (191).

Aspergillus niger releases into the medium relatively large amounts of a substance or substances not citrate which bind ferric iron (67) the possible ecological value is obvious (158).

A few of the metabolic effects of iron may be ascribed to direct use of the element: increases in catalase (257) and in pulcherrimin (201) are of this type. Other metabolic effects of iron concentration e.g. on organic acid formation (45 57 184 185) on penicillin production (108) on the balance of gentisyl alcohol and patulin (Chapter 6) and on sporulation and spore color (118 211) are known but the operative mechanisms are obscure. Only one persistent pattern may be discerned: the maximum formation of certain metabolites—streptomycin (37) penicillin (100) and citrinin (9)—requires more iron than is needed for maximum growth. The data on organic acid accumulation as affected by iron are too much in conflict to allow any generalization (36 57 183) as discussed elsewhere (Chapter 6) it is most probable that deficiencies or excesses of iron and other inorganic constituents act positively on acid formation when they do by virtue of interference with normal metabolism.

6 ZINC

The essentiality of zinc for growth in fungi has been confirmed many times (35). Reported optima determined of course under different conditions range from 0.001 to 0.5 ppm probably 0.5–1.0 ppm is adequate in routine work (5 36 37 76 211). Higher concentrations of zinc may be toxic especially at high pH (117 174) the toxicity is reversed by certain other divalent ions (130). Inhibition of sporulation by zinc is reversed by iron (174 200). High levels of zinc induce genetically stable variation in *Fusarium* sp. (46) and *Helminthosporium sativum* (145). Partial replacement of zinc by cadmium (101) probably results from contamination of the cadmium with zinc (227).

The metabolic effects of zinc deficiency almost defy enumeration: several reviews consider them in detail (35 57 58 183). The major types of metabolic effects are

1. Accumulation of organic acids at levels of zinc too low for normal growth. Associated with this is a positive effect of the metal on the

economic coefficient and on carbon dioxide evolution. The effect of zinc on citric acid formation is in almost all reported instances to decrease the per cent conversion of carbohydrate to citrate (35). This zinc effect is not specific: citric acid accumulation is favored by other mineral deficiencies (36).

2 Lowering of specific enzyme concentrations in cells grown in low zinc media. Enzymes affected include pyruvic carboxylase (59, 212) and ethanol dehydrogenase and the tryptophan-synthesizing enzyme (156). Diphosphopyridine nucleotidase on the other hand is increased by moderate zinc deficiency (156, 157). It should not be assumed however that these effects are specific to zinc.

3 Decrease by zinc of the accumulation of certain antibiotics (141, 245). In *Aspergillus ochraceus* the specific 6 β hydroxylation of steroids is much retarded by zinc deficiency (51).

4 Increase in cytochrome synthesis in *Ustilago sphaerogena* at high zinc (76) with an accompanying decrease in ferrichrome accumulation (158).

5 Changes in cell composition (127, 189, 214).

6 Stimulation by zinc of the *in vitro* formation of vesicles by *Puccinia coronata* (218).

7 A shift in the proportions of related ring compounds produced by *Penicillium urticae* (52).

8 Effects of zinc supply on sporulation reviewed by Foster (57).

Zinc is known to activate a few enzymes specifically (126) and others non specifically (207). Ethanol dehydrogenase and glutamic dehydrogenase are zinc proteins (249, 250). The diverse effects of zinc on metabolism suggest in addition that it is concerned with the synthesis of enzyme proteins: this hypothesis applies especially to the second, third, fourth and seventh effects just listed.

7 COPPER

Copper is required by fungi for normal growth and sporulation at about 0.01–0.1 ppm (36, 37, 100, 241). Higher concentrations are of course toxic although two fungi make visible growth even in a saturated solution of copper sulfate (226). The uptake of copper is somewhat greater at moderate pH than at low pH (215).

Perhaps the most striking physiological effect of copper deficiency is the reduction in pigmentation of the colored spores of *Aspergillus* spp (26, 102, 142, 150) and *Trichoderma viride* (27). Although other metal

deficiencies also affect spore pigmentation (58) copper is probably the most important at least for the dark spore pigments

It has been mentioned earlier (Chapter 6) that copper may play a role in chlorine metabolism of fungi. The possibility that as in animals (134) copper is essential for normal iron metabolism has not been explored

8 MANGANESE

The essentiality of manganese in small amounts about 0.005–0.01 ppm has been demonstrated for a number of fungi (23, 36, 53, 57, 86, 137, 190, 199, 227, 241). We may assume that fungi in general require it although the effects of deficiency are not apparent for all. The requirement is more easily shown at low pH than at high (121).

Manganese deficiency has several physiological effects (57, 183) including a pronounced decrease in sporulation. The frequent role of the metal in activation of enzymes especially those of the citric acid cycle (136) suggests a role for it in metabolism at this point. Manganese concentration also affects the concentration of other enzymes in the cell (86).

The biological oxidation of manganese in soil can be demonstrated perhaps best in soil percolation experiments (129). It is probable that fungi as well as other organisms carry out this reaction. The limited evidence is reviewed by Mulder and Gerretsen (153) and by Starkey (225).

9 MOLYBDENUM

The essentiality of molybdenum for *Aspergillus niger* was discovered by Steinberg (229). The requirement is small—estimates range from 0.1 parts per billion (152, 168) to 10 ppb (241)—and *A. niger* responds to as little as 0.02 ppb (152). Vanadium does not replace molybdenum (168).

It has been observed repeatedly that the molybdenum requirement of both *Aspergillus niger* and *Neurospora crassa* is higher if nitrate is used as the source of nitrogen than if ammonium nitrogen is provided (152, 170, 230). The explanation for this phenomenon is the first unequivocal demonstration of a specific enzymatic basis for a deficiency symptom in fungi. Molybdenum participates as electron carrier in the reduction of nitrate by enzymes of *Neurospora crassa*, *Escherichia coli* and soybean. The evidence for this conclusion is reviewed by Nason

(155) The valence states of molybdenum which function in electron transport are the +5 and the +6 (171)

Molybdenum must however have other functions inasmuch as a molybdenum deficiency can be demonstrated in media not containing nitrate (152 168 170)

10 CALCIUM

The effect of calcium on the growth of fungi has only recently been recognized although the first report of such an effect was in 1922 (258) Species in 19 genera including representatives of all the major groups of fungi and of the genus *Streptomyces* have been found to respond to added calcium by an increase in dry weight (62 63 73 99 113a 117 128 166 175 178 198 239 246 253) Reported optima range from 0.5 to 20 ppm Some fungi it should be noted cannot be shown to require calcium (198 249) Figure 3 illustrates the response to calcium

Strontium partially replaces calcium for *Allomyces arbuscula* (99) and several other fungi (117 198 239) Strontium is firmly bound by mycelium of *Rhizopus* sp (116)

One may still ask however whether the so-called calcium require




Figure 3 The growth of *Coprinus ephemerus* at different levels of calcium (milligrams per liter) Redrawn from Fries (63) by permission of the Svensk Baktisk Tidsskrift

ment of fungi is specific. Calcium in many organisms including fungi (117) protects against the injurious effects of certain common monovalent cations especially hydrogen sodium and potassium. In bacteria calcium and related ions also protect against zinc toxicity (121). These non nutritional effects the wide range of optima reported the extent of replacement by strontium and the fact that some fungi do not respond at all to calcium all argue for some caution in interpretation of the data as reflecting an absolute and specific requirement. It is noticeable that in a casamino acids medium with high sodium concentration calcium affects the growth of *Aspergillus oryzae* (222) but the related *A. niger* in the usual low sodium media does not respond to calcium (239). The crucial experiment would be of course to determine whether an apparent calcium requirement can be eliminated by adjusting the concentration of other ions.

It has been found that calcium affects the time required for perithecial formation in *Chaetomium* spp. (14, 15) and the intensity of conidiation in *Trichoderma viride* (27).

Calcium like magnesium may be made less available if chelating agents e.g. citrate are present in the medium and is less available because of precipitation at high pH than at low.

11 COBALT

In spite of early claims of the essentiality of cobalt for fungi reviewed by Marston (138) there is at present no conclusive evidence of a cobalt requirement: some more recent data are suggestive however (11). Cobalt is required by animals but not apparently by higher plants (138). As mentioned elsewhere (Chapter 10) vitamin B₁₂ is made by *Streptomyces* spp.—and for the synthesis cobalt must be required—but whether the vitamin is produced by true fungi is uncertain. Furthermore neither the fungi nor the actinomycetes are known to require the vitamin. Although cobalt activates some enzymes e.g. the dipeptidase of *Aspergillus oryzae* (42) its only known essential function in organisms generally is as a constituent of vitamin B₁₂. In view of the very great technical difficulty in establishing a cobalt requirement if the metal is only needed for B₁₂ synthesis (91) it may be more profitable to explore the B₁₂ metabolism of fungi than to attack the cobalt problem directly. Preliminary results (168) suggest that neither B₁₂ nor cobalt is essential to *Aspergillus niger*.

Cobalt is accumulated by *Neurospora crassa* (12) the degree of accumulation—up to 23 times the external concentration—being a function of the amount of cobalt in the medium. Up to 40 per cent of the

cobalt so accumulated is bound to a protein. Cobalt accumulation in *N. crassa* is affected indirectly by the iron concentration (11).

Cobalt at appreciable concentrations is toxic to fungi and other organisms. Toxicity is reduced by histidine (181, 182, 212) which presumably forms a complex with the cobalt ion. The uptake of cobalt by *Aspergillus niger* is sharply reduced by the addition of relatively small amounts of magnesium (3). A third complication in cobalt toxicity is the observation that although iron and cobalt do not simply compete, the enzymatic symptoms of cobalt toxicity in *Neurospora crassa* resemble closely the symptoms of iron deficiency (80).

12 OTHER INORGANIC NUTRIENTS

Aspergillus niger responds slightly to vanadium (17, 18, 19) and fleshy fungi take up vanadium from the soil (20) but satisfactory proof of an essential role is not at hand. Vanadium is essential however for the alga *Scenedesmus obliquus* (6).

Steinberg has reported growth increases from addition of gallium (231) and—only with glycerol as carbon source—of scandium (233). Results on gallium have not however proved reproducible (237).

Claims of an essential or growth promoting role of boron (256) and organic compounds of silicon (87) have not as yet been confirmed. Winfield (255) reports negative results with boron. Chlorine appears to have no necessary role in the nutrition of fungi (60) although chlorinated metabolites are not uncommon (Chapter 6) and chlorine is essential to higher plants (28). Other metals e.g. aluminum and chromium often exert physiological effects and may even under peculiar circumstances improve growth but it has not been suggested that they are essential.

13 BIOASSAY OF MINERALS

The response of fungi to nutrients is sensitive enough so that a number of bioassays for inorganic elements in soil have been developed to the point of practical utility. Almost all the work has been with *Aspergillus niger* although a radial growth assay for phosphorus with *Cunninghamella* spp. was devised in 1934 (139, 140).

Bioassays for minerals are more laborious than chemical determinations and are intrinsically less accurate: precision is especially low in those assays based on visual estimates of growth or spore color (151). The principal advantage of the bioassay method and the reason for its popularity is that it may more nearly approximate a determination of




Figure 4 The form of assay curves employing *Aspergillus niger* for zinc molybdenum and copper Redrawn from D J D Nicholas *The Analyst* Vol 77 p 699 649 (1952) by permission of the Society for Analytical Chemistry

the minerals actually available to a crop plant. The assumption made therefore is that the fraction of a given mineral which the plant cannot absorb is equally unavailable to the fungus. Consequently the major criterion for the practicability of a bioassay must be the degree to which the assay results predict the behavior of crop plants in a given soil. Correlation with chemical determinations is of secondary importance. Correlative studies have usually shown at least fair agreement between growth of *Aspergillus niger* and growth of higher plants in soils deficient in phosphorus potassium zinc copper and molybdenum (47 73 147 150 168). The fungus is less suitable for manganese determination because metabolic acids make more manganese available to it than is available for higher plants (153).

Typical assay curves are shown in simplified form in Figure 4. It is to be noted that the response is not rectilinear. Detailed methods are described by Donald et al (47) Gerretsen (73) Mulder (151) and Nicholas (168).

14 PROBLEMS IN MINERAL NUTRITION

Two general approaches may be used in determining the essentiality of a given element. One may design a medium free of the element in question and containing all other materials necessary for growth. Essentiality is shown by a quantitatively determined growth response to graded amounts of some soluble compound containing the element. Obviously the medium must be purified rigorously for the study of

micronutrients and equally important the compound added must not contain any growth promoting substances

The second approach little used so far is to determine that some metabolite which contains the element is essential to growth. For example if the vitamins B₁₂ prove to be essential then cobalt must be essential.

Details of the methods for purification of media for studies of micronutrient elements lie outside the scope of this work. There is no one best method for all materials and the choice depends on the particular element under study. The most useful over all methods involve adsorption on calcium carbonate (240) or aluminum compounds (48, 221). Special methods have been described for copper (151, 167, 168), manganese (168) and molybdenum (81, 168). It appears that the chelating agents (8-hydroxyquinoline, diphenylthiocarbazone, etc.) are not only less efficient for most micronutrient elements but are likely to be toxic to the organism. Detailed comparisons of different methods have been published by Steinberg (240), Donald et al. (48) and Nicholas (168).

Methods of obtaining inorganic salts of maximum purity are discussed by Hutner et al. (94).

The sources of contamination in micronutrient work are well known by now: the inoculum, glassware, distilled water from a metal still, sugars and polyatomic alcohols, hydroxy acids, inorganic salts and complex extracts of biological origin (92, 168, 229, 230, 232). Amino acids are often so heavily contaminated that they cannot be employed at all (235).

Chelating agents may be used to exaggerate a requirement and make it more readily detectable. Thus Hutner (93) added citrate to hold calcium and other elements in solution: the calcium requirement in the absence of the complexing anion is estimated by extrapolation.

Work with the major inorganic nutrients is of course somewhat simpler inasmuch as deficiencies are not made up by minor amounts of contamination. Methodological problems do however arise even here. Thus the usual practice in working at low or zero potassium is to supply phosphate as the sodium salt: this of course ignores the possibility that sodium partially replaces potassium. In this particular instance the solution is to use an amine in place of the alkali metal to control pH (126).

In reporting results the concentration permitting half maximal growth is a more useful datum than that at which maximum growth occurs.

Although an optimum amount of a given element may be defined for rigidly specified conditions (236, 237) it must be realized that there

is no absolute optimum concentration of an element for even one species. Some of the complicating factors may be mentioned. First the need for inorganic nutrients is proportional to the carbohydrate supply (241). Second the need for an element may be increased or decreased by the presence of a second element, the important nutritional factor becoming then the ratio of the two (60, 129). In practice this means that equally good growth is observed over a range of total salt concentration (43, 77, 128). Third the nutrients and metabolites which form chelate complexes with minerals are present in continually changing amounts, although a chelating agent may on the one hand solubilize a mineral and so make it more available, on the other hand the cell must compete with the chelate complex for the element. Fourth the pH of the medium—again a factor not usually constant for the entire life cycle—affects the solubility of most inorganic salts (62, 78, 88, 230), the state of oxidation of manganese (225) and the stability of chelated complexes (25). Finally changes in the conditions of culture may change the apparent requirement for a mineral, as in the case of the magnesium requirement of *Aspergillus terreus* (119, 120).

The use of metabolically inert chelating agents, particularly ethylenediaminetetraacetic acid (EDTA, versene) solves some of the practical problems of supplying a fungus with minerals; the compound has been used successfully at 2–5 ppm for several fungi (194, 209a, 252) and for *Streptomyces nitificans* (213). However this and similar agents may prove to be toxic (63, 128, 243). Added citrate or malate, unless utilized by the organism, can be used to keep minerals soluble (92).

The fungi accumulate minerals against a concentration gradient; the data are particularly clear for cobalt (11, 12), silver and cesium (144) and potassium (54). These examples and the accumulation of rubidium by fleshy basidiomycetes (21) make it clear in addition that accumulation of non-essential elements is common. The energy for accumulation must come of course from respiration. The very striking effect of magnesium on the uptake of cobalt by *Aspergillus niger* (3) and of copper by *Sclerotinia fruticola* (132) suggests that reactions at a semi-permeable membrane are of importance in mineral nutrition. In yeast there is good evidence for carrier systems in the transport of cations across the membrane (61).

The functions of the minerals required by fungi deserve further study. Phosphorus and sulfur occur in known essential compounds and their major role appears to be constitutive. The other essential mineral elements are today more or less conventionally assigned a catalytic function as activators or active centers of enzymes. Reasons exist of course for this general belief: the function of molybdenum in

nitrate reductase and zinc in ethanol dehydrogenase the occurrence of copper and iron in purified enzymes and the known activating effect of magnesium manganese and calcium on isolated enzymes. Further study of possible enzymatic roles will undoubtedly be rewarding but too narrow a dedication to this task may mean neglect of other as yet unknown physiological reactions of the metals

BIBLIOGRAPHY

- 1 Abbott E V 1923 *Soil Sci* 16 207-216
- 2 Abbott L D Jr 1947 *Arch Biochem* 15 205-214
- 3 Abelson I H and E Aldous 1950 *J Bacteriol* 60 401-413
- 4 Allen M B and D I Arnon 1955 *Physiol Plantarum* 8 653-660
- 5 Anderson Kotô I and G C Hevesy 1949 *Biochem J (London)* 44 407-409
- 6 Arnon D I and G Wessl 1953 *Nature (London)* 172 1039-1040
- 7 Auhagen E and S Grzycki 1953 *Biochem Z* 265 217-222
- 8 Baba S 1913 *Proc Imp Acad (Tokyo)* 19 70-76
- 9 Bailey J M and C J Cavallito 1913 *J Bacteriol* 45 30-31
- 10 Bajaj V S F Damle and P S Krishnan 1954 *Arch Biochem Biophys* 50 451-460
- 11 Ballantine R 1953 *J Cellular Comp Physiol* 42 413-426
- 12 Ballantine R and D G Stephens 1951 *J Cellular Comp Physiol* 37 369-387
- 13 Barner H D and E C Cantino 1929 *Am J Botany* 39 746-751
- 14 Basu S N 1951 *J Gen Microbiol* 5 231-238
- 15 Basu S N 1952 *J Gen Microbiol* 6 199-201
- 16 Bayliss M D Glick and R Siene 1918 *J Bacteriol* 53 307-316
- 17 Bertrand D 1941a *Bull soc chim biol* 23 467-471
- 18 Bertrand D 1941b *Compt rend (Paris)* 213 254-257
- 19 Bertrand D 1942 *Ann inst Pasteur* 68 226-244
- 20 Bertrand D 1913 *Bull soc chim biol* 25 194-197
- 21 Bertrand G and D Bertrand 1949 *Ann inst Pasteur* 76 199-202
- 22 Bhargava K S 1915 *Proc Indian Acad Sci B* 21 314-319
- 23 Blank L M 1941 *J Agr Research* 62 179-180
- 24 Blumenthal H J A Hemerline and S Roseman 1956 *Bacteriol Proc (Soc Am Bacteriologists)* 1956 109
- 25 Bohtelsky M and J Jorlan 1913 *J Am Chem Soc* 67 1804-1831
- 26 Bortels H 1927 *Biochem Z* 182 301-308
- 27 Brian P W and H G Hemming 1950 *Trans Brit Mycol Soc* 33 132-141
- 28 Broyer C C A B Carlton C M Johnson and P R Stout 1954 *Plant Physiol* 29 56-57
- 29 Butkevitch W S (Butkevich V S) and A C Timofeeva 1934 *Microbiol ogya* 3 574-581
- 30 Cantino E C 1920 *Quart Rev Biol* 25 269-277
- 31 Cantino E C 1925 *Quart Rev Biol* 30 133-149
- 32 Challenger F and Y C Liu 1950 *Rec trav chim* 69 334-342
- 33 Chang S C 1910 *Soil Sci* 49 197-210
- 34 Chattaway F W C C Thompson and A G E Barlow 1951 *Biochim et Biophys Acta* 14 583-584
- 35 Chesters C G C and G N Robinson 191a *Biol Revs Cambridge Phil Soc* 26 239-5

- 36 Chesters C G C and G N Rolinson 1951b *J Gen Microbiol* 5 553-558
- 37 Chesters C G C and G N Rolinson 1951c *J Gen Microbiol* 5 559-565
- 38 Cochrane V W and P L Hawley 1956 *J Bacteriol* 71 308-314
- 39 Contardi A and A Ercole 1933 *Biochem Z* 261 275-309
- 40 Cook A H and C A Slater 1956 *J Chem Soc (London)* 1956 4130-4133
4133 4135
- 41 Courtois J and G Joseph 1917 *Bull soc chim biol* 29 951-955
- 42 Crewther W G and F G Lennox 1953 *Australian J Biol Sci* 6 410-477
- 43 Darby R T and G R Mandels 1954 *Mycologia* 46 276-288
- 44 Davis B D 1955 *Advances in Enzymol* 16 247-312
- 45 Di Capua A 1933 *Gazz chim ital* 63 296-302
- 46 Dimock A W 1936 *Zentr Bakteriell Parasitenk Abt II* 95 341-341
- 47 Donald C B I Passey and R J Swaby 1952a *Australian J Agr Research*
3 305-325
- 48 Donald C B I Passey and R J Swaby 1952b *J Gen Microbiol* 1 911-
220
- 49 Dox A W 1911 *J Biol Chem* 10 71-80
- 50 Dox A W and R Golden 1911 *J Biol Chem* 10 183-186
- 51 Dulaney E L E O Stapley and C Hlavac 1955 *Mycologia* 47 464-474
- 52 Ehrensward G 1955 *Exp Cell Research Suppl* 3 102-109
- 53 English M P and N H Barnard 1955 *Trans Brit Mycol Soc* 38 78 89
- 54 Eno C F and H W Reuzer 1950 *Soil Sci* 80 199-209
- 55 Feeney R E and J A Garibaldi 1948 *Arch Biochem* 17 447-458
- 56 Forrest I S A Grauer and C Neuberg 1954 *Enzymologia* 16 305-310
- 57 Foster J W 1939 *Botan Rev* 5 207-239
- 58 Foster J W 1949 *Chemical Activities of Fungi* New York Academic Press
pp 648
- 59 Foster J W and F W Denison Jr 1950 *Nature (London)* 166 833-834
- 60 Fothergill P G and R Ashcroft 1955 *J Gen Microbiol* 12 381-395
- 61 Foulkes E C 1956 *J Gen Physiol* 39 687-704
- 62 Fries L 1945 *Arkiv Botan* 32 (10) 1-8
- 63 Fries L 1956 *Scensk Botan Tidskr* 50 47-96
- 64 Fries N 1945 *Arkiv Botan* 32 (8) 1-9
- 65 Fries N 1946 *Svensk Botan Tidskr* 40 127-140
- 66 Fromageot C 1950 In J B Sumner and K Myrback (eds) *The Enzymes*
Vol 1 (1) p 517-526 New York Academic Press
- 67 Garibaldi J A and J B Neilands 1956 *Nature (London)* 177 596-597
- 68 Garreau Y 1941 *Compt rend soc biol* 135 508-510
- 69 Gayet J 1948a *Bull soc chim biol* 30 488-496
- 70 Gayet J 1948b *Bull soc chim biol* 30 542-547
- 71 Gayet J 1949a *Bull soc chim biol* 31 796-800
- 72 Gayet J 1949b *Bull soc chim biol* 31 1046-1051
- 73 Gerretsen F C 1948 *Anal Chim Acta* 2 782-799
- 74 Gollmick F 1936 *Zentr Bakteriell Parasitenk Abt II* 93 491-442
- 75 Golueke C G 1957 *J Bacteriol* 74 337-344
- 76 Grimm P W and P J Allen 1954 *Plant Physiol* 29 369-377
- 77 Haenseler C M 1921 *Am J Botany* 8 147-163
- 78 Halvorson H O and R L Starkey 1927 *J Phys Chem* 31 626-631
- 79 Harley J I C C McCready and J K Brierley 1953 *New Phytologist* 5
194-132
- 80 Healy W B S Cheng and W D McElroy 1955 *Arch Biochem Biophys*
54 206-214
- 81 Hewitt E J and D G Hallas 1951 *Plant and Soil* 3 466-408
- 82 Hockenhull D J D 1948 *Biochem J (London)* 43 498- 01
- 83 Hockenhull D J D 1949 *Biochim et Biophys Acta* 3 326-33
- 84 Hockenhull D J D 1950 *J Exp Botany* 1 194-200

- 8 Hofmann E and H Scheck 1950 *Biochem Z* 321 98-106
- 86 Hofmann E, H Scheck and K Saffert 1950 *Biochem Z* 320 126-13
- 91 Holzapfel L and W Engel 1954 *Z Naturforsch* 9b 600-606
- 88 Hopkins E F and F B Wann 1956 *Botan Gaz* 81 353-356
- 99 Horowitz N H 1950 *Advances in Genet* 3 33-51
- 90 Howell A Jr and R J Fitzgerald 1953 *J Bacteriol* 66 437-44
- 91 Hungate F P and T J Mannell 1950 *Genetics* 35 709-719
- 92 Hutner S H 1946 *J Bacteriol* 52 13-291
- 93 Hutner S H 1948 *Trans N Y Acad Sci* 11 10 136-141
- 94 Hutner S H, L Provasoli, A Schatz and C P Haskins 1950 *Proc Am Phil Soc* 94 1 2 1-10
- 95 Ingelman B 1950 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 511-516 New York Academic Press
- 96 Ingelman B and H Malmgren 1947 *Acta Chem Scand* 1 4 4-13
- 97 Ingelman B and H Malmgren 1948 *Acta Chem Scand* 2 36 380
- 98 Ingelman B and H Malmgren 1949 *Acta Chem Scand* 3 15, 16
- 99 Ingraham J L and R Emerson 1954 *Am J Botany* 41 146-15
- 100 Jarvis P G and M J Johnson 1950 *J Bacteriol* 59 51-60
- 101 Javiller M 1913 *Ann inst Pasteur* 27 1021-1038
- 102 Javiller M 1939 *Ann fermentations* 5 371-381
- 103 Keilin D and A Tissot 1953 *Nature (London)* 17 393-394
- 104 Keitt G W and D M Boone 1954 *Phytopathology* 44 360-361
- 105 Kitasato T 1958a *Biochem Z* 197 237-258
- 106 Kitasato T 1958b *Biochem Z* 201 206-211
- 107 Kluyver A J, J P van der Walt and A J van Triet 1953 *Proc Natl Acad Sci U S A* 39 583-593
- 108 Koffler H, S G Knight and W C Frazier 1947 *J Bacteriol* 53 115-193
- 109 Krishnan P S 1951a *Arch Biochem* 3 30-251
- 110 Krishnan P S 1951b *Nature (London)* 168 171
- 111 Krishnan P S 1952 *Arch Biochem Biophys* 37 24-931
- 112 Krishnan P S and V Bajaj 1953a *Arch Biochem Biophys* 4 174-181
- 113 Krishnan P S and V Bajaj 1953b *Arch Biochem Biophys* 47 39-5
- 113a Lamprecht L 1957 *Arch Mikrobiol* 27 180-18
- 114 Lavollay J and F Laborey 1941 *Ann fermentations* 6 129-140
- 115 Lehninger A L 1950 *Physiol Revs* 30 393-419
- 116 Lewin S, Z P J Lucchesi and J E Vance 1953 *J Am Chem Soc* 75 60-8
- 117 Landeberg G 1944 *Symbolae Botan Upsalenses* 8 (2) 1-183
- 118 Linossier G 1910 *Compt rend (Paris)* 151 1075-1076
- 119 Lockwood L B and C E N Nelson 1946 *Arch Biochem* 10 365-374
- 120 Lockwood L B and M D Reeves 1945 *Arch Biochem* 6 455-469
- 121 Lohnis M P 1944 *Antonie van Leeuwenhoek J Microbiol Serol* 10 100-112
- 122 Lohrmann W 1940 *Arch Mikrobiol* 11 39-367
- 123 Lwoff S and E L Lumberg 1938 *Compt rend acad sc U R S S* 91 194-198
- 124 Lwoff A and H Ionesco 1948 *Ann inst Pasteur* 71 44-450
- 125 McCallan S E and F Wilcoxon 1931 *Contrib Boyce Thompson Inst* 3 13-38
- 126 McElroy W D and A Mason 1954 *Ann Rev Plant Physiol* 5 1-30
- 127 McHargue J S and R K Calfee 1931 *Botan Gaz* 91 183-193
- 128 Machlis L 1953 *A J Botany* 40 450-459
- 129 MacLeod R A and E E Snell 1948 *J Biol Chem* 176 39-52
- 130 MacLeod R A and E E Snell 1950 *Ann N Y Acad Sci* 52 1249-1250
- 131 McVeigh I and E Bell 1951 *Bull Torrey Botan Club* 78 134-144
- 132 Malmgren H 1950 *Acta Chem Scand* 6 16-6

- 133 Mann P J G and J H Quastel 1946 *Nature (London)* 158 154-156
- 134 Mann T 1944a *Biochem J (London)* 38 339-340
- 135 Mann T 1944b *Biochem J (London)* 38 345-351
- 136 Marsh I B 1910 *Phytopathology* 30 54-61
- 137 Marshall B H Jr 1935 *Phytopathology* 45 616-680
- 138 Marston H R 1902 *Physiol Revs* 32 66-121
- 139 Mehlich A E B Fred and E Truog 1934 *Soil Sci* 38 440-458
- 140 Mehlich A E B Fred and E Truog 1930 *J Am Soc Agron* 27 876-83
- 141 Menzel A E O O Wintersteiner and J C Hoogerheide 1914 *J Biol Chem* 152 419-479
- 142 Metz O 1930 *Arch Mikrobiol* 1 197-251
- 143 Michel Durand E 1938 *Bull soc chim biol* 20 399-412
- 144 Miller L P S E A McCallan and R M Weed 1903 *Contribs Boyce Thompson Inst* 17 151-171 173-195 283-298
- 145 Milkkan C R 1940 *J Australian Inst Agr Sci* 6 203-205
- 146 Molliard M 1900 *Compt rend (Paris)* 170 949-951
- 147 Mooers C A 1938 *Soil Sci* 46 211-227
- 148 Mothes K 1939 *Planta* 29 67-109
- 149 Mucke D 1904 *Flora* 141 30-50
- 150 Mulder E G 1939 *Arch Mikrobiol* 10 72-96
- 151 Mulder E G 1948a *Anal Chim Acta* 2 793-800
- 152 Mulder E G 1948b *Plant and Soil* 1 94-119
- 153 Mulder E G and F C Gerretsen 1909 *Advances in Agron* 4 909-977
- 154 Muntz J A 1947 *J Biol Chem* 171 653-660
- 155 Nason A 1906 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 109-136 Baltimore The Johns Hopkins Press
- 156 Nason A N O Kaplan and S P Colowick 1901 *J Biol Chem* 188 391-406
- 157 Nason A N O Kaplan and H A Oldewurtel 1903 *J Biol Chem* 901 435-441
- 158 Neillands J B 1907 *Bacteriol Revs* 73 101-111
- 159 Neuberg C and H A Fischer 1938a *Compt rend trav lab Carlsberg Sér Chim* 22 366-374
- 160 Neuberg C and H A Fischer 1938b *Enzymologia* 2 241-257
- 161 Neuberg C A Grauer and J Mandl 1900 *Enzymologia* 14 157-163
- 162 Neuberg C and K Kurono 1903 *Biochem Z* 140 295-298
- 163 Neuberg C and J Wagner 1905 *Biochem Z* 161 492-505
- 164 Neuberg C and J Wagner 1926a *Biochem Z* 171 485-500
- 165 Neuberg C and J Wagner 1926b *Biochem Z* 174 457-463
- 166 Newton W 1946 *Sci Agr* 26 303-304
- 167 Nicholas D J D 1900 *J Sci Food Agr* 1 339-344
- 168 Nicholas D J D 1902 *Analyst* 77 629-642
- 169 Nicholas D J D 1906 *J Gen Microbiol* 15 470-477
- 170 Nicholas D J D A Nason and W D McElroy 1954 *J Biol Chem* 201 341-344
- 171 Nicholas D J D and H M Stevens 1906 In W D McElroy and B Glass (eds) *Symposium on Inorganic Nitrogen Metabolism* p 178-183 Baltimore The Johns Hopkins Press
- 172 Nickerson W J 1901 *Trans N Y Acad Sci* 11 13 140-140
- 173 Nickerson W J 1953 *J Infectious Diseases* 93 43-56
- 174 Nickerson W J and J B Chadwick 1946 *Arch Biochem* 10 81-100
- 175 Norkrans B 1900 *Symbolae Botan Upsalienses* 11(1) 1-1.6
- 176 Orskov S L 1918 *Acta Pathol Microbiol Scand* 20 277-283
- 177 Osharov K E 1937 *Compt rend acad sci U R S S* 16 461-464 (*Chem Abstr* 32 1740 1938)
- 178 Painter H A 1904 *J Gen Microbiol* 10 177-190

- 179 Pathak M A and A Sreenivasan 1955 *Arch Biochem Biophys* 59 306-312
- 180 Perkins D D 1949 *Genetics* 34 601-606
- 181 Perlman D 1948 *Am J Botany* 35 36-41
- 182 Perlman D 1949a *Am J Botany* 36 180-184
- 183 Perlman D 1949b *Botan Per* 15 19-290
- 184 Perlman D 1951 *Am J Botany* 38 6 2-6 8
- 185 Perlman D W W Dorrell and M J Johnson 1946 *Arch Biochem* 11 130-143
- 186 Pine L 1954 *J Bacteriol* 68 671-679
- 187 Pisano M A B H Olson and C L San Clemente 1954 *J Bacteriol* 68 444-449
- 188 Pontecorvo G *et al* 1953 *Advances in Genet* 5 141-238
- 189 Porges N 193 *Botan Gaz* 94 197-205
- 190 Purdy L H Jr and R G Grogan 1954 *Phytopathology* 44 26-39
- 191 Quilico A and A Di Capua 1933 *Atti accad naz Lincei Ser 6* 17 177-182
- 192 Raistrick H and J M Vincent 1948 *Biochem J (London)* 43 90-99
- 193 Rautanen N and A E Kyla Siurola 1954 *Acta Chem Scand* 8 106-111
- 194 Reischer H S 1951 *Mycologia* 43 142-155
- 195 Rennerfelt E 1934 *Planta* 29 1-239
- 196 Rippel A and G Behr 1934 *Arch Mikrobiol* 5 561-577
- 197 Rippel A and G Behr 1936 *Arch Mikrobiol* 7 584-589
- 198 Rippel A and U Stoess 1932 *Arch Mikrobiol* 3 492-506
- 199 Robbins W J and A. Hervey 1944 *Bull Torrey Botan Club* 71 253-266
- 200 Robert M 1928 *Centr Bakteriell Pa astenkl Abt II* 74 333-370
- 201 Roberts C 1946 *Am J Botany* 33 237-244
- 202 Roberts R B I Z Roberts and D B Cowie 1949 *J Cellular Comp Physiol* 34 259-292
- 203 Robertson H E and P D Boyer 1956 *Arch Biochem Biophys* 62 380-395 396-401
- 204 Roche J 1950 In J B Sumner and K Myrback (eds) *The Enzymes* Vol 1 (1) p 473-510 New York Academic Press
- 205 Rothstein A and R Meier 1949 *J Cellular Comp Physiol* 34 97-114
- 206 Sadasivan V 1950 *Arch Biochem* 28 100-110
- 207 Sadasivan V 1952 *Arch Biochem Biophys* 37 17-185
- 208 Sakseena R K S K Jagan and S M H Jafri 1953 *J Indian Botan Soc.* 31 81-86
- 209 Salvin S B 1949 *J Infectious Diseases* 84 27-283
- 209a Sanwal B D and R S Sandhu 1956 *Experientia* 12 380-381
- 210 Sarasin A 1953 *Ber schweiz botan Ges* 63 287-316
- 211 Sauton B 1910 *Compt Rend (Paris)* 151 241-243
- 212 Schade A I 1949 *J Bacteriol* 58 811-819
- 213 Schatz A R R Mohan and G S Trelawny 1955 *Antonie van Leeuwenhoek J Microbiol Serol* 1 225-238
- 214 Schulz G 1937 *Planta* 27 196-218
- 215 Schwartz W and H Steinhart 1933 *Arch Mikrobiol* 4 301-325
- 216 Scianini L J and F F Nord 1943 *Arch Biochem* 3 261-267
- 217 Semenuk C 1944 *Iowa State Coll J Sci* 18 325-338
- 218 Sharp E L and F C Smith 1955 *Phytopathology* 42 581-582
- 219 Shepherd C J 1956 *J Gen Microbiol* 15 29-38
- 220 Shibata C 1957 *Centr Bakteriell Pa astenkl Abt II* 71 232-247
- 221 Shu P and M J Johnson 1948 *J Bacteriol* 56 577-585
- 222 Simonart P and K Y Chow 1954 *Antonie van Leeuwenhoek J Microbiol Serol* 20 210-216
- 223 Smith V M 1949 *Arch Biochem* 23 416-470
- 224 Stahl W H B McQue G R Mandels and R G H Siu 1949 *Arch Biochem* 4 430

- 225 Starkey R L 1935 *Soil Sci* 19 1-14
- 226 Starkey R L and S A Waksman 1945 *J Bacteriol* 45 509-519
- 227 Steinberg R A 1935 *Bull Torrey Botan Club* 62 81-90
- 228 Steinberg R A 1936a *Botan Gaz* 97 666-671
- 229 Steinberg R A 1936b *J Agr Research* 52 439-448
- 230 Steinberg R A 1937 *J Agr Research* 55 891-902
- 231 Steinberg R A 1938 *J Agr Research* 57 569-574
- 232 Steinberg R A 1939a *J Agr Research* 59 731-748
- 233 Steinberg R A 1939b *J Agr Research* 59 749-764
- 234 Steinberg R A 1941 *J Agr Research* 63 109-127
- 235 Steinberg R A 1942 *J Agr Research* 64 435-475
- 236 Steinberg R A 1943a *Plant Physiol* 20 600-608
- 237 Steinberg R A 1943b *Soil Sci* 60 185-189
- 238 Steinberg R A 1946 *Am J Botany* 33 210-214
- 239 Steinberg R A 1948 *Science* 107 423
- 240 Steinberg R A 1950 *Arch Biochem* 28 111-116
- 241 Steinberg R A and J D Bowling 1939 *J Agr Research* 58 717-737
- 242 Strauss B S 1953 *Arch Biochem Biophys* 44 200-210
- 243 Sussman A S 1954 *J Gen Physiol* 38 59-77
- 244 Tandon M P 1950 *Proc Indian Acad Sci B* 32 7-11
- 245 Texera D 1948 *Phytopathology* 38 70-81
- 246 Treschow C 1944 *Dansk Botan Arkiv* 11 (6) 1-180
- 247 Tsuyuki H and R A MacLeod 1951 *J Biol Chem* 190 711-719
- 248 Turian G 1956 *Experientia* 12 24-28
- 249 Vallee B L S J Adelstein and J A Olson 1955 *J Am Chem Soc* 77 5196
- 250 Vallee B L and F L Hoch 1955 *Proc Natl Acad Sci U S A* 41 377-384
- 251 Varma T N R and K S Srinivasan 1953 *Enzymologia* 17 116-127
- 252 Vishniac H S 1955 *J Gen Microbiol* 12 455-463
- 253 Waksman S A 1953 *Neomycin* New Brunswick Rutgers Univ Press pp 219
- 254 Whiffen A J 1945 *J Florida Mitchell Sci Soc* 61 114-123
- 255 Winfield M E 1949 *Australian J Exp Biol Med Sci* 23 267-272
- 256 Yogeswari L 1949 *Proc Indian Acad Sci B* 28 177-201
- 257 Yoshimura F 1939 *Botan Mag (Tokyo)* 53 125-138
- 258 Young H C and C W Bennett 1929 *Am J Botany* 9 459-469
- 259 Zalokar M 1953 *Arch Biochem Biophys* 44 330-337

10. Vitamin Requirements of Fungi

Fungi like all other organisms so far known require minute amounts of specific organic compounds for growth. The cell may synthesize its own supply of one of these growth factors or it may be dependent in whole or in part on an exogenous supply. Commonly we refer to the dependence on exogenous supply as a requirement realizing that the organism which synthesizes its own growth factors is equally dependent on them for normal metabolism.

The definition of growth factors may be broad or narrow. We may for example include as growth substances all compounds required in small amounts and not used for energy (260). This broader definition includes amino acids, purines and choline as growth factors. A narrower concept the one used here excludes compounds that function as structural materials even though they are needed in small amounts.

Most of the known vitamins have a catalytic function in the cell as coenzymes or constituent parts of coenzymes. This function is stressed in the definition of vitamins employed in this chapter: organic molecules required in small amounts and not so far as we know used as sources of either energy or structural materials of protoplasm. This definition is arbitrary and provisional. Arbitrary because it separates constituents of coenzymes (e.g. riboflavin) from constituents of enzymes (e.g. amino acids) and cannot separate for example the role of adenine in adenosine triphosphate from its role in the structure of nucleic acids. Provisional because the cofactor role of some of the vitamins especially inositol is uncertain.

This chapter then includes substances primarily the water soluble vitamins which are thought to be catalytic in function the amino acids and nitrogen bases are considered in Chapter 8 as is the plant growth factor indoleacetic acid

The history of our knowledge of the vitamin requirements of fungi is reviewed by Janke (117) Robbins and V Kavanagh (213) Schopfer (239) and Knight (122) Although the work on yeasts and yeastlike fungi really began in the nineteenth century the vitamin needs of the filamentous fungi have been studied intensively only since 1934 i.e. since the papers of Schopfer (236 237) on thiamine and the growth of *Phycomyces blakesleeianus* Most work has been devoted to the requirements of naturally occurring fungi and more recently to the requirements of induced mutants both types of material have provided significant information on the biosynthesis and the function of vitamins in living things generally

1 PROBLEMS OF METHOD

The central methodological problem in vitamin studies is that of the degree to which the investigator knows the composition of the medium Almost any natural product e.g. sucrose or asparagine is contaminated with significant amounts of some vitamins procedures for purification will be found in the literature on particular growth factors

For very limited purposes e.g. in formulating a medium for routine use it may be sufficient to use the approximate methods so common in the literature on vitamin nutrition However a serious study of vitamin nutrition requires at the least that three methodological criteria be met

- 1 The measurement of growth should be objective and quantitative Although growth on agar has been used fairly successfully in particular problems it cannot be relied upon and dry weight or turbidity is to be preferred (Chapter 1) Pyridine extracted agar (214) may however be used for some problems

- 2 Over some range of concentration of the vitamin it should be shown that growth is a function of dose (concentration or total amount of the vitamin)

- 3 If growth occurs on the basal medium lacking added vitamin an independent assay of the medium for the vitamin should be made One very simple test consists of growing on the basal medium an organism known to require the factor absolutely Other problems of method are summarized by Robbins and V Kavanagh (213)

Especially with negative results i.e. the finding that an organism does not need a vitamin still further precautions may be mentioned. First the basal medium must be free of the vitamin even cotton plugs may contribute growth factors (248-251). Second it is essential that the organism be serially subcultured in the vitamin free medium for at least three transfers. In some investigations (131) the inoculum has no effect but this cannot be assumed. The criterion of serial subculture is the one most frequently disregarded and in consequence most reports of a lack of need for a vitamin—especially those e.g. biotin for which the requirement is always small—are not wholly reliable.

The use of crude extracts of natural products in the preliminary search for vitamin requirements is of course helpful and within limits valuable. However the simple observation that such an extract increases growth has no necessary bearing on vitamin nutrition since so many other materials known and unknown are present in these extracts. Thus the effect of agar on growth of several fungi (132) and the effect of yeast extract on cytochrome synthesis by *Ustilago sphaerogena* (89) are both attributable to the zinc content of the natural product not at all to organic materials.

The vitamin content of many common constituents of culture media is known (34-44-95-206-217-224-264) occurrence in other natural products is reviewed in general works on the vitamins (225-249).

2 THE NATURE OF VITAMIN REQUIREMENTS

It is generally assumed that a fungus or other organism which is independent of an externally supplied vitamin is so by virtue of its ability to synthesize the compound—a thiamine independent fungus synthesizes thiamine etc. The validity of this assumption is borne out by experimental evidence in all cases so far investigated and is logically deduced from the premise that the known vitamins have essential functions in all cells. This premise it should be noted extends only to the water soluble or B vitamins.

Further evidence if such is needed for the essentiality of vitamins in those fungi which do not require an external supply is found in the fact that deficient mutants can usually be obtained by appropriate techniques examples in addition to the classical *Neurospora crassa* include species of *Penicillium* (17) and *Aspergillus* (115-195-199) genera in which naturally occurring vitamin deficiencies are uncommon.

The simplest situation in vitamin studies is that of complete inability of a given fungus to synthesize one or more vitamins. In this circumstance the cell depends absolutely on an external supply and at

This chapter then includes substances primarily the water soluble vitamins which are thought to be catalytic in function the amino acids and nitrogen bases are considered in Chapter 8 as is the plant growth factor indoleacetic acid

The history of our knowledge of the vitamin requirements of fungi is reviewed by Jinke (117) Robbins and V Kavanagh (213) Schopfer (239) and Knight (122) Although the work on yeasts and yeastlike fungi really began in the nineteenth century the vitamin needs of the filamentous fungi have been studied intensively only since 1931 i.e. since the papers of Schopfer (236-237) on thiamine and the growth of *Phycomyces blakesleeana* Most work has been devoted to the requirements of naturally occurring fungi and more recently to the requirements of induced mutants both types of material have provided significant information on the biosynthesis and the function of vitamins in living things generally

1 PROBLEMS OF METHOD

The central methodological problem in vitamin studies is that of the degree to which the investigator knows the composition of the medium Almost any natural product e.g. sucrose or asparagine is contaminated with significant amounts of some vitamins procedures for purification will be found in the literature on particular growth factors

For very limited purposes e.g. in formulating a medium for routine use it may be sufficient to use the approximate methods so common in the literature on vitamin nutrition However a serious study of vitamin nutrition requires at the least that three methodological criteria be met

1 The measurement of growth should be objective and quantitative Although growth on agar has been used fairly successfully in particular problems it cannot be relied upon and dry weight or turbidity is to be preferred (Chapter 1) Pyridine-extracted agar (214) may however be used for some problems

2 Over some range of concentration of the vitamin it should be shown that growth is a function of dose (concentration or total amount of the vitamin)

3 If growth occurs on the basal medium lacking added vitamin an independent assay of the medium for the vitamin should be made One very simple test consists of growing on the basal medium an organism known to require the factor absolutely Other problems of method are summarized by Robbins and V Kavanagh (213)

which the vitamin is essential will obviously reduce the demand for the exogenous vitamin. The replacement of nicotinic acid by large amounts of tryptophan represents replacement by a precursor (20). The replacement of *p* aminobenzoic acid by methionine (269) may be interpreted as evidence that at least one function of the vitamin is in the biosynthesis of methionine and a similar mechanism is invoked to explain the partial replacement of biotin by aspartic acid (190).

The third type of conditioned requirement is the most interesting and the least studied—the possibility that fungi change in their synthetic capacity and in their vitamin needs during development. It must be realized that the typical vitamin experiment performed with fungi actually measures in the first instance the capacity of the spores to germinate in the absence of the vitamin in question: if the spore germination process requires the growth factor the organism as such is recorded as having the requirement. It is entirely possible that spore germination may require factors which mature mycelium can synthesize for itself. Thus the growth of *Memnoniella echinata* from a spore inoculum requires biotin which is not replaceable by desthiobiotin. However, if desthiobiotin is added to a growing culture supplied with limiting amounts of biotin the desthiobiotin elicits a significant growth response (191). Evidently the mycelium is able to convert desthiobiotin to biotin but the spore cannot. Successful use of the differential germination method for the detection of vitamin-deficient mutants (69–308) shows that spore germination is affected by growth factor deficiencies.

Several fungi—*Alternaria solani* (138) *Fusarium solani* (285) and *Penicillium digitatum* (61)—are reported to be accelerated in early growth by vitamins the requirements for which are not evident if total growth over a long period is measured. In *Myrothecium verrucaria* a requirement for biotin is detectable only at a stage shortly after spore germination (157).

Among the dimorphic fungi (Chapter 1) a single study (229) suggests that the yeast and the mycelial phases of some forms differ in their biotin requirement.

The onset of reproduction as a phase of growth may, although the data are far from decisive, involve changes in vitamin needs and responses. There is however no evidence of a singular association of one vitamin with sporulation *per se* independent of growth. This problem is considered in Chapter 11.

Multiple requirements are especially common in the yeasts: five or six vitamins may be needed (27–29–39). Among the filamentous fungi there are several instances of a requirement for three different vitamins

least over a certain range growth will be proportional to the supply of the required factor or factors. Many natural and induced deficiencies are or appear to be of this type which is the most easily detectable.

A somewhat more complicated situation arises if a fungus is able to synthesize a vitamin, but so slowly that under the usual conditions of culture the rate of all other processes is potentially faster than vitamin synthesis. Here we find the organism growing slowly in the absence of exogenous vitamin but responding to an external supply by a faster rate of growth. Such partial deficiencies are also common perhaps more common than complete deficiencies. Examples include the requirements of particular fungi for thiamine (16, 65, 183, 186), biotin (144, 216), pyridoxine (66, 67, 216, 217), nicotinic acid (291) and inositol (238).

It also follows that any growth regulating process—including vitamin synthesis—may be limiting under some conditions and not under others. A vitamin requirement that is manifest only under certain circumstances is termed a *conditioned requirement*. The known types of conditioned deficiency in fungi may be classified as follows:

1. The deficiency is apparent or more severe at particular temperatures, pH levels, or salt concentrations.
2. The requirement for a vitamin is reduced or more rarely eliminated by provision in the medium of a precursor or of a metabolite for the synthesis of which the vitamin is essential.
3. The deficiency is limited to or more acute at a particular stage of development.

Environmental factors conditioning vitamin deficiencies include temperature, acidity, and salt concentration. A mutant of *Neurospora crassa* for example requires riboflavin only at relatively high temperatures (172). *Sclerotinia camelliae* requires more inositol at 26° C than at 18° but is inhibited by inositol at 27° (8). A temperature-dependent inositolless mutant of *N. crassa* is also known (113). *Sordaria fimicola* requires thiamine only if the initial pH of the culture is less than 4.0, possibly because thiamine synthesis is inhibited by hydrogen ion (139). A similar phenomenon in mutants of *Neurospora* spp. which require pyridoxine appears, however, to reflect a requirement for free ammonia rather than a direct effect of acid (268). The influence of pH on the *p*-aminobenzoic acid requirement seems finally to be exerted through permeability effects (325). *Pythium butleri* requires exogenous thiamine only in a high salt medium (211).

The second type of conditioned requirement is more easily explained: a precursor of the vitamin or a metabolite in the synthesis of

thiamine biotin pyridoxine *p*-aminobenzoic acid pantothenic acid and inositol (239 259 281). Usually dry weight is the criterion although pyruvic acid formation (93) pigment production (289) and rate of growth on agar (10 277 282) have all been proposed.

Assays based on the dry weight of vitamin requiring fungi have in general the advantages and disadvantages of any biological assay (171 259). However the relatively more tedious procedures necessary with the fungi have persuaded most investigators to use whenever possible bioassays based on yeast or bacterial growth or metabolism. In special cases fungi may prove more valuable thus different fungi able to use either the pyrimidine or the thiazole moiety of thiamine were used by Krampitz and Woolley (121) in a study of the thiaminase of carp.

Bioassays using fungi are critically reviewed by Snell (259) others have been described for biotin (221) pantothenate (290) and thiamine (93 289). The *Phycomyces* assay for thiamine is reviewed by Ødegård (185). A survey of the response of many fungi to graded doses of water soluble vitamins (136) brings out some of the difficulties inherent in biological assay procedures.

5 THIAMINE

Thiamine (aneurin vitamin B₁) is the first vitamin to have been studied as a known entity in the nutrition of fungi and is the vitamin most frequently required by fungi. Thiamine deficiencies occur in all major taxonomic groups—although the ascomycetes have been less thoroughly investigated than others—and in fungi from many different natural environments.

Reported requirements vary widely (65 226 300) time of harvest temperature and the composition of the medium can be expected to affect thiamine demand. It seems at least possible that those fungi with very small requirements are able to synthesize thiamine once growth has been initiated. A typical response curve is shown in Figure 1.

The metabolically active form of thiamine is the pyrophosphate long known as cocarboxylase because of its coenzyme function in the decarboxylation of pyruvic acid. Thiamine pyrophosphate is also the coenzyme for transketolase and for a variety of enzymatic reactions of α keto acids (118). In the fungi its role in the metabolism of pyruvic acid is clear from its acceleration of ethanol formation (42 217) and the accumulation of pyruvate in thiamine-deficient cultures (64 93 301). Acceleration of glucose utilization (101) and decrease in oxalate accumulation (175) may operate through the same mechanism.

e.g. in *Ascoidea rubescens* (67) *Blastocladia pringsheimii* (35) *Ceratostomella microspora* (216) and *Trichophyton discoides* strains (159 222)

Inhibitory effects of vitamins have been reported occasionally. Thiamine depresses the growth of *Fusarium* spp. (55 285 304) and *Rhizopus* spp. (142 210 238) biotin may reduce growth or enzyme formation (11 228 253 316). Large amounts of pyridoxine do not however appear to depress growth (67). The mechanism of inhibition is not always known: thiamine inhibition has been attributed to ammonia accumulation in *Trichoderma viride* (16) and to ethanol accumulation in *Rhizopus stolonatus* (247).

3 THE METABOLISM OF VITAMINS

Apart from their catalytic role in metabolism, vitamins are themselves metabolites and are both synthesized and destroyed in the cell. More or less comprehensive surveys of vitamin synthesis have been reported for *Penicillium* spp. (37 100 178 274) and a good deal of information is available on vitamin synthesis by *Aspergillus* spp. (37 60 85 127 181 233 262) and *Streptomyces* spp. (47 100 103 105 106 151). The vitamin content of microorganisms is reviewed comprehensively by Van Lanen and Tanner (288). As mentioned earlier, we may assume that a fungus which does not require one of the water-soluble vitamins known to be required by other fungi is able to synthesize it.

Vitamins synthesized by a fungus usually appear in the medium and are detected there by bioassay; however, in *Penicillium chrysogenum* the greater part of the thiamine, riboflavin and nicotinic acid produced remains in the cell (100) and the same is true of thiamine in some other fungi (133) and of riboflavin in *Aspergillus niger* (60).

The destruction of vitamins by fungi has been less thoroughly studied. *Phycomyces blakesleeanus* and *Sclerotium rolfsii* destroy the thiazole moiety of thiamine (21 120). A mutant of *Neurospora crassa* destroys pyridoxine and this destruction is inhibited competitively by thiamine (96).

4 FUNGI IN THE BIOASSAY OF VITAMINS

Analysis for individual vitamins by a biological assay is often, for one reason or another, superior to chemical analysis. This is especially true if a biologically active vitamin exists in more than one chemical form. Bioassays employing filamentous fungi have been devised for

Table 1 Response of Fungi to Thiamine and Its Moieties*

Minimum Requirement	Examples
Intact thiamine	<i>Ceratostomella fimbriata</i> (220) <i>Chalaropsis thielavii</i> (220) <i>Cortinarius glaucopus</i> (167) <i>Phytophthora</i> spp (188 204) <i>Tilletia tritici</i> (46) <i>Trichophyton discoides</i> (222)
Both moieties	<i>Boletus variegatus</i> (167) <i>Ceratostomella</i> spp (220) <i>Marasmius perforans</i> (145) <i>Phycomyces</i> spp (130 205 210) <i>Piricularia oryzae</i> (284) <i>Polyporus versicolor</i> (220) <i>Tilletia horrida</i> (245) <i>Ustilago</i> spp (15)
Thiazole	<i>Collybia velutipes</i> † (156) <i>Endomyces magnusii</i> (165) <i>Mucor ramanianus</i> (173 174) <i>Stereum frustulosum</i> (182)
Pyrimidine	<i>Allomyces kniepi</i> (196) <i>Blakeslea trispora</i> (130) <i>Blastocladiella emersonii</i> (7) <i>Ceratostomella</i> spp (220) <i>Cercospora herpotrichoides</i> (45) <i>Collybia tuberosa</i> (130) <i>Coprinus lagopus</i> (130) <i>Fomes</i> spp (332) <i>Hypholoma fasciculare</i> (245) <i>Marasmius fulvobulbosus</i> (145) <i>Nocardia</i> spp (157) <i>Polyporus</i> spp (332) <i>Sclerotium</i> spp (205) <i>Sphaerulina trifolii</i> (210) <i>Stereum murranii</i> (208) <i>Trichobolus tonsurans</i> (273)

* A more inclusive list is given by Robbins and V. Kavanagh (213)

† Not definitely established

to live with either moiety alone does in fact make the other and synthesize thiamine (14 21 133 220 239) neither pyrimidine nor thiazole by itself has a vitamin function

A great deal of attention has been devoted to the specificity of the structures of thiazole and pyrimidine for fungi. In general the requirements are quite specific (213 225 239) most utilizable modifications e.g. the substitution of an acetyl for a hydroxyl group or of an ethyl for a methyl group are either minor or are very likely transformed biologically into the natural product. Allithiamine fully replaces thiamine in the nutrition of *Endoconidiophora fimbriata* *Mucor ramanianus* and other fungi and can be converted into thiamine in the cell (141)

It might be assumed *a priori* that biosynthesis of thiamine proceeds via a coupling of the two moieties. However there are some difficulties with this simple concept as applied to *Neurospora crassa* and Harris (97 98) on the basis of indirect evidence from nutritional studies has proposed that thiamine biosynthesis proceeds in this fashion

Pyrimidine + unidentified thiazole precursor →

thiamine-like intermediate → thiamine (1)

Direct coupling of pyrimidine and thiazole may occur as a minor pathway

6 BIOTIN

Biotin (vitamin H coenzyme R) is required by many yeasts and fungi; all the major taxonomic groups of the fungi are represented by biotin requiring types (5 35 67 128 161 214 219 240 241). The absolute requirement is usually less than 5 μg per liter (Figure 2).


The role of biotin is not certain in bacteria; it or a derivative of it is probably involved in several enzymatic systems (126). Evidence from studies of several fungi (111 155 190) implicates biotin in the synthesis of aspartic acid inasmuch as aspartate partially replaces biotin in nutrition. It seems likely that the reaction affected by biotin is either the carboxylation of pyruvic acid to yield oxalacetic acid or the conversion of oxalacetate to aspartate (92).




A second probable function of biotin in fungi is in the synthesis of essential fatty acids; oleic acid partially replaces biotin for a *Neurospora crassa* mutant (111) and for *Ophiostoma pini* (163). It must be said that evidence for the essentiality of fatty acids for fungi is incomplete; the possibility is raised by (a) the stimulating effect of oleic acid on growth (12 84 332) and (b) the occurrence in *Neurospora crassa* of mutants requiring fatty acids (129).

Less well-defined and probably indirect effects of biotin on fungi include inhibition of growth of a pantothenicless mutant (228), inhibition of enzyme synthesis in one strain of *Neurospora crassa* (11) and change in the growth pattern of the dimorphic *Sporotrichum schenckii* (160). *Mitrella paludosa* requires biotin only if ammonium nitrogen is used in the medium (67).

In relation to biosynthesis the structures of three compounds are of interest:



Desethyl biotin



We may assume that synthesis of the pyrimidine moiety follows that of other pyrimidines (Chapter 8) The origin of thiazole in *Neurospora crassa* is apparently related to amino acid metabolism (50)

Two fungi *Phycomyces blakesleeanus* and *Sclerotium rolfsii* which can use the two moieties in place of thiamine also destroy or convert to an inactive form part of the thiazole whether added as thiamine or as free thiazole *Phytophthora cinnamomi* which requires intact thiamine does not destroy free or combined thiazole Thus the destruction of thiazole seems to be unrelated to the use of thiamine in the growth processes of the fungi (120) The effect of temperature on the enzyme which destroys thiazole may explain the observation that the efficiency of thiamine for growth increases with decreasing temperature (212)

The compound pyrithiamine is inhibitory to those fungi e.g. *Phytophthora cinnamomi* which require intact thiamine and does not inhibit those e.g. *Neurospora crassa* which synthesize thiamine organisms requiring only one moiety of thiamine are intermediate in susceptibility (312-314) However the problem needs further study in view of the fact that thiamineless mutants of *Neurospora crassa* do not respond according to expectations (279) Further it is believed that pyrithiamine is a mixture of different compounds neopyrithiamine has been synthesized (303) and has the structure of a pyridine analogue of thiamine originally assigned to pyrithiamine. Neopyrithiamine competitively inhibits thiamine metabolism in animals (162) Pyrithiamine can be used by pyrimidine-deficient fungi as a source of the pyrimidine ring (207)




Figure 2 The response of *Stachybotrys* sp (SV 1) to biotin From data of Marsh and Bollenbacher (161)

biotin *dl* Oxybiotin (O heterobiotin) has slight activity for fungi (190) as it does for other organisms (92). Specific anti biotin activity against fungi is reported for avidin (317) biotin sulfone (321) and γ (3,4 ureylenecyclohexyl)butyric acid (191).

7 PYRIDOXINE

Pyridoxine (vitamin B₆, adermin) appears to be required by fewer fungi than is biotin or thiamine. Several species of *Ceratostomella* (*Ophiostoma*) require it (67, 215) as do *Ascoidea rubescens* and *Trichophyton discoides* (67, 215, 222). The requirement is highly species specific within the genus *Ceratostomella*.


Pyridoxineless mutants of *Neurospora crassa* and *N. sitophila* grow better at limiting pyridoxine concentration if thiamine is present (263, 279). This relation between two vitamins has been explained by the postulate that thiamine competitively inhibits the endogenous destruction of pyridoxine (96). Conversely pyridoxine inhibits the biosynthesis of thiamine probably by preventing the incorporation of the pyrimidine moiety (98).

As mentioned above certain mutants of *Neurospora crassa* require pyridoxine only at low pH. The mechanism of this conditioned deficiency is a requirement for free ammonia in the absence of pyridoxine at low pH. Insufficient free ammonia is present and the vitamin is therefore needed for growth (268).

The coenzyme form of pyridoxine is pyridoxal 5 phosphate known to be the coenzyme for enzymes of amino acid metabolism specifically decarboxylases, transaminases and racemases (286). Insofar as these enzymes are present in fungi (Chapter 8) we may assume the involvement of the coenzyme also. In addition pyridoxal phosphate has been specifically shown to be active in several isolated enzyme systems of *Neurospora crassa* (201, 268, 287, 327). Pyridoxine deficiency in *N. crassa* mutants is accompanied by accumulation of nitrite from nitrate indicating a role of pyridoxal phosphate in nitrite reduction (255).

Although pyridoxine, pyridoxamine and pyridoxal are equally usable by fungi (261) other modifications of the basic molecule are usually inactive or toxic (218); the few that are active can easily be guessed to be converted in the cell to pyridoxine (198, 218).

The growth of a pyridoxineless mutant of *Neurospora sitophila* is inhibited by 4 deoxypyridine; the inhibition is competitively reversed by pyridoxine and its derivatives (197, 198).



Tatum (276) studied the nutrition of biotinless mutants of *Penicillium chrysogenum*. One mutant can use desthiobiotin as well as biotin; a second mutant accumulates desthiobiotin but cannot use it. Production of desthiobiotin in this strain is increased by provision of pimelic acid [$\text{HOOC} \text{---} (\text{CH}_2)_6 \text{---} \text{COOH}$]. Equation 3 therefore summarizes the known steps in biosynthesis:




Biotin deficient fungi isolated from nature fall into two groups. The first, exemplified by *Ceratostomella ips*, *Neurospora crassa*, *Blastomyces dermatitidis* and yeast (49, 95, 143), can grow at least to a degree with desthiobiotin in place of biotin; these fungi are presumably blocked in a step prior to desthiobiotin synthesis. The second group, which includes *Ceratostomella pini* and *Sordaria fimicola*, cannot use desthiobiotin (135, 143) and is therefore blocked in the conversion of desthiobiotin to biotin. Those fungi known only to be unable to use pimelic acid in place of biotin (219) cannot yet be placed in either group. We have noted earlier the different responses of spores and pregrown cultures of *Memnoniella echinata* to desthiobiotin.

Biotin sulfoxide is formed by *Aspergillus niger* in synthetic medium since pimelic acid and desthiobiotin increase the yield, and since added biotin is quantitatively converted to the sulfoxide, it is concluded that *A. niger* synthesizes biotin as in Equation 3 and then transforms it to biotin sulfoxide (54, 317, 320, 321, 322). Biotin sulfoxide is as active as biotin for *Neurospora crassa* but not for yeast or the bacteria so far tested (321). *Aspergillus niger* utilizes azelaic acid also in biotin synthesis (318).

Biocytin, ϵ -N-biotinyl-L-lysine, a naturally occurring form of biotin, is as active as biotin for *Neurospora crassa* and for a biotin requiring mutant of *Penicillium chrysogenum* (319). The oxygen analogue of

gossypii and *Eremothecium ashbyii* are less sensitive to iron (75 301). Moderate levels of glucose, a slightly acid pH, and aeration all improve the yield of riboflavin (202 243 301). Riboflavin formation by *Aspergillus niger* is favored by any of a variety of nutritional factors, all of which somewhat retard growth (230).

The structure of riboflavin is



Provision of certain purines, e.g., adenine or guanine, to *Eremothecium ashbyii* increases the yield of riboflavin (26 153 155). The entry of isotopic formate, carbon dioxide, and acetate (194) and of labeled adenine (163) into the molecule also suggests that the synthesis of ring C is similar to the synthesis of the pyrimidine ring of a purine or that the pyrimidine ring of a purine is incorporated intact (156).

Inhibition studies with 1,2-dichloro-4,5-diaminobenzene led Woolley (310 311) to suggest that the natural precursor of ring A—and of cyanocobalamin (p. 343)—is 1,2-dimethyl-4,5-diaminobenzene.

The cellular function of riboflavin is to provide the prosthetic group of a number of oxidizing enzymes, the flavin enzymes (114). Flavin enzymes known in fungi include glucose aerodehydrogenase and other respiratory enzymes (Chapter 7) and several enzymes of nitrogen metabolism—nitrate reductase, nitrite reductase, hydroxylamine reductase, and the amino acid oxidases (Chapter 8). Evidence from organisms other than the fungi indicates that riboflavin plays a role in the synthesis of nicotinic acid (41).

Zalokar (335) has suggested that a riboflavin-protein complex may be the photoreceptor in the light-activated synthesis of carotene by *Neurospora crassa* (Chapter 6).

9 NICOTINIC ACID

Deficiencies for nicotinic acid—or its amide, which appears from limited evidence to be physiologically equivalent to the free acid in fungi—occur in relatively few fungi. *Microsporium audouinii* (5) *Blas*

8 RIBOFLAVIN

Riboflavin (vitamin B₂ lactoflavin) is only rarely required by microorganisms isolated from nature. Several of the lactic acid bacteria are riboflavin deficient (192) the yeasts appear not to require it (29). A single report without data indicates that *Poria vaillantii* requires riboflavin (119) this is the first such claim for a filamentous fungus. *Diclyostelium* spp. have an absolute requirement (271).

However riboflavinless mutants can usually be isolated by appropriate techniques (23 77 115 121 195). Of these the most interesting is the mutant of *Neurospora crassa* isolated by Mitchell and Houlahan (172). This strain does not require exogenous riboflavin at moderate temperatures but only if grown at 28° C or higher (Figure 3). Even at the elevated temperature the mutant synthesizes riboflavin if it is provided initially with some of the vitamin.

Riboflavin has been shown directly to be formed by many yeasts and fungi (60 91 100 137 172 189 288 302 335). In two yeastlike fungi *Eremothecium ashbyii* and *Ashbya gossypii* yields are high enough so that commercial use is practicable (75) it is believed that up to 6 mg per milliliter of free riboflavin is produced by *A. gossypii* in a complex medium. Although esters of riboflavin especially flavin adenine dinucleotide occur in the cell only free riboflavin is found in the medium (252 326).

The physiological conditions of riboflavin synthesis in the yeastlike fungi have been of both practical and theoretical interest. Although even relatively small amounts of iron inhibit riboflavin synthesis by *Clostridium acetobutylicum* (109) and *Candida* spp. (275) *Ashbya*





Figure 3 The response of a *Neurospora crassa* mutant to exogenous riboflavin. Curve 1 at 30° C curve 2 at 25°. Redrawn from Nicholas (180) by permission of the *Journal of General Microbiology*.

gossypii and *Eremothecium ashbyii* are less sensitive to iron (75 301) Moderate levels of glucose a slightly acid pH and aeration all improve the yield of riboflavin (202 243 301) Riboflavin formation by *Aspergillus niger* is favored by any of a variety of nutritional factors all of which somewhat retard growth (230)

The structure of riboflavin is



Provision of certain purines e.g. adenine or guanine to *Eremothecium ashbyii* increases the yield of riboflavin (26 153 155) The entry of isotopic formate carbon dioxide and acetate (194) and of labeled adenine (163) into the molecule also suggests that the synthesis of ring C is similar to the synthesis of the pyrimidine ring of a purine or that the pyrimidine ring of a purine is incorporated intact (156)

Inhibition studies with 1,2-dichloro-4,5-diaminobenzene led Woolley (310 311) to suggest that the natural precursor of ring A—and of cyanocobalamin (p. 343)—is 1,2-dimethyl-4,5-diaminobenzene

The cellular function of riboflavin is to provide the prosthetic group of a number of oxidizing enzymes the flavin enzymes (114) Flavin enzymes known in fungi include glucose acrodehydrogenase and other respiratory enzymes (Chapter 7) and several enzymes of nitrogen metabolism—nitrate reductase nitrite reductase hydroxylamine reductase and the amino acid oxidases (Chapter 8) Evidence from organisms other than the fungi indicates that riboflavin plays a role in the synthesis of nicotinic acid (41)

Zalokar (335) has suggested that a riboflavin protein complex may be the photoreceptor in the light activated synthesis of carotene by *Neurospora crassa* (Chapter 6)

9 NICOTINIC ACID

Deficiencies for nicotinic acid—or its amide which appears from limited evidence to be physiologically equivalent to the free acid in fungi—occur in relatively few fungi *Microsporum audouinii* (5) *Blas*

tocladia pringsheimii (35) *Lenturia inaequalis* (63) *Phlyctothiza variabilis* (227) and individual isolates of *Trichophyton equinum* (79) and *Glomerella cingulata* (270). Deficiencies may be complete or partial.

The known function of nicotinic acid is as a constituent of the respiratory coenzymes diphosphopyridine nucleotide and triphosphopyridine nucleotide (257). Diphosphopyridine nucleotide has been isolated from *Fusarium tricothecioides* (87).

The biosynthesis of nicotinic acid by mutants of *Neurospora crassa* has been intensively investigated with important results. These studies are reviewed by Bonner and Yanofsky (20) and a suggested scheme is shown in Figure 4. Evidence for this pathway comes primarily from the ability of compounds to replace nicotinic acid in the nutrition of




Figure 4 The probable pathway of conversion of tryptophan to nicotinic acid in *Neurospora crassa*. Modified from Greenberg (88) by permission of Academic Press Inc.

deficient mutants and from the accumulation of intermediates by deficient strains. Thus one nicotiniless mutant cannot use quinolinic acid and accumulates detectable amounts of this compound indicating that it is blocked in the conversion of quinolinate to the vitamin (19, 102). Other mutants accumulate 3-hydroxyanthranilic acid (18) or a derivative of kynurenine (328) or convert one intermediate into others (330).

As indicated in Figure 4, the steps from 3-hydroxyanthranilic acid to nicotinic acid are still uncertain and may follow one of the two possibilities shown via unstable intermediates (83).

Preliminary studies of nicotiniless mutants of *Aspergillus nidulans* (19c) suggest that a second and shorter pathway occurs in addition to the reactions of Figure 4; this shorter sequence cannot be detected in *Neurospora crassa* (187).

The *Neurospora* pathway is probably followed also in the rat, for which some direct enzymatic evidence is available (257). In the bacteria so far studied, the pathway does not appear to be applicable (328). Evidence on other fungi is limited to the finding mentioned earlier that tryptophan and certain other intermediates of the *Neurospora* pathway replace nicotinic acid for *Trichophyton equinum* (51, 79).

10. INOSITOL

Requirements for inositol (*meso*-inositol, *D*-inositol, *myo*-inositol) are rather rare in fungi; complete or partial deficiencies have been reported in, for example, yeasts (30), *Trichophyton faviforme* (82), *Lophoderium pinastri* (123), *Sclerotinia camelliae* (8), *Rhizopus stolonatus* (238), *Eremothecium ashbyi* (243, 331), *Diplocarpon rosae* (253), and *Diaporthe phaseolorum* var. *batalicola* (283). Inositolless mutants have been found in *Neurospora crassa* (9) and other fungi (17, 70, 121).

It is doubtful whether inositol should be considered a vitamin as the word is used here. No coenzyme function is known in any organism; rather, the finding that most of the inositol of *Neurospora crassa* mycelium is associated with phospholipid suggests that it is a structural component of the cell (74).

The magnitude of the inositol requirement is much greater than that of the other vitamins (136, 259, 281), again suggesting a structural rather than a catalytic role. The weight of dry cells per gram thiamine needed is about $0.5\text{--}2.5 \times 10^6$, and the comparable value for inositol is only 600–900 (65). Figure 5 illustrates the growth response to inositol.

The inositol requirement is highly specific (12c, 242). Phytin, the salt of inositol hexaphosphoric acid, replaces inositol for *Eremothecium*




Figure 5 The response of an inositolless mutant of *Neurospora crassa* to inositol. From Leonian and Lilly (136) courtesy of the West Virginia University Agricultural Experiment Station

ashbyii (170) but neither phytin nor lipositol is utilizable by an inositolless mutant of *Neurospora crassa* (9)

As mentioned earlier no general function of inositol is known in fungi. Inositol overcomes the inhibition caused by high levels of some other vitamins (238-253) and has been stated to cause a conidial type of growth in *Ophiostoma multiannulatum* (72). The response of *Sclerotinia camelliae* to inositol is conditioned by temperature (8).

The insecticide γ hexachlorocyclohexane (gammexane) has been investigated as a possible inhibitory analogue of inositol; it is not however structurally an analogue of *meso* inositol but of *muco* inositol (125). Gammexane inhibition of *Neurospora crassa* appears to be based on two mechanisms: an interference with the metabolism of exogenous inositol in an inositol requiring mutant and a second mechanism which is independent of inositol metabolism (93).

11 PARA AMINOBENZOIC ACID

Only one fungus *Blastocladia ramosa* deficient for *p* aminobenzoic acid (PABA) is known to occur naturally (39a); a yeast *Rhodotorula aurantiaca* is reported to show an absolute requirement (221) and several bacteria require the compound (323). Mutants of several filamentous fungi require PABA (17, 68, 83, 115, 195) so it is presumably a normal metabolite. The deficient mutant of *Neurospora crassa* grows maximally with PABA at about 5 μg per liter (281); the requirement is lower at acid pH than at neutral pH, presumably by reason of a faster penetration of the undissociated acid into the cell (325). The requirement of the *N. crassa* mutant is highly specific; of closely related compounds only 2-fluoro *p* aminobenzoic acid has appreciable activity (323).

Although the folic acid molecule contains the structure of *p* amino benzoic acid it seems doubtful that the major function of PABA is in the formation of folic acid for example folic acid does not replace PABA for *Neurospora crassa* (1 333) Several metabolic roles of PABA have been suggested especially a function in the introduction of 1-carbon intermediates into other compounds (254) The fact that methionine replaces PABA for a mutant of *Neurospora crassa* supports this possibility the vitamin here probably functions in the methylation of homocysteine (269) Other functions no doubt exist (266) but they have not yet been shown in the fungi

The biosynthesis of PABA by *Neurospora crassa* is a special case of aromatic synthesis A single gene mutant is deficient for PABA tryptophan phenylalanine and tyrosine shikimic acid replaces all these requirements (280) The probable pathway of the synthesis is considered elsewhere (Chapter 8)

PABA in high concentrations is toxic to fungi (38) but in *Neurospora crassa* it is converted to a pigment of unknown structure and to N acetyl *p* aminobenzoic acid (200) The acetylation of PABA occurs also in animal tissues (323)

The metabolic importance of PABA was first recognized by Woods (307) who found that the bacteriostatic effect of sulfanilamide is competitively antagonized by PABA The same antagonism has been observed in fungi (24 48 99) and we may assume that the same mechanism prevails

Emerson and Cushing (58) isolated a sulfonamide resistant mutant of *Neurospora crassa* and a second mutant requiring sulfonamides for growth The requirement is absolute at 35 C partial at lower incubation temperatures (56) Later studies by Zalokar (333) suggest that the sulfa requiring strain is abnormally sensitive to the PABA which is formed in its own cells the beneficial effect of sulfonamides is in this view based on its antagonism of the toxic effect of PABA Some of the difficulties of this hypothesis are discussed by Mitchell (171) a relationship to amino acid metabolism is apparently involved (57 334) The present uncertain status of the sulfonamide requirement reflects the limitations of growth methods used by themselves for the solution of essentially biochemical problems

12 PANTOTHENIC ACID

Requirements for pantothenic acid although fairly common in yeasts (27 29 134) are rare or unknown in the filamentous fungi *Penicillium digitatum* is stimulated in the early stages of growth by

pantothenate (61 315) and it has been reported that *Polyporus texanus* requires pantothenate (332) Pantothenicless mutants have been obtained frequently in the fungi (83 121 195 278)

Pantothenic acid functions in metabolism as a constituent of the larger molecule coenzyme A an essential cofactor in a number of acyl transfer and related reactions (147 184) The effect of pantothenate on carotenogenesis in *Mucor hiemalis* presumably results from an increase in the rate of acetyl transfer (90)

Presumably all fungi except those needing exogenous pantothenate synthesize the vitamin this is borne out by analysis of for example *Boletus edulis* (40) *Neurospora crassa* (297) and a few other fungi (181) The ultimate step in pantothenate synthesis is known to be the coupling of pantoic acid and β alanine This coupling is carried out by growing cells resting cells and a cell free preparation (acetone powder) of *Neurospora crassa* (296) Surprisingly the enzyme is found also in a pantothenicless mutant the cells of which do not under the usual conditions of still culture form pantothenate from the two precursors (295) However it has been found that the coupling reaction above does proceed in pantothenicless cells grown under aeration (297) Either we have here in still culture a gene controlled suppression of an enzyme or the mutant strain is unable under limited aeration to provide the energy necessary for the formation of the peptide bond of pantothenic acid

The origin of the pantoic acid moiety in fungi is not known in bacteria *p* aminobenzoic acid is involved and there is some evidence that α ketoisovaleric acid is a precursor of pantoic acid (266)

13 PTEROYLGLUTAMIC ACID

Pteroylglutamic acid (vitamin Bc *Lactobacillus casei* factor folic acid folacin) and its relatives are required by microorganisms and higher animals probably the several active compounds are all converted to the same form (265) *Penturia inaequalis* is reported to be stimulated in its growth by pteroylglutamic acid (63) no fungi or actinomycetes are known to require the factor

A substance active for *Lactobacillus casei* is produced by several fungi including species of *Absidia* *Aspergillus* *Chaetomium* *Circinella* *Fusarium* *Monascus* *Mucor* *Neurospora* *Penicillium* and *Rhizopus* (30 164 178) The identity of this product is uncertain *Citrovorum* factor another member of this group of compounds is formed by *Streptomyces griseus* (59)

Rhizopterins N formylpteronic acid is formed by *Rhizopus nigricans*

(306) and replaces pteroylglutamic acid for strains of *Streptococcus lactis* (265). Presumably it and the factor active for *L. casei* are both utilized by the fungi producing them in the same biological reactions for which the coenzyme form of pteroylglutamic acid is required in bacteria and animals.

14 THE VITAMINS B₁₂

The vitamins B₁₂ comprise a family of compounds the chemical and biological relationships of which are not fully understood. Vitamin B₁₂ itself or cyanocobalamin was the first discovered and its structure is now known (22, 110). Other cobalamins e.g. hydroxocobalamin can be converted to it by cyanide treatment. A second group of compounds differs from the cobalamins in the nucleotide moiety; these are active for some microorganisms only (25, 62). Vitamin B₁₂ is believed to function in the cell in the synthesis of 1-carbon fragments especially the labile methyl group (25).

Vitamin B₁₂ and related compounds are required by several bacteria and algal flagellates (62) not so far as is now known by actinomycetes or fungi. Higher animals require the vitamin; green plants do not.

The synthesis of cobalamins occurs in certain of the true bacteria and especially in species of *Streptomyces* and *Nocardia* of the Actinomycetales (32, 43, 53, 94, 116, 203, 209, 232, 267, 305). Commercial supplies of the vitamin can in fact be obtained as byproducts of antibiotic production by *Streptomyces* spp. (324). Cobalt added to the culture medium increases the yield of cyanocobalamin from *Streptomyces griseus* (104).

In addition to cyanocobalamin species of *Streptomyces* form hydroxocobalamin (vitamin B₁₂, B₁₂_H) in which the cyanide of cyanocobalamin is replaced by hydroxyl and vitamin B₁₂_C containing a nitrite radical in place of cyanide. All three forms have been isolated from one species *S. griseus* (43).

Synthesis of vitamin B₁₂ in the filamentous fungi has not been demonstrated conclusively. Substances active for *Lactobacillus leichmanii* are formed by *Ashbya gossypii* (258) and *Aspergillus niger* (179). However the *L. leichmanii* assay is not specific enough to support a final identification (62).

It has been mentioned earlier that inhibition studies point to 1,2-dimethyl-4,5-diaminobenzene as a precursor of riboflavin. These investigations (310, 311) implicate this compound also as a precursor of the cobalamins; the structure is known to be part of the larger cobalamin molecule. Other evidence bears out the suggestion that the

biosyntheses of riboflavin and of cobalamins are related (69 266) *Streptomyces olivaceus* incorporates labeled 5-6-dimethylbenzimidazole without dilution into the cobalamin molecule (299) more recent and still preliminary evidence is in favor of 2-aminolevulinic acid as a precursor (250)

15 VITAMINS K

Vitamins K₁ and K₂ are required by higher animals. Both are derivatives of 2-methyl-1,4-naphthoquinone and it appears that any compound containing this nucleus is active in animals (3).

Many bacteria produce vitamin K (2) and the phthiocol of *Mycobacterium tuberculosis*—chemically 2-methyl-3-hydroxy-1,4-naphthoquinone—has K activity (3). The six fungi that have been tested (7) do not have detectable vitamin K activity. Other 1,4-naphthoquinones are known to be produced by fungi (Chapter 6) but none of these is a 2-methyl derivative.

No requirement for the K vitamins is known in the fungi. Almquist (3) reviews the conflicting evidence for a K requirement in *Mycobacterium paratuberculosis* (313).

Some compounds structurally related to vitamin K, specifically 2-chloro-1,4-naphthoquinone (246) and 2,3-dichloro-1,4-benzoquinone are fungitoxic (Chapter 14). Menadione (2-methyl-1,4-naphthoquinone) competitively inhibits the action of both of these compounds in certain biological systems (246 309). However the fungitoxicity and the reversal by menadione are not adequate evidence for a role of the K vitamins in fungal metabolism; one may equally well consider the antagonists as foreign toxicants competing for the same site of action (309).

16 OTHER GROWTH FACTORS

The only well characterized growth factor for fungi that has not so far been mentioned is coprogen, required by species of *Pilobolus* and first isolated from manure (107). Bioassays indicate that this factor is manufactured by some other fungi and several actinomycetes (108) and crystalline coprogen has been isolated from *Penicillium* sp. (193).

Coprogen is related chemically and biologically to two other materials: ferrichrome and the *Arthrobacter terregens* factor. Ferrichrome and ferrichrome A are nitrogenous iron-containing pigments of *Ustilago sphaerogena* (76 176); the effect of zinc on synthesis suggests that ferrichrome is a precursor of cytochrome c (177). The

growth factor for *Arthrobacter terregens* although not identical with either coprogen or ferrichrome is evidently related to them thus both ferrichrome and coprogen are active for *A. terregens* (33 148 149) and *Pilobolus* spp (177a)

Sterols are produced by many fungi (Chapter 6) so far only *Labyrinthula vitellina*—of uncertain taxonomic position but possibly a myxomycete—has been shown to have a steroid requirement (293 294) It has been claimed (4) that anaerobic growth of *Saccharomyces cerevisiae* in a synthetic medium is dependent on ergosterol or other sterols

Many still unknown factors have been postulated for particular fungi Mycorrhizal fungi are stimulated by materials from plant roots and forest litter and the effect cannot be traced to known vitamins or amino acids (166 167 168) The myxomycetes apparently require for their growth one or more bacterial proteins (112 272) not vitamins as we have here used the term As mentioned earlier growth stimulating effects of crude natural materials do not necessarily implicate a vitamin

Other vitamins—ascorbic acid and vitamins A D and E—required by animals do not appear to play any essential role in the fungi (28 86) ascorbic acid may however affect growth through its effect on the oxidation reduction potential

17 THE OCCURRENCE OF VITAMIN DEFICIENCIES IN THE FUNGI

It will be clear from the preceding discussion of individual vitamins that some deficiencies occur frequently others more rarely The vitamin most often required by fungi from nature is thiamine usually the deficiency is for the pyrimidine moiety and we have noted only a few instances of a deficiency for thiazole (Table 1) Biotin is very often required and deficiencies in pyridoxine or its derivatives are frequently encountered Pantothenic acid and nicotinic acid requirements are rare in nature and we have at present only single reports of requirements for riboflavin and *p*-aminobenzoic acid among filamentous fungi

This lopsided distribution of vitamin requirements demands an attempt at an explanation Assuming that spontaneous mutations in nature are like those induced by mutagens forms deficient for riboflavin or *p*-aminobenzoic acid must arise from time to time Their failure to survive—if it is real and not an artifact of screening methods—may result from either or both of two conditions viz.

- 1 The required vitamin is not available in the environment
- 2 The derangement of metabolism associated with the deficiency is so great that even with the missing factor available in the environment the mutant is at a selective disadvantage and cannot propagate itself

The second of these hypotheses is the more attractive inasmuch as even the plant pathogens living in close contact with higher plant cells do not seem to develop strains deficient for certain vitamins

18 VITAMINS TAXONOMY AND ECOLOGY

Comparative physiology often affords clues to taxonomic and phylogenetic relationships. Vitamin deficiencies do not, so far as can be determined, offer such clues. From the frequency of induced mutations one would expect that vitamin deficiencies occur at random within species, genera and higher categories; this appears to be true. Marked differences between the vitamin demands of strains of a single species are often encountered: e.g. *Trichophyton* spp. and other animal pathogens (5, 80, 81, 152); *Lenzites trabea* (140); *Boletus granulatus* (169) and *Glomerella cingulata* (270). In culture both *Fusarium avenaceum* (214) and *Ustilago major* (15) give rise to clones which differ in requirements from the parent strain.

Within a genus again species differ widely in vitamin demands: examples include the genus *Ophiostoma* and its asexual form *Ceratostomella* (66, 67, 215, 217, 224); *Trichophyton* (31, 80, 222, 223); *Ustilago* (245); *Candida* (241); *Boletus* (169); *Marasmius* (145) and *Absidia* (213).

In the light of this intra- and interspecific variation it is difficult to imagine that vitamin needs per se can separate taxons or can yield valid evidence of phylogeny.

The theory of Lwoff (150) and Knight (122) that metabolic deficiencies have arisen in the evolution of microorganisms by loss of functions originally present has been generally accepted. If we grant in view of the argument just made that this loss of function has not followed phylogenetic or taxonomic divisions we may ask if there is any relation of vitamin needs to ecology.

First it appears that some genera as isolated from nature rarely have vitamin deficiencies. Only a tentative listing can be made at present in view of the methodological difficulty in establishing the absence of a requirement: *Aspergillus*, *Chaetomium*, *Fusarium*, *Mucor*, *Penicillium*, *Rhizopus*, *Streptomyces* and *Zygorhynchus* (213). Exceptions occur in most of these genera: i.e., some isolates are deficient or pur

tially deficient for a given vitamin (61 195 214) and deficient mutants may be obtained in both *Aspergillus* and *Penicillium* (17 115 19.) These genera are all vigorous saprophytes widespread in nature and frequently isolated from soil. They seem to form therefore an ecological type independent of taxonomic boundaries at present however it is fruitless to speculate on the biochemical genetic or environmental determinants involved.

Soil inhabiting fungi are not however as a group independent of vitamins many soil fungi requiring them for maximal growth (6). Soil especially soil of high organic matter contains several vitamins (234 235) probably formed by higher plants (298) or algae (78) it is not surprising therefore to find deficient soil fungi.

Certain major ecological groups show no general pattern of vitamin dependence these include animal pathogens (5 52 223) plant pathogens (45 46 55 215 217 244 336) and the aquatic phycomycetes as a group (36). Among the insect pathogens however limited data suggest a tendency for species of wide host range e.g. *Isaria farinosa* to be independent of growth factors and more specialized pathogens e.g. *Hirsutella* spp. to require vitamins and organic nitrogen (154).

The nearest approach to a correlation of a taxonomic group with a vitamin requirement is in the Agaricales almost all the fungi in this group that have been investigated require thiamine replaceable for most by the pyrimidine moiety (71 119 145 146 166 183 213 332). Ecologically these include wood-destroying litter decomposing and mycorrhizal groups i.e. groups living in environments in which it is likely that thiamine is present. A similar situation may prevail in some of the orders of the lower phycomycetes (36) but too few species have been investigated far fewer than among the Agaricales.

BIBLIOGRAPHY

- 1 Agarwala S. C. and W. H. Peterson. 1950. *Arch. Biochem.* 27: 304-315.
- 2 Almquist H. J. 1941. *Physiol. Revs.* 21: 194-216.
- 3 Almquist H. S. 1944. In W. H. Sebrell Jr. and R. S. Harris (eds) *The Vitamins* Vol. 2, p. 389-399. 415-418. New York: Academic Press.
- 4 Andreasen A. A. and T. J. B. Suer. 1953. *J. Cellular Comp. Physiol.* 41: 23-36.
- 5 Arêa Leão A. E. de and A. Cury. 1950. *Mycopath. et Mycol. Appl.* 5: 65-90.
- 6 Atkinson R. G. and J. B. Robinson. 1955. *Can. J. Botany* 33: 81-288.
- 7 Barner H. D. and E. C. Cantino. 195. *Am. J. Botany* 33: 746-751.
- 8 Barnett H. L. and V. G. Lilly. 1948. *Am. J. Botany* 35: 297-307.
- 9 Beadle G. W. 1944. *J. Biol. Chem.* 156: 683-689.
- 10 Beadle G. W. and F. L. Tatum. 1941. *Proc. Natl. Acad. Sci. U. S.* 27: 499-506.
- 11 Bender A. E., H. A. Krebs and N. H. Horowitz. 1949. *Biochem. J. (London)* 4: xxxi-xxxi.

- 12 Benham R W 1941 *Proc Soc Exp Biol Med* 46 176-178
- 13 Benham R W 1945 *Proc Soc Exp Biol Med* 58 199-201
- 14 Bhargava K S 1946 *Lloydia* 9 13-23
- 15 Blumer S 1940 *Mitt naturforsch Ges Bern* 1940 19-36
- 16 Blumer S 1944 *Ber schweiz botan Ges* 54 605-624
- 17 Bonner D M 1946 *Am J Botany* 33 788-791
- 18 Bonner D M 1948 *Proc Natl Acad Sci U S* 34 5-9
- 19 Bonner D M and C Yanofsky 1949 *Proc Natl Acad Sci U S* 35 546-581
- 20 Bonner D M and C Yanofsky 1951 *J Nutrition* 44 603-616
- 21 Bonner J and E R Buchman 1939 *Proc Natl Acad Sci U S* 25 164-171
- 22 Bonnet R *et al* 1935 *Nature (London)* 176 398-330
- 23 Bradley S G and J Lederberg 1936 *J Bacteriol* 72 219-225
- 24 Brian P W 1944 *Nature (London)* 153 83-84
- 25 Briggs G M and F S Daft 1955 *Ann Rev Biochem* 24 339-39
- 26 Brown E G T W Goodwin and S Pendlington 1933 *Biochem J (London)* 61 37-46
- 27 Burkholder P R 1943 *Am J Botany* 30 206-211
- 28 Burkholder P R and I McVeigh 1942 *Science* 93 197-128
- 29 Burkholder P R I McVeigh and D Moyer 1944 *J Bacteriol* 48 383-391
- 30 Burkholder P R I McVeigh and K Wilson 1943 *Arch Biochem* 7 987-303
- 31 Burkholder P R and D Moyer 1943 *Bull Torrey Botan Club* 70 370-311
- 32 Burton M O and A G Lochhead 1951 *Can J Botany* 29 330-359
- 33 Burton M O F J Sowden and A G Lochhead 1954 *Can J Biochem and Physiol* 32 400-406
- 34 Cannon M D R K Boutwell and C A Elvehjem 1943 *Science* 100 509-530
- 35 Cantino E C 1948 *Am J Botany* 35 238-242
- 36 Cantino E C 1955 *Quart Rev Biol* 30 138-149
- 37 Carpenter C C and E W Friedlander 1942 *Science* 93 603
- 38 Cavill G W K and J M Vincent 1945 *Nature (London)* 153 301
- 39 Clark F M N C Banister and W R Mitchell 1948 *Arch Biochem* 17 3-9
- 39a Crasemann J M 1957 *Am J Botany* 44 218-224
- 40 Dagys J and P Bluzmanas 1943 *Ber deut botan Ges* 61 49-66
- 41 Dalglish C E 1955 *Advances in Protein Chem* 10 33-150
- 42 Dammann E O T Rotimi and F F Nord 1938 *Biochem Z* 297 185-200
- 43 Darken M A 1933 *Botan Rev* 19 99-130
- 44 Day D 1942 *Bull Torrey Botan Club* 69 11-20
- 45 Defago G 1939 *Ber schweiz botan Ges* 49 413-414
- 46 Defago G 1940 *Phytopathol Z* 13 293-315
- 47 De Vries W H *et al* 1950 *J Am Chem Soc* 72 4838
- 48 Dimond N S 1941 *Science* 94 420-421
- 49 Dittmer K D B Melville and V du Vigneaud 1944 *Science* 99 203-203
- 50 Doudney C O and R P Wagner 1933 *Proc Natl Acad Sci U S* 39 1043-1052
- 51 Drouhet E 1954 *Congr Intern Botan 8 Congr Paris Rapports Communs 8 Sect* 18-20 112-113
- 52 Drouhet E and F Mariat 1952 *Ann inst Pasteur* 82 337-347
- 53 Dulaney E L and P L Williams 1953 *Mycologia* 45 345-358
- 54 Eakin R A and E A Eakin 1942 *Science* 96 187-188
- 55 Elliott E S 1949 *Proc West La Acad Sci* 20 65-68
- 56 Emerson S 1947 *J Bacteriol* 54 197-207
- 57 Emerson S 1949 *Cold Spring Harbor Symposia Quant Biol* 14 40-48
- 58 Emerson S and J E Cushing 1946 *Federation Proc* 5 379-389

- 59 Emery W B & A Lees and A D Walker 1950 *Biochem J* (London) 46 512-574
- 60 Emenegger T 1954 *Ber schweiz botan Ges* 61 453-486
- 61 Fergus C L 1950 *Mycologia* 44 183-189
- 62 Ford J F and S H Hutner 1950 *Vitamins and Hormones* 13 101-136
- 63 Fothergill P G and R Ashcroft 1950 *J Gen Microbiol* 10 387-390
- 64 Friend J and T W Goodwin 1954 *Biochem J* (London) 57 434-437
- 65 Fries N 1938 *Symbolae Botan Upsalienses* 3 (2) 1-188
- 66 Fries N 1940 *Svensk Botan Tidskr* 36 451-466
- 67 Fries N 1943 *Symbolae Botan Upsalienses* 7 (2) 5-73
- 68 Fries N 1945 *Aktu Botan* 30 (8) 1-9
- 69 Fries N 1947 *Nature* (London) 159 199
- 70 Fries N 1948 *Hereditas* 31 338-350
- 71 Fries N 1949 *Svensk Botan Tidskr* 43 316-341
- 72 Fries N 1950 *Aktu Botan* Ser 2 1 271-287
- 73 Fuller R C R W Barratt and E L Tatum 1950 *J Biol Chem* 186 893-897
- 74 Fuller R C and E L Tatum 1956 *Am J Botany* 43 361-363
- 75 Gaden E L Jr D N Petsiavas and J Winoker 1954 *J Ag Food Chem* 2 634-638
- 76 Garibaldi J A and J B Nelands 1955 *J Am Chem Soc* 77 2409-2430
- 77 Garnjobst L and E L Tatum 1956 *Am J Bot* 43 149-157
- 78 Gaumann E and O Jaag 1950 *Phytopathol Z* 17 218-228
- 79 Georg L K 1949a *Proc Soc Exp Biol Med* 7 65-65
- 80 Georg L K 1949b *Trans N Y Acad Sci* 11 281-86
- 81 Georg L K 1950 *Mycologia* 42 693-699
- 82 Georg L K 1951 *Mycologia* 43 297-309
- 83 Giles N H 1946 *J Bacteriol* 50 501
- 84 Goldschmidt M C and H Koffler 1950 *Ind Eng Chem* 42 1819-1823
- 85 Gorca H J W H Petersson and H Steenbock 1935 *J Nutrition* 9 691-699
- 86 Gottlieb D and G M Culligan 1946 *Arch Biochem* 10 163-164
- 87 Gould B S A A Tytell and H Jaffe 1942 *J Biol Chem* 146 219-224
- 88 Greenberg D M 1954 In D M Greenberg (ed) *Chemical Pathways of Metabolism* Vol 2 p 113-147 New York Academic Press
- 89 Grimm P W and P J Allen 1954 *Plant Physiol* 29 369-377
- 90 Grob E C A Grundbacher and W H Schopfer 1954 *Experimenta* 10 378-379
- 91 Guilhermon A M Fontaine and A Raffy 1935 *Compt rend (Paris)* 201 1077-1080
- 92 Gyorgy P 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 1 p 57-388 New York Academic Press
- 93 Haag E 1940 *Compt rend oc phys et hist nat Genève* 57 136-139
- 94 Hall H F et al 1953 *Appl Microbiol* 1 191-199
- 95 Halliday W J and E McCoy 1955 *J Bacteriol* 70 461-468
- 96 Harris D L 1950 *Arch Biochem Biophys* 41 294-304
- 97 Harris D L 1955 *Arch Biochem Biophys* 57 240-251
- 98 Harris D L 1956 *Arch Biochem Biophys* 60 3-43
- 99 Hartel H and K Roholt 1946 *Compt rend trav lab Carlsberg Ser physiol* 24 163-171
- 100 Hata T et al 1951 *J Int Bot Soc (Japan)* 5 31-39
- 101 Hawker L F 1911 *Ann Botany* 8 79-90
- 102 Henderson L M 1949 *J Biol Chem* 181 677-68
- 103 Hendon D M C Cavelle A J Peters and T R Wood 1950 *J Biol Chem* 180 61-69

- 104 Hendlin D and M L Ruger 1950 *Science* 111 541
- 105 Herrick J A. and C J Alexopoulos. 1942 *Bull Torrey Botan Club* 69 569-572
- 106 Herrick J A and C J Alexopoulos 1943 *Bull Torrey Botan Club* 70 369-371
- 107 Hesseltine C W et al 1952 *J Am Chem Soc* 74 1362
- 108 Hesseltine C W et al 1953 *Mycologia* 45 7-19
- 109 Hickey R J 1945 *Arch Biochem* 8 439-447
- 110 Hodgkin D C et al 1955 *Nature (London)* 176 325-328
- 111 Hodson A Z 1949 *J Biol Chem* 179 49-52
- 112 Hok K A 1954 *Am J Botany* 41 792-799
- 113 Horowitz N H 1950 *Advances in Genet* 3 33-71
- 114 Huennkens F M 1956 *Experientia* 12 1-6
- 115 Iguchi N 1952 *J Agr Chem Soc Japan* 26 146-151
- 116 Janicki J et al 1954 *Acta Microbiol Polon* 3 3-9
- 117 Janke A 1939 *Zentr Bakteriell Parasitenk Abt II* 100 409-459
- 118 Jansen B C P 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 3 p 426-442 New York Academic Press
- 119 Jennison M W M D Newcomb and R. Henderson 1955 *Mycologia* 41 275-304
- 120 Kavanagh F 1942 *Bull Torrey Botan Club* 69 669-692
- 121 Keitt G W and D M Boone 1954 *Phytopathology* 44 362-370
- 122 Knight B C J G 1945 *Vitamins and Hormones* 3 103-228
- 123 Kogl F and N Fries 1937 *Z physiol Chem Hoppe Seyler's* 249 93-110
- 124 Krampitz L O and D W Woolley 1944 *J Biol Chem* 152 9-17
- 125 Lardy H A 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 2 p 323-329 342-351 New York Academic Press
- 126 Lardy H A and H Peanasky 1953 *Physiol Revs* 33 560-565
- 127 Lavollay J and F Laborey 1941 *Ann fermentations* 6 129-142
- 128 Leaver F W J Leal and C R Brewer 1947 *J Bacteriol* 51 401-403
- 129 Lein J S and P S Lein 1949 *J Bacteriol* 58 595-599
- 130 Leonian L H and V G Lilly 1938 *Phytopathology* 28 531-548
- 131 Leonian L H and V G Lilly 1939 *Phytopathology* 29 592-596
- 132 Leonian L H and V G Lilly 1940a *Am J Botany* 27 18-26
- 133 Leonian L H and V G Lilly 1940b *Plant Physiol* 15 515-525
- 134 Leonian L H and V G Lilly 1942 *Am J Botany* 29 459-464
- 135 Leonian L H and V G Lilly 1945a *J Bacteriol* 49 291-298
- 136 Leonian L H and V G Lilly 1945b *West Va Univ Agr Exp Sta Bull* 319 1-35
- 137 Levine H et al 1949 *Ind Eng Chem* 41 1665-1668
- 138 Lewis R W 1952 *Phytopathology* 42 657-659
- 139 Lilly V G and H L Barnett 1947 *Am J Botany* 34 131-138
- 140 Lilly V G and H L Barnett 1948 *J Agr Research* 77 290-300
- 141 Lilly V G H L Barnett and B G Anderson 1953 *Science* 118 548-549
- 142 Lilly V G and L H Leonian 1940 *Proc West Va Acad Sci* 14 44-49
- 143 Lilly V G and L H Leonian 1944 *Science* 99 205-206
- 144 Lindeberg G 1939 *Svensk Botan Tidskr* 33 85-90
- 145 Lindeberg G 1944 *Symbolae Botan Upsalienses* 8 (?) 1-183
- 146 Lindeberg G and K Molin 1949 *Physiol Plantarum* 2 138-144
- 147 Lipmann F 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 2 p 598-625 New York Academic Press
- 148 Lochhead A G and M O Burton 1953a *Congr intern microbiol 6th Congr* 1 298-301
- 149 Lochhead A G and M O Burton 1953b *Can J Botany* 31 7-29
- 150 Lwoff A 1936 *Ann fermentations* 2 419-477

- 151 Mackinnon J E. 1942 *Bull Torrey Botanical Club* 69 21-26
- 152 Mackinnon J E and R C Arttagaveytia Allende. 1948 *J Bacteriol* 56 91-96
- 153 MacLaren J A. 1952 *J Bacteriol* 63 233-241
- 154 MacLeod D M. 1954 *Ann N Y Acad Sci* 60 (1) 58-70
- 155 McNutt W S Jr. 1954 *J Biol Chem* 210 511-519
- 156 McNutt W S Jr. 1956 *J Biol Chem* 219 365-373
- 157 Mandels G R. 1955 *Am J Botany* 42 921-929
- 158 Marczyński R. 1943 *Am Midland Naturalist* 30 164-170
- 159 Mariat F. 1954 *Ann inst Pasteur* 87 233-236
- 160 Mariat F and E Drouhet. 1953 *Ann inst Pasteur* 84 659-662
- 161 Marsh P B and K Bollenbacher. 1946 *Am J Botany* 33 245-249
- 162 Martin G J. 1951 *Biological Antagonism*. New York: The Blakiston Co. pp 516
- 163 Mathiesen A. 1950 *Physiol Plantarum* 3 93-102
- 164 Matsuda T. 1949 *Vitamins (Kyoto)* 2 121-124 144-145 (*Chem Abstr* 45 10300 1951)
- 165 Meissel M N. 1947 *Nature (London)* 160 269-270
- 166 Melin E. 1948 *Trans Brit Mycol Soc* 30 92-99
- 167 Melin E. 1954 *Svensk Botan Tidskr* 48 86-94
- 168 Melin E and V S R Das. 1954 *Physiol Plantarum* 7 851-8 8
- 169 Melin E and B Nyman. 1941 *Arch Mikrobiol* 12 254-259
- 170 Minoura K. 1950 *J Fermentation Technol (Japan)* 28 125-129 (*Biol Abstr* 26 9227 1951)
- 171 Mitchell H K. 1950 *Vitamins and Hormones* 8 127-150
- 172 Mitchell H K and M B Houlahan. 1946 *Am J Botany* 33 31-35
- 173 Muller F W. 1941 *Ber schweiz botan Ges* 51 165-256
- 174 Müller W and W H Schopfer. 1937 *Compt rend (Paris)* 205 687-689
- 175 Naeata Y A Matuda and M Hiura. 1954 *Research Bull Fac Agr Gifu Univ (Japan)* No 3 100-101
- 176 Neilands J B. 1952 *J Am Chem Soc* 74 4846-4847
- 177 Neilands J B. 1953 *J Biol Chem* 205 647-650
- 177a Neilands J B. 1957 *Bacteriol Revs* 23 101-111
- 178 Newell G W W H Peterson and C A Elvehjem. 1947 *Poultry Sci* 26 84-88
- 179 Nicholas D J D. 1952 *Analyst* 77 6 9-64
- 180 Nicholas D J D. 1956 *J Gen Microbiol* 15 470-477
- 181 Nielsen N and V Hartelius. 1945 *Compt rend t av lab Ca lsbe g s r physiol* 24 117-14
- 182 Noecker N L and M Reed. 1943 *Am Midland Naturalist* 30 171-174
- 183 Norström B. 1950 *Symbolae Botan Upsalienses* 11 (1) 1-16
- 184 Novelli G D. 1953 *Physiol Revs* 33 5 5-543
- 185 Odgård K. 1950 *Physiol Plantarum* 5 583-608
- 186 Otani Y. 1950 *Ann Phytopathol Soc Japan* 7 9-15
- 187 Partridge C W H D M Bonner and C Yanofsky. 1950 *J Biol Chem* 194 769-78
- 188 Payette A and C Perrault. 1944 *Can J Research C* 22 127-132
- 189 Peltzer G L and R Borchers. 1947 *J Bacteriol* 54 519-520
- 190 Perlman D. 1948 *Am J Botany* 35 36-41
- 191 Perlman D. 1951 *Am J Botany* 38 652-658
- 192 Peterson W H and M S Peterson. 1945 *Bacteriol Revs* 9 49-109
- 193 Pidacks C et al. 1953 *J Am Chem Soc* 75 6064-6065
- 194 Plaut G W E. 1954 *J Biol Chem* 208 513-50
- 195 Pontecorvo G et al. 1953 *Advances in Genetics* 5 141-238
- 196 Quantz L. 1943 *Jahrb wiss Botan* 91 150-168

- 197 Rabinowitz J C and E E Snell 1953a *Arch Biochem Biophys* 43 399-401
- 198 Rabinowitz J C and E E Snell 1953b *J Am Chem Soc* 75 998-999
- 199 Raper K B R D Coghill and A Hollander 1945 *Am J Botany* 32 163-176
- 200 Reid L B E G Iritchett and J E Cushing 1952 *J Biol Chem* 199 443-450
- 201 Reissig J L 1952 *Arch Biochem Biophys* 36 234-235
- 202 Renaud J and M Lachaux 1945 *Compt rend (Paris)* 291 187-188
- 203 Rickes E L et al 1948 *Science* 108 634-636
- 204 Robbins W J 1938a *Bull Torrey Botan Club* 65 267-276
- 205 Robbins W J 1938b *Proc Natl Acad Sci U S* 24 53-56
- 206 Robbins W J 1939 *Am J Botany* 26 712-777
- 207 Robbins W J 1941 *Proc Natl Acad Sci U S* 27 419-429
- 208 Robbins W J and A Hervey 1955 *Mycologia* 47 155-169
- 209 Robbins W J A Hervey and M E Stebbins 1950 *Bull Torrey Botan Club* 77 423-441
- 210 Robbins W J and F Kavanagh 1938a *Am J Botany* 25 229-236
- 211 Robbins W J and F Kavanagh 1938b *Bull Torrey Botan Club* 65 453-461
- 212 Robbins W J and F Kavanagh 1944 *Bull Torrey Botan Club* 71 1-10
- 213 Robbins W J and V Kavanagh 1942 *Botan Rev* 8 411-411
- 214 Robbins W J and R Ma 1941 *Bull Torrey Botan Club* 68 446-46
- 215 Robbins W J and R Ma 1942a *Am J Botany* 29 835-849
- 216 Robbins W J and R Ma 1942b *Arch Biochem* 1 219-229
- 217 Robbins W J and R Ma 1942c *Bull Torrey Botan Club* 69 184-203
- 218 Robbins W J and R Ma 1942d *Bull Torrey Botan Club* 69 349-359
- 219 Robbins W J and R Ma 1942e *Science* 96 406-407
- 220 Robbins W J and R Ma 1943 *Bull Torrey Botan Club* 70 190-197
- 221 Robbins W J and R Ma 1944 *Science* 100 85-86
- 222 Robbins W J and R Ma 1945 *Am J Botany* 32 509-523
- 223 Robbins W J J E Mackinnon and R Ma 1949 *Bull Torrey Botan Club* 69 509-521
- 224 Robbins W J and M B Schmidt 1939 *Bull Torrey Botan Club* 66 139-150
- 225 Robinson F A 1951 *The Vitamin B Complex* New York John Wiley and Sons pp 688
- 226 Rother W 1954 *Arch Mikrobiol* 20 89-108
- 227 Rothwell F M 1956 *Am J Botany* 43 28-32
- 228 Ryan F J C Kunin R Ballentine and W Maas 1953 *J Bacteriol* 65 434-439
- 229 Salvin S B 1949 *J Infectious Diseases* 81 275-283
- 230 Sarasin A 1953 *Ber Schweiz botan Ges* 63 287-316
- 231 Sarett H P and V H Cheldelin 1944 *J Biol Chem* 156 91-100
- 232 Saunders A P R H Otto and J C Sylvester 1952 *J Bacteriol* 64 728-728
- 233 Scheunert A and M Schieblisch 1936 *Biochem Z* 286 66-71
- 234 Schmidt E L 1951 *Soil Sci* 71 129-140
- 235 Schmidt E L and R L Starkey 1951 *Soil Sci* 71 221-231
- 236 Schopfer W H 1934a *Arch Mikrobiol* 5 511-549
- 237 Schopfer W H 1934b *Ber deut botan Ges* 52 308-312
- 238 Schopfer W H 1942 *Compt rend soc phys et hist nat Ceres* 59 101-106
- 239 Schopfer W H 1943 *Plants and Vitamins* Waltham Mass Chronica Botanica Co pp 293

- 240 Schopfer W H 1944a *Compt rend soc phys et hist nat Genève* 61 147-150
- 241 Schopfer W H 1944b *Compt rend soc phys et hist nat Genève* 61 230-236
- 242 Schopfer W H 1944c *Helv Chim Acta* 27 408-471
- 243 Schopfer W H 1944d *Helv Chim Acta* 27 1017-1030
- 244 Schopfer W H and S Blumer 1938 *Arch Mikrobiol* 9 305-367
- 245 Schopfer W H and S Blumer 1940 *Arch Mikrobiol* 11 203-214
- 246 Schopfer W H and E C Grob 1949 *Helv Chim Acta* 32 89-838
- 247 Schopfer W H and M Guilleud 194 *Z Vitaminforsch* 16 181-296
- 248 Schopfer W H and W Rytz Jr 1937 *Arch Mikrobiol* 8 244-248
- 249 Sebrell W H Jr and R S Harris (eds) 1954 *The Vitamins* New York Academic Press 3 vol
- 250 Shemin D J W Corcoran C Rosenblum and I M Miller 1956 *Science* 124 272
- 251 Sherwood M B and E D Singer 1944 *J Biol Chem* 150 361-369
- 252 Shmida S S Chava and K M Noura 1950 *J Fermentation Technol (Japan)* 30 15-22 (*Chem Abstr* 48 3449 1951)
- 253 Shirakawa H S 1955 *Am J Botany* 42 379-384
- 254 Shive W 1951 *Vitamins and Hormones* 9 13-150
- 255 Silver W and W D McElroy 1954 *Arch Biochem Biophys* 51 379-391
- 256 Sinclair H M 1937 *Nature (London)* 140 361
- 257 Slinger T P and E B Kearney 1954 *Advances in Enzymol* 15 79-139
- 258 Smiley K L et al 1951 *Ind Eng Chem* 43 1380-1384
- 259 Snell E E 1948 *Physiol Revs* 28 5-98
- 260 Snell E E 1951 In F Skoog (ed) *Plant C and the Substances* p 431-446 Madison Univ of Wisconsin Press
- 261 Snell E E and A N Rannefeld 194 *J Biol Chem* 154 415-489
- 262 Srinivasan K S and C V Ramakrishnan 1950 *Biochim et Biophys Acta* 9 156-160
- 263 Stokes J L J W Foster and C R Woodward Jr 1945 *Arch Biochem* 2 235-245
- 264 Stokes J L M Cunniff and J W Foster 1944 *J Bacteriol* 47 293-299
- 265 Stokstad E L R 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 3 p 89-191 124-216 New York Academic Press
- 266 Stokstad E L R H P Broquist and N H Sloane 1955 *Ann Rev Microbiol* 9 111-144
- 267 Stokstad E L R T H Jukes J Pierce A C Page Jr and A L Franklin 1949 *J Biol Chem* 180 617-654
- 268 Strauss B S 1951 *Arch Biochem* 30 290-303
- 269 Strehler B L 1950 *J Bacteriol* 59 103-111
- 270 Struble F B and G W Keitt 1950 *Am J Botany* 37 563-566
- 271 Sussman M 1951 *Ann Rev Microbiol* 10 21-50
- 272 Sussman M and S C Bradley 1954 *Arch Biochem Biophys* 51 428-433
- 273 Swartz H E and L K Georg 1955 *Mycologia* 47 475-493
- 274 Tanner F W Jr S E Pfeiffer and J M Van Lanen 1953 *Arch Biochem* 8 29-36
- 275 Tanner F W Jr C Vojnovich and J M Van Lanen 1954 *Science* 101 180-181
- 276 Tatum E L 1945 *J Biol Chem* 160 455-459
- 277 Tatum E L and G W Beadle 1940 *Proc Natl Acad Sci U S A* 28 234-243
- 278 Tatum E L and G W Beadle 1943 *Ann Missouri Botanical Garden* 30 125-139

- 279 Tatum E L and T T Bell 1946 *Am J Botany* 33 15-20
- 280 Tatum E L S R Gross G Ehrensward and L Garnjobst 1954 *Proc Natl Acad Sci U S* 40 271-276
- 281 Tatum E L M G Ritchey E V Cowdry and L F Wicks 1946 *J Biol Chem* 163 675-682
- 282 Thompson R C E R Isbell and H K Mitchell 1943 *J Biol Chem* 148 281-287
- 283 Timnick M B V G Lilly and H L Barnett 1951 *Phytopathology* 41 337-336
- 284 Tomizawa C 1952 *Ann Phytopathol Soc Japan* 27 113-118
- 285 Turel F 1952 *Phytopathol Z* 19 307-312
- 286 Umbreit W W 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 3 p 234-242 New York Academic Press
- 287 Umbreit W W W A Wood and I C Gunsalus 1947 *J Biol Chem* 165 731-732
- 288 Van Lanen J M and F W Tanner Jr 1948 *Vitamins and Hormones* 6 163-224
- 289 Villaneuva J R 1955 *Nature (London)* 176 465
- 290 Villela G G 1917 *Bull soc chim biol* 29 763-767
- 291 Villela G G and A Cury 1950 *J Bacteriol* 59 1-11
- 292 Vinson L J L R Cerecedo R P Mull and F F Nord 1945 *Science* 101 388-389
- 293 Vishniac H S 1955 *J Gen Microbiol* 12 464-472
- 294 Vishniac H S and S W Watson 1953 *J Gen Microbiol* 8 248-255
- 295 Wagner R P 1949 *Proc Natl Acad Sci U S* 35 185-189
- 296 Wagner R P and B H Guirard 1948 *Proc Natl Acad Sci U S* 34 398-402
- 297 Wagner R P and C H Haddox 1951 *Am Naturalist* 85 319-330
- 298 West P M 1939 *Nature (London)* 144 1050-1051
- 299 Weygand F H Klebe and A Trebst 1954 *Z Naturforsch* 9b 449-450
- 300 White N H 1941 *J Australian Council Sci Ind Research* 14 137-146
- 301 Wickerham L J M H Flickinger and R M Johnston 1946 *Arch Biochem* 9 95-98
- 302 Willstaedt H 1935 *Ber deut chem Ges* 68B 333-340
- 303 Wilson A N and S A Harris 1949 *J Am Chem Soc* 71 2231-2233
- 304 Wirth J C and F F Nord 1942 *Arch Biochem* 1 143-163
- 305 Witkus E R F Gentile L Kelly and L Murphy 1954 *Can J Botany* 32 503-505
- 306 Wolf D E et al 1947 *J Am Chem Soc* 69 2753-2759
- 307 Woods D D 1940 *Brit J Exp Pathol* 21 74-89
- 308 Woodward V W J R DeZeeuw and A M Srb 1954 *Proc Natl Acad Sci U S* 40 192-200
- 309 Woolley D W 1945 *Proc Soc Exp Biol Med* 60 225-228
- 310 Woolley D W 1950 *Proc Soc Exp Biol Med* 75 745-746
- 311 Woolley D W 1951 *J Exp Med* 93 13-24
- 312 Woolley D W 1952 *A Study of Antimetabolites* New York John Wiley and Sons pp 269
- 313 Woolley D W and J R McCarter 1940 *Proc Soc Exp Biol Med* 45 357-360
- 314 Woolley D W and A G C White 1943 *J Exp Med* 78 489-497
- 315 Wooster R C and V H Cheldelin 1945 *Arch Biochem* 8 311-320
- 316 Worley C L 1947 *Plant Physiol* 17 218-288
- 317 Wright L D and E L Cresson 1954 *J Am Chem Soc* 76 4156-4160
- 318 Wright L D E L Cresson and C A Driscoll 1955 *Proc Soc Exp Biol Med* 89 234-236

- 319 Wright L D E L Cresson K V Liebert and H R Skeggs 1952 *J Am Chem Soc* 74 2004-2006
- 320 Wright L D E L Cresson J Valiant D E Wolf and K Folkers 1954a *J Am Chem Soc* 76 4160-4163
- 321 Wright L D E L Cresson J Valiant D E Wolf and K Folkers 1954b *J Am Chem Soc* 76 4163-4166
- 322 Wright L D and C A Driscoll 1954 *J Am Chem Soc* 76 4999-5000
- 323 Wright L D and P A Favormina 1951 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 3 p 13-2 New York Academic Press
- 324 Wuest H M 1954 In W H Sebrell Jr and R S Harris (eds) *The Vitamins* Vol 1 p 417-421 New York Academic Press.
- 325 Wyss O V G Lilly and L H Leonian 1944 *Science* 99 18-19
- 326 Yagi K Y Matsuoka S Kuyama and M Tada 1956 *J Biochem (Tokyo)* 43 93-100
- 327 Yanofsky C 195 *J Biol Chem* 198 343-352
- 328 Yanofsky C 1954 *J Bacteriol* 68 577-584
- 329 Yanofsky C and D M Bonner 1950 *Proc Natl Acad Sci U S* 36 167-176
- 330 Yanofsky C and D M Bonner 1951 *J Biol Chem* 190 211-218
- 331 Yaw K E 1952 *Mycologia* 44 307-317
- 332 Yusef H 1953 *Bull Torrey Botan Club* 80 43-64
- 333 Zalokar M 1948 *Proc Natl Acad Sci U S* 34 32-36
- 334 Zalokar M 1950 *J Bacteriol* 60 191-203
- 335 Zalokar M 1955 *Arch Biochem Biophys* 56 318-325
- 336 Zscheile F P 1951 *Phytopathology* 41 1115-1124

11. Reproduction

The attention of those interested in physiological aspects of the reproductive process in fungi has been focused almost exclusively on the problem of the initiation of sporulation and these studies occupy the bulk of this chapter. The further development of the reproductive structures has in the main been treated as a problem in morphology and with some exceptions its physiological and biochemical determinants are unknown.

Naturally enough any physiological study tends to emphasize the one or two factors in reproduction which are manipulated by the experimenter. It should of course be obvious that reproduction requires that all genetic and environmental factors be favorable. This complexity is particularly apparent in ecological studies of the higher fungi (115).

Although some tentative conclusions on fungi in general have been proposed it should be emphasized that any organism must be studied individually; there is no one formula for sporulation.

It is fair to say that most of our knowledge of reproduction in the fungi comes from essentially qualitative data. Sporulation occurs under some conditions but not under others and data may be reported in simple yes or no categories. No objection can be made to this procedure in general; however it is becoming clear that some problems require quantitative data on the intensity of sporulation. This is particularly true of nutritional studies in which the response is likely to be quantitative rather than all or none.

It has not been possible with present information to consider the physiology of resting structures e.g. chlamydospores and sclerotia separately or in detail. Chlamydospore formation is probably a response to unfavorable environmental conditions (214 215 231 266).

Sclerotium development is of course often a prelude to sexual reproduction and may be expected to be influenced by the same factors allowing for a higher nutritional level entailed by the larger mass (149 252 290). The sclerotia of myxomycetes are purely resting structures and appear to develop primarily as a response to desiccation (110 154 166).

The geotropism of large sporophores is an established fact of observation (27 49 50 278) but its physiological correlates have not yet been uncovered.

Finally in the list of omissions purely genetic problems are not here considered. These are of course most important to a thorough understanding of reproduction but they lie outside the field of physiology.

1 LIGHT

The reactions of fungi to visible and ultraviolet light are of three principal types inductive effects inhibition and trophic responses. Inductive effects include both absolute light requirements for initiation or maturation of reproductive structures and quantitative responses i.e. an increase in the number of reproductive structures upon illumination.

Light is absolutely required for the formation of various types of reproductive organs in certain species. As examples we may cite the requirement of light for the formation of sporophores in the Agaricales (226 253) of perithecia (264) of apothecia (248 293 319) of basidiocarps in the Gasteromycetes (298) of pycnidia (105 183 185) of sporangia in the Mucorales (221) of conidia in the Mucorales (18) and Fungi Imperfecti (97 159) and of Myxomycete sporangia (117). Smith (263) reviews still other examples.

As might be expected the effect of light is not always exerted at the same point in these diverse fungi to take two extremes light is required for the first steps in sexual reproduction in *Pyronema confluens* (248) and for all stages in *Polyporus arcularius* (1107) but only for pileus development in some Agaricales (27 106). Whether all developmental stages of the *Coprinus lagopus* sporophore require light is in dispute (196 294).

Light of the visible range often has the effect of increasing the number of reproductive structures formed (82 89 120 124 130 264). More striking effects of this type are induced by ultraviolet light in certain ascomycetes and Fungi Imperfecti both sexual and asexual structures are affected (11 66 160 198 232 262 271 272 273 274). Ultraviolet effects are exerted in the cell not in the medium (74 271).

and cannot be ascribed to temperature effects (262) The effective wavelength is still in doubt (88 232) but it is probably longer than the well known germicidal wavelength of $254\text{ m}\mu$ Ultraviolet light of course in high doses represses both sporulation and growth

Most of the work on induction and stimulation by visible light has been at best semiquantitative The available data on the action spectrum indicate fairly consistently that the effective wavelengths are in the blue region of the spectrum (37 83 117 198 248 250a 263 277 319) Morphogenetic action on *Coprinus lagopus* is maximal at $440\text{--}460\text{ m}\mu$ (255) close to the peak of the action spectrum of phototropism

The amount of light needed for inductive response is similarly uncertain it is much less than normal daylight exposure (105 168) The maximal effect of light on fruiting in *Coprinus lagopus* is exerted by exposure for 5 seconds at 0.1 foot-candles intensity (196)

So much attention has been focused on the positive effects of light that the possibly more important inhibitory action may be overlooked Induction of conidiation in *Choanephora cucubitarum* requires a dark as well as a light period (18) and light of wavelength below $580\text{ m}\mu$ appears to be inhibitory (80) Dark periods are required for maximum production of ascospores by *Diaporthe phaseolorum* (286) and of conidiophores by *Alternaria brassicae* (312) For the Myxomycete *Didymium eunigripes* Leith (189 190) presents evidence for both induction and inhibition of sporangial formation by different wavelengths Finally the type of zonation in which spore production is greater during dark periods (see below) is most easily interpreted as resulting from a light inhibition of sporulation

The diurnal cycle of *Pseudoperonospora humuli* and other downy mildew fungi in nature involves both light inhibition and light stimulation although effects on the host metabolism cannot be excluded (315) Other diurnal cycles in nature (314 316) and the frequently observed nocturnal production of conidia (161 162) may also result from light inhibition although the dominant factor in nocturnal formation of spores by *Sclerospora* spp. has been thought to be water (306 307)

Zonation—the formation of fruiting structures in concentric rings on an agar surface or on the host plant—is a common phenomenon and may best be considered at this point since light is usually the primary environmental factor The occurrence of light induced zonation is reviewed by Hall (214) and by Hafiz (123) A simple explanation is rendered difficult by the observation that two extreme types of response to an alternation of light and dark periods are known viz

Type 1 The fungus sporulates well in darkness much less or not at all during the light period Examples are *Cephalothecium roseum* (145) and *Sclerotinia fructicola* (124)

Type 2 The organism e.g. *Fusarium* spp. (29 124 320) sporulates poorly in the dark and forms a zone of spores after even a brief exposure to light

It seems necessary to explain Type 1 reactions as the result of light inhibition of sporulation Type 2 responses are more difficult we may propose however that light inhibits mycelial growth—reports of such inhibition are reviewed in Chapter 13—and that stimulation to sporulation results from the check upon vegetative growth The light effect in these fungi is exerted only upon actively growing hyphal tips (29 45 264)

Temperature alternations also induce the formation of zones of reproductive structures (94 123 131 250a 320) it appears especially from the data of Ellis (94) and Hafiz (123) that the immediate stimulus for sporulation is a check in growth

Zonation of reproductive structures in some fungi is dependent neither upon light nor upon temperature fluctuations (40 75 146 206) presumably nutritional factors are involved here but they are not understood The pH of the medium has an effect (212 240)

Any unitary hypothesis to explain the action of light on sporulation will have to take account of several facts viz

- 1 The indifference of many fungi to light stimuli

- 2 The high specificity of the response Thus the formation of conidia of *Choanephora cucurbitarum* is governed by light but the development of the sporangia is independent of light (18) and the formation of zygospores is inhibited by light at low temperature (20)

- 3 The fact that light may either stimulate or inhibit sporulation

- 4 The partial replacement of the light effect by high temperature (29 37 185 212) by periodic exposure to air (206) by treatment with oxidizing agents (82) or by mechanical agitation (37 275)

- 5 The observation of Houston and Oswald (159) that *Helminthosporium gramineum* forms spores in the dark if growing on host tissue but requires light for sporulation on agar media

- 6 The extreme sensitivity of the response (29 168) which suggests that an enzymatic mechanism is triggered by the stimulus

- 7 The frequent efficacy of ultraviolet radiation known to be injurious to all organisms in sufficient dose Sporulation of *Alternaria solani* is induced by either ultraviolet or mechanical damage (74)

A unified theory should take as its principal postulate that light is inhibitory to one or more regulatory systems which in the normal culture prevent or retard reproduction and favor continued vegetative growth. That is to say there is in truth no such thing as stimulation by light in the induction of reproductive activity; instead growth is checked and the chain of events so initiated leads to reproduction if other factors are not limiting. Alternative explanations are possible and several have held specifically that stimuli for reproduction do not check growth; nevertheless the weight of the evidence favors the hypothesis presented. Confirmation must wait upon studies which determine accurately and quantitatively the effects of light on both reproduction and growth simultaneously.

Phototropic responses are common in fungi; examples include the orientation of basidiomycete sporophores (37, 278, 298) and of asci or ascocarps (8, 41, 53). Gametic phototaxis has been reported (180). The most dramatic and the most thoroughly investigated phototropic reaction is that of the sporangiophores of *Phycomyces* spp. and *Pilobolus* spp., observed also in others of the Mucorales (96).

Consideration of phototropism in *Phycomyces* spp. should begin with the observation of Blaauw (32), confirmed by Castle (68), that a sporangiophore illuminated equally from two sides exhibits a measurable temporary acceleration in vertical growth, the light growth reaction. A more recent study of this phenomenon (86) brings out several important aspects: (1) the response probably involves no net increase in total growth; (2) the effect of the light stimulus depends in a predictable fashion on the level of illumination prior to the application of the stimulus; (3) there is a latent period of about 2.5 seconds; and (4) the magnitude of the increased growth is proportional to the strength of the stimulus. This last observation is important in showing that the light growth reaction is not an all or none or threshold reaction.

Granted that illumination can increase growth, the fact of phototropism—unequal growth—requires some mechanism whereby the sporangiophore wall distal to the light grows more rapidly than the proximal wall. Castle (71, 72) argues that the size, shape, and refractive index of the sporangiophore combine in the lens effect to concentrate unilateral illumination in such a way that the distal region of the protoplasm receives more light than the proximal. This is consistent with the observation that sporangiophores immersed in paraffin oil become negatively phototropic (14, 48), with the effect of ultraviolet light (66a) and with the demonstration by Banbury (14) that

illumination of only one edge of the sporangiophore causes localized extension of that side and consequent curvature


This theory and others of the same general type (32) fail as Delbrück and Reichardt (86) point out to account for the extreme localization of the receptor which is implied by the unidirectionality of the phototropic curvature if the entire growing zone were affected unilateral illumination and the known spiral habit of growth would combine to generate a helical shape of the sporangiophore

The action spectrum of phototropism in *Phycomyces blakesleeanus* (Figure 1) and *Pilobolus kleini* has a maximum at about 430–500 m μ (54 55 69 99) This has often been taken as evidence for a carotenoid receptor pigment carotenoids being present in these fungi and absorbing maximally in the same region of the spectrum (297) However as Galston (107) points out riboflavin or a derivative of it could with equal probability be the receptor Carotenogenesis in *Phycomyces* sp can be suppressed at least below detectable levels without impairment of the phototropic response (241)

Nor are there any solid data on chemical mediators between the light absorbing system and the growth (or wall extension) response In higher plants it has been suggested that auxin (indoleacetic acid) is involved but its role is not yet entirely certain (107) Fungi do produce auxin (Chapter 8) and it is found in fact in the sporangiophore of *Phycomyces nitens* (151) but there is at present no reason to accept—and there is considerable evidence against—an auxin theory of phototropism in the fungi (14)

Phototropism in *Pilobolus kleini* has been investigated especially by Buller (53) who reviews the earlier literature In this species a pig

Figure 1 The action spectrum of phototropism in *Phycomyces blakesleeanus* Redrawn from Castle (69) by permission of The Rockefeller Institute for Medical Research



mented ocellus at the base of the subsporangium is believed to be the photosensitive area but it should be noted that photosensitivity is apparent before the formation of the subsporangium (222) and that other phototropic forms do not have an ocellus

Other light responses of *Pilobolus* spp may be briefly noted. Light induces the formation of the trophocyst or sporangial primordium of some isolates. Page (221) raises the possibility of a flavin receptor in this reaction. Second alternations of light and dark, or of temperature, set up a periodic rhythm in the development of sporangia (175-196). This rhythm, once established, persists for two or three 24-hour cycles after the removal of the stimulus (254-291). Finally, the forcible discharge of the sporangium is hastened by light (55).

Light has other effects on the reproductive apparatus, notably a positive effect on the size and septation of macrospores of *Fusarium* spp (130-264). Pigment production is also enhanced by light (189-264) but attempts to connect this phenomenon with induction of sporangia in *Didymium eumigris* were unsuccessful (189).

Exposure to light, unless appropriate filters and cooling devices are used, means an increase in temperature. Thus the apparent attraction of myxomycete pseudoplasmodia to light is a temperature effect: the plasmodium moves toward a source of heat along a temperature gradient as small as 0.05°C per centimeter (35). Similarly, a negative phototropic response of *Phycomyces* sp. to intense light was shown by Castle (70) to be a heat effect. Many reports on light effects have not included quantitative determinations of the ambient temperature; possibly some of the apparent complexities may disappear when such data become available.

2 TEMPERATURE

Only two generalizations can be formulated with respect to temperature and sporulation. First, the temperature range permitting reproduction is usually narrower than that permitting growth. Second, each spore form has its own temperature optimum, which may or may not coincide with that of vegetative growth or of other spore forms of the same species.

The relative narrowness of the temperature limits of reproduction is virtually universal in the fungi. Examples include both sexual and asexual spores in the phycomycetes (174) and ascomycetes (148-194, 281), sporophores of basidiomycetes (5-182) and spores of the Fungi Imperfecti (2-204, 287). Usually the differences in range are not

great *Phoma apicola* is an extreme with mycelial growth on agar at 5-28 C but pycnidium formation only at 13-26 (25)

The optimum temperature for sporulation of a given species may be either close to the growth optimum as in *Piricularia oryzae* (149) and *Rhizopus* spp (303) or rather different lower in *Fusarium conglutinans* (287) and higher in *Sphaerotheca pannosa* (194)

Different spore forms of a given species often differ in their temperature requirements. Thus the zygotes of *Sporodinia grandis* and of *Saprolegnia mixta* form at lower temperatures than do the corresponding asexual sporangia (172-173). Perithecia of *Aspergillus* spp (171-256) and of *Ceratostomella* (*Ophiostoma*) *fimbriata* (17) are produced in abundance at relatively high temperature and are replaced by conidia at lower. *Pseudopeziza ribis* forms conidia at 20-24 C microconidia at lower temperatures (33)

It must be borne in mind that most of these data are only semi-quantitative and that the same uncertainties are involved in the definition of an optimum for reproduction as in the definition of a growth optimum (Chapter 1). One such complication is illustrated by the findings of Robbins and Schmitt (245) on *Phycomyces blakesleeana*: the failure of progametes to form at 26 C is the result of the accumulation of acid in the medium before the two strains meet. Presumably progametes would form at 26 if the pH were controlled.

One may speculate that sexual stages adapted to survive a cold winter are formed preferentially at low temperature and that fungi which must live through a hot and dry season will tend to form their dormant spores at higher temperature. However there is as yet no ecological or physiological data bearing on this possibility and the failure of sexual reproduction in *Neurospora crassa* at high temperature (305) speaks against the idea. The asci of *Erysiphe graminis* mature somewhat more rapidly if chilled briefly but they do mature without cold treatment (114). The apothecia of *Claviceps purpurea* appear from limited data to develop only if the sclerotia bearing them are exposed to low temperature (170). Zygosporangium formation in *Thamnidium* spp is also best at low temperature (150).

Temperature induced zonation has been considered elsewhere with the conclusion that the positive effect of high or low temperature on spore formation is probably effected by retardation of mycelial growth.

Morphological effects of temperature are easily observed: as examples we may cite the finding that the conidia of *Cercospora sesami* are larger at an intermediate temperature than at higher or lower tem-

peratures (79) and the effect of temperature on the morphology of the conidial fructification in *Aspergillus janus* (239)

3 WATER RELATIONS

Satisfactorily quantitative determinations of the moisture requirements of sporulation are rare. Indications are however that specific fungi have specific requirements. At one extreme *Colletotrichum lindemuthianum* and other anthracnose fungi sporulate much more profusely at high than at low humidity (318). Conidiation in *Sclerotinia* spp. (306-307) and in *Peronospora destructor* (317) is also favored by high humidity. The formation of sexual structures in ascomycetes and basidiomycetes (10-111-283) and sporangium formation in *Sporodinia grandis* (147) are believed to proceed more rapidly at high than at low moisture levels.

Somewhat lower humidities are tolerated but are not necessarily required by some of the rust fungi (318) by *Botrytis cinerea* (140) and by *Bremia lactucae* (228). Species of *Alternaria* in culture sporulate rather better if the culture is allowed to dry out (90-233). Finally the powdery mildew fungi appear to be relatively independent of humidity or possibly favored in both sexual and asexual reproduction by lower atmospheric moisture (47-125-318). Asci of *Erysiphe graminis* are reported to form only if the perithecium is alternately wetted and dried (76) but it seems doubtful that this is a general requirement among the powdery mildew fungi.

The weight of evidence favors the generalization that reproductive processes in most fungi at least are favored by moderate rather than high atmospheric humidity. However one may see from even a superficial glance at the original work that accurate control of the environment and quantitative determination of the intensity of sporulation have not usually been achieved.

Different spore forms of a single species occasionally show different humidity responses. Examples include the two asexual spore forms of *Choanephora cucurbitarum* (19) and the zygote and sporangium of *Sporodinia grandis* (13-172-217).

Miscellaneous reported effects of atmospheric humidity include negative hydrotropism of sporangiophores (172-300) transitory acceleration of the growth of sporangiophores of *Phycomyces nitens* at high humidity (300) and changes in the size or morphology of conidia (98-119) and sporophores (39).

Spore discharge in fungi has been studied in some detail the sum

mary of Ingold (162) has been supplemented more recently by investigations of spore discharge in rust fungi (220) and *Nigrospora sphaerica* (302). With the possible exception of drop excretion discharge in basidiomycetes (50 51 52 229) all these discharge mechanisms rely in some way on either substrate or atmospheric water. We infer that osmotic forces are at work. In immature asci (162) and in the immature active layer of *Sphaerobolus* spp. (299) glycogen is abundant; its disappearance at maturity suggests that the uptake of water is mediated by enzymatic conversion of the osmotically inactive polysaccharide to osmotically active sugars. Such a mechanism, if suitably quantified, might be extended to cover all discharge mechanisms which depend on increasing turgor of a cell or cell layer.

The elevation of the receptaculum in the Phallales is accompanied by disappearance of glycogen, but in this instance the increase in size may be true growth (58 95). Elongation of the *Agaricus campestris* sporophore has been shown to be a growth process rather than simple water uptake (36).

1 OXYGEN, CARBON DIOXIDE AND VOLATILE SUBSTANCES

It is a common observation that sporulation is reduced in closed containers (82 126 185) and that submerged mycelium fails to develop spores (81 201 321). Although this is usually ascribed to an oxygen deficit, it seems more likely that the accumulation of carbon dioxide is the critical factor. *Pyronema confluent*, for example, does not fruit in a closed tube, but if alkali is present to absorb carbon dioxide, perithecia develop (248). Carbon dioxide accumulation inhibits the asexual reproduction of *Choanephora cucurbitarum* (19) and sporophore development in *Agaricus campestris* (181) and *Collybia velutipes* (227).

In large culture containers the sporulation of *Piricularia oryzae* is dependent upon aeration of the vessel. Henry and Anderson (149) suggest that metabolically produced ammonia is the inhibitory factor. An unidentified volatile substance, probably of microbial origin, inhibits sporophore development in *Psalliota campestris* (199).

The best available data on the oxygen requirement for sporulation are those of Denny (87). Any reduction of oxygen below atmospheric pressure delays the appearance of perithecia. Further, usually estimated mycelial growth appeared to be normal at oxygen pressures too low for reproduction. The same phenomenon—a higher oxygen requirement for fruiting than for growth—is reported by Ternetx (283).

Plant pathogenic fungi usually and with an obvious ecological advantage form their reproductive structures on the surface of the host. It is reasonable to speculate that either the higher oxygen requirement or the lower carbon dioxide tolerance of sporulation is the mechanism that ensures extramatrical sporulation. It is significant that active anaerobic respiratory mechanisms are common in the lower phycocetes and the genus *Fusarium* forms in which resting spores or conidia often occur within host tissues (93).

Several *Penicillium* spp. sporulate in shaken cultures but the determinants are not known (102-122). In our laboratory we have observed that *Fusarium solani* forms abundant macrospores in shake culture particularly in certain media. Conidia of *Streptomyces griseus* appear in aerated cultures at the time of autolysis (67).

In a very brief note Richards (212) reported that ozone appears to stimulate sporulation in *Alternaria* spp. *Mycosphaerella citrullina* forms more pycnidia if exposed to ozone but the spores are non viable.

The striking effect of carbon dioxide and bicarbonate on sporangial development in *Blastocladia* and *Blastocladiella* spp. is considered later in this chapter.

5 ACIDITY

Although pH has definite effects on reproduction no unitary hypothesis or generalization is possible as mentioned elsewhere (Chapter 1). pH effects are exerted in a variety of ways. Furthermore the common failure to determine the final pH casts doubt on much of the available data. In general it appears that the pH range for sporulation is somewhat narrower than that for growth (5, 20, 23, 192, 193, 276). Within the physiological range an acid reaction is often less favorable than a neutral or mildly alkaline reaction (170, 245, 276) but exceptions to this rule have been reported (5, 156).

Lockwood (192) showed that in several fungi perithecia form over a wide range of pH values but asci mature only at neutral or slightly alkaline reaction. This finding is of particular importance because often the only data submitted in pH studies concern macroscopically visible fruit bodies.

In *Mycosphaerella pinodes* pycnidia, pseudothecia and chlamydo spores are reported to have different pH optima (266) but other data on the same species are not easily reconciled with this claim (23). The pH range for sporangium formation by *Physarum polycephalum* is narrowed by high temperature cultivation (116).

6 CARBON NUTRITION

The classical work on the influence of nutrition and for that matter of all environmental factors on reproduction in fungi is that of Klebs (172 173 174). In relation to nutrition his major points may be formulated briefly

1 Sexual and asexual reproduction are most likely to occur when a vigorous mycelium exhausts its nutrients or is transferred to a medium low in nutrients

2 Formation of fruiting structures therefore occurs over a narrower range of nutrient concentrations than does vegetative growth

3 Fruiting structures of different fungi or different spore forms of the same fungus have specific requirements for nutrient concentration. In *Saprolegnia mixta* and *Sporodinia grandis* the sexual spore stage requires a higher nutrient level than the asexual stages

With minor modifications—most importantly that the concentration of the carbon source is usually more decisive than that of other nutrients—these generalizations still stand. It would be tedious indeed to list all the fungi that conform to Klebs' predictions; some of the more significant modern results with pure cultures deserve however some attention.

Suppression relative or absolute of reproductive stages at higher concentrations of carbohydrate—concentrations that permit heavy vegetative growth—has been found in studies of ascomycetes (59 137 141), phycmycetes (267) and imperfect fungi (44 45 82). More extensive reviews of the literature on this topic are available (140 185).

The same phenomenon can be demonstrated more neatly by transfer experiments in which a mycelium growing in a concentrated medium is transferred to or allowed to grow into a more dilute medium. Such experiments almost always confirm the first principle of Klebs (6 81 82 142 185 219 296). In general transfer to water is less successful than transfer to a dilute medium.

Finally it follows from the first principle that spore formation should occur at the point in the culture cycle at which the carbon source is substantially exhausted. Studies on myxomycetes (62 238) demonstrate this principle clearly. The same response has been shown in *Melanconium fuligineum* in a different manner: heavy inoculation of an agar surface accelerates sporulation because of the more rapid utilization of nutrients (285).

It has often of course been found in studies of particular fungi that Klebs first principle does not hold or is overridden by other factors (78 184 185) this is best documented in studies on fleshy basidiomycetes (185 22b) in which the relatively greater mass of the fruit body might be expected *a priori* to change the usual relation

The mechanism by which high nutrient concentrations inhibit sporulation is not definitely established. One cause which must be operative in some cultures is simply the accumulation of toxic metabolites. Effects of culture age and of concentration have been attributed to this phenomenon (104 251). It seems however unlikely that such accumulation can be a general explanation of the concentration effect. For the present we may posit no doubt oversimplifying the picture that a limitation of growth per se is the stimulus to reproduction and that the effect of high nutrient levels is merely to postpone the onset of starvation.

The second principle of Klebs is a corollary of the first. Granted that some minimum amount of growth must precede fruiting and that too high concentrations delay or inhibit fruiting then the range of concentration suitable for reproduction must necessarily be narrower than that suitable for vegetative growth.

The third principle of specificity of requirements is obvious insofar as different species are concerned. Even within a species however different spore forms have different requirements. Examples include the sexual and asexual stages of *Sporodinia grandis* (13 96) and the alternative asexual stages of *Blakeslea trispora* (112).

The concentration of the carbon source affects the type of asexual spore formed in *Cytosporina ludibunda* and related fungi (218) and the septation of macrospores of *Fusarium* sp. (157).

The type of carbon source used strongly influences the formation of reproductive structures both sexual and asexual. In general it is found that oligosaccharides and polysaccharides support more fruiting than do the simple hexoses. Cellulose a very slowly available compound for most fungi is the best carbon source for sporulation of some forms (85 319).

The mechanism of specific carbon source effects has been thoroughly investigated by Hawker and her collaborators (133 137 141) and by Bretzlöff (42). There seems little doubt that the primary mechanism is actually a concentration effect i.e. it exemplifies Klebs first principle. Essentially provision of a slowly utilizable carbon compound is the same as use of a low concentration of a more available sugar. That is the fungus hydrolyzes (or phosphorylates) the complex carbohydrate slowly the result being a steady maintenance of a low con




Figure 2 The response of growth and fruiting to carbohydrate concentration M and F mycelial growth and fruiting respectively with a readily available carbohydrate M and F the same with a more slowly available carbohydrate M and F the same with a very slowly utilizable carbohydrate. From Hawker (140) by permission of the University of London Press Ltd.

concentration of hexose. This point is shown admirably in Figure 2 from Hawker (140).

This principle explains many of the effects of particular carbon sources on reproductive activity. Thus the optimum sucrose concentration for fruiting in *Melanospora destruens* is ten times that in *M. zamae* indicating—and this is borne out by sucrase assays—that the latter species utilizes sucrose much more rapidly than the former (141).

Hawker (137–138) has suggested that the favorable effect of sucrose is in part to be attributed to the formation of phosphorylated hexoses from it by the fungus. This it must be admitted seems unlikely and is certainly not established by data available at present. Phosphorylated compounds have however been tentatively implicated in studies on stimulation by contaminating organisms (p. 372).

7 NITROGEN NUTRITION

The concentration of the nitrogen source in the medium is important for fruiting, i.e. an optimum concentration can be determined

under specified conditions account must be taken however of the carbon source and its concentration (257 305) In general high concentrations of nitrogen suppress sporulation (109 153 250)

Specific effects of particular compounds are often reported but can not be combined in a general theory pH effects and contamination of organic nitrogen sources with growth factors unduly complicate interpretation of experimental data Asparagine so often a favorable source of nitrogen for growth generally represses sporulation (5 28 208 226 257) The asparagine effect may arise from ammonia accumulation or may be another example of the competition between vegetative growth and reproductive activity It should also be noted that asparagine of the sources of nitrogen tested is the best for perithecium formation in *Diaporthe phaseolorum* (286)

The effect of hypoxanthine (or guanine) on zygospore formation in *Phycomyces blakesleeanus* is probably not specifically a reproductive phenomenon and is considered elsewhere (Chapter 12) The very striking positive effect of low concentrations of histamine on zygospore formation in *P. blakesleeanus* (250) merits reinvestigation

8 MINERAL NUTRITION

Numerous observations some of which have been mentioned in Chapter 9 attest to the influence of mineral nutrition on sporulation these are reviewed in more detail by Foster (101) As expected either deficiencies or excesses of particular elements reduce or inhibit sporulation these reactions require no further comment The important question is whether the mineral requirements for sporulation differ qualitatively or quantitatively from those for growth

There is no evidence for a qualitative difference i.e. sporulation never requires an element which is non-essential to growth Quantitatively however there are differences most of which are subsumed in the generalization that sporulation is reduced by partial deficiencies in certain elements much more than growth is reduced by the same deficiency Experiments with copper almost invariably show that sporulation of *Aspergillus niger* is reduced drastically by a deficiency which has a small or even no effect on dry weight (205 211 246 269 270 313) This copper effect is found also in work with *Penicillium* spp and *Phoma betae* (205) The effect of copper deficiency on spore pigment has been noted earlier (Chapter 9)

The same phenomenon—sharp reduction in sporulation with only a small decrease in growth—occurs with partial deficiencies of iron (246) or manganese (26 155 193 269 270) It is a general rule even

with a major element like potassium (209) or magnesium (230) that sporulation is prevented by deficiencies which do not completely inhibit growth. That is sporulation requires more of a mineral than the minimum necessary for measurable growth.

Much has been made of the repressive effect of added zinc on sporulation (38 113 152 165 246). Toxicity of course is possible but it seems more likely that in the concentrated media used provision of zinc simply postpones sporulation by allowing more luxuriant vegetative growth.

The response of *Chaetomium globosum* to calcium (21 22) is an apparent contradiction to the principle that sporulation and growth have qualitatively the same mineral requirements: calcium does not affect dry weight but does increase perithecial formation. The fact that many fungi including other *Chaetomium* spp. do respond in growth to calcium and the possibility that calcium is an antagonist rather than a nutrient (Chapter 9) are arguments against immediate revision of the principle. It may be noted that sporulation in *Aspergillus niger* is inhibited by high levels of NaCl without any apparent effect on growth (201).

One rather serious reservation attaches to most of the results mentioned in this section—sporulation is estimated visually rather than measured quantitatively. A second limitation is that most of the work has been with *Aspergillus niger*; study of other organisms might cause us to modify some of the conclusions now accepted.

9 VITAMINS AND UNIDENTIFIED FACTORS

Under this head we consider only substances which exert a non-specific effect: hormones and hormone-like substances are taken up in the next section.

The data available show that most fungi which require biotin or thiamine need more of the vitamin for adequate sporulation than they do for measurable vegetative growth (17 134). Thus *Sordaria fimicola* in a medium without biotin makes about one third maximal growth but does not form perithecia unless the vitamin is supplied; in this species biotin deficiency results in abortion or failure to mature of ascospores (16).

However one cannot specify an optimum level of a vitamin without reference to other constituents of the medium. *Ceratostomella fimbriata* for example requires thiamine in proportion to the total concentration of the medium (17). Conversely it is well established for numerous fungi that provision of thiamine raises the level of

carbohydrate optimal for fruiting (59 132 133 134 135 140) The simplest explanation of this phenomenon is based on the fact that thiamine accelerates glucose utilization (136), this is in accord with our previous conclusion that exhaustion of the carbon source is required for the initiation of reproduction

There is no conclusive evidence that any fungus requires a known vitamin specifically for the reproductive process and independently of growth effects The available data can more or less satisfactorily be explained on the basis of the two generalizations just stated that fruiting may require more of a vitamin than measurable growth and that vitamins may accelerate sugar utilization and so hasten the onset of reproduction However at least the possibility of a more direct relation is raised by the work of Barnett and Lilly (17) on *Ceratostoma fimbriata* a thiamine starved mycelium begins to fruit if transferred to a thiamine solution containing no other nutrients

Unidentified materials in crude plant and microbial extracts often increase fruiting Examples include both sexual reproduction (164 186 226 259 322) and asexual reproduction (121 144 169)

It is often found usually as a result of accidental contamination of cultures that other fungi and bacteria encourage fungal reproduction Many such cases are reviewed by Asthana and Hawker (6) and by Raper (236) other instances may also be cited here (24 143 243 258) However there is no assurance that these are vitamin effects—pH for example or nutrient concentration could be the decisive factor Stimulation of conjugation of *Zygosaccharomyces* spp by *Aspergillus niger* has been described the effect of the fungus is replaced by a mixture of riboflavin and an organic acid (216 217) The association in nature of *Gonatorrhodiella highleyi* with *Nectria coccinea* may be explained by some chemical influence of the latter on the reproduction of *G. highleyi* (7)

Another series of investigations reviewed by Buston and Khan (60) suggests that the effect of contaminating organisms on peritheciium formation in *Melanospora destruens* and *Chaetomium globosum* is to be attributed to organic phosphorus compounds particularly 3-phosphoglyceric acid liberated by the inducing fungus Whether this is a direct effect is however not yet clear In *C. globosum* perithecial production is associated with a high level of organic phosphate esters (61)

Several apparently unrelated synthetic organic compounds are reported to increase sporulation without a corresponding increase in growth possibly the effect is an antagonism of inhibitors present in the growth medium (158 295)

10 SPECIFIC REPRODUCTIVE HORMONES

Hormones—endogenous chemical substances which influence or determine reproductive processes—have been unequivocally demonstrated in certain of the Oomycetes notably *Achlya* spp. by J. R. Raper who reviewed the entire field in 1952 (236).

Certain strains of *Achlya bisexualis* and *A. ambisexualis* are heterothallic, i.e. the male and female sex organs—antheridia and oogonia respectively—are borne on different thalli. This type of heterothallism is perhaps more clearly described as haplodioecism. When thalli of opposite sex are in the same medium the development of the sex organs is under hormonal control. The hormones whose existence and action must be inferred from convincing physiological and morphogenetic evidence are summarized in Table 1. It is apparent that the

Table 1 The Sexual Hormones of *Achlya* Spp

Hormone	Produced by	Action
A A ²	Female vegetative plant	Induces antheridial hyphae on male plant
A ¹	Male vegetative plant	Augments A and A ²
A ³	Male vegetative plant	D presses action of A and A ²
B	Antheridial hyphae	Induces oogonial initials on female plant
C	Oogonial initials	(a) Induces antheridial hyphae to grow along a concentration gradient toward the oogonium (b) induces the formation and delimitation of the antheridium in the male plant
D	Antheridia	Induces (a) delimitation of the oogonium and (b) differentiation of oospheres

From Raper (236)

entire sequence of events leading up to fertilization is under specific and sequential hormonal control. The hormones are produced in extremely small amounts so small that only a start has been made toward chemical fractionation (237). Biological activities indicate that the battery of hormones produced by *A. bisexualis* is not fully identical with that of *A. ambisexualis* (234).

Others of the Saprolegniaceae—*Thraustotheca clavata* and the homothallic *Achlya americana*—are governed in reproduction by similar but not identical hormone systems (235, 236). Preliminary evidence of a hormonal system in *Sapromyces reinschii* has also been obtained (30).

Burgess (56) studied *Mucor mucedo*, a heterothallic Zygomycete which copulates by the fusion of identical gametangia when two strains of opposite mating type are brought together. The two morphologically identical types are called for convenience plus (+) and minus (-) historically at least the term heterothallism is appropriately applied to this situation. Burgess showed that at least two of the steps in reproduction are initiated by a chemical diffusible stimulus. Zygomorph development in each strain is begun only when the opposite strain is close enough for the stimulus to reach receptive hyphae. The somewhat clumsy term telemorphosis has been used for this reaction. A second reaction also initiated by a diffusible metabolite has been called zygotropism. zygophores of each strain bend toward those of the opposite strain preparatory to contact and fusion.

These reactions in *Mucor mucedo* have been confirmed (167-176, 225) and found also to prevail in *Pilobolus crystallinus* (177-178). Limited evidence suggests that the zygotropic reaction is effected by a volatile product (15-167). Such a material would have to be of very high activity to induce a differential growth response in an organ as small as the zygophore.

Banbury (14) reports evidence that zygophores of the same strain in *Mucor mucedo* exert a mutual repulsion also by a volatile substance. This recalls the numerous examples of aversion or barrage, a zone of inhibition develops between certain colonies of the same or of different species and not between others. This phenomenon is reviewed by Cayley (73) and by Takemaru (282). little physiological work has been done although it has been suggested that a volatile material is responsible (293).

A final type of interstrain chemical influence is shown by a slight increase in oxygen uptake when (+) and (-) mycelia occupy the same respirometric vessel (57).

None of the hormonal activities of the Zygomycetes has been separated physiologically or biochemically from the others. it is not yet possible therefore to describe any such elegant sequential mechanism as that of *Achlya* spp.

At this point we may briefly mention the numerous attempts to find physiological or biochemical differences between (+) and (-) strains of species of the Mucorales. Various workers have claimed that the strains differ in growth rate on certain sugars, sucrase or catalase formation, lipid or carotenoid content, sugar content, reductive capacity, susceptibility to poisons, and so on. Most of these are reviewed by Ronsdorf (250) and by Raper (236). Study of the original papers and of the divergent results obtained leaves one skeptical of all this work.

The most recent claimed difference—a color reaction with strong alkali (292)—was found invalid almost immediately (46). Enzyme differences between opposite mating types in *Neurospora crassa* (103) and growth differences between (+) and (−) strains of *Ustilago* spp. (234) are exposed until confirmed to a similar skepticism.

The long debate over the existence of perithecial inducing hormones in *Neurospora sitophila* is reviewed by Raper (236); the work of Aronescu (3, 4) in particular seems conclusively negative. This conclusion must stand unless challenged by new evidence. However in *Glomerella cingulata* like *Neurospora* a member of the Sphaeriales there is preliminary evidence for a diffusible metabolite which induces the formation of perithecia (92, 200). Since the various steps of sexual reproduction in *G. cingulata* are gene-controlled (308) it is perhaps not too fanciful to envision a one gene—one hormone relation.

In *Ascobolus stercorarius* the induction of ascogonia and other early steps in sexual reproduction are controlled by diffusible materials, presumably hormones (31). The branching of the trichogyne in *Neurospora sitophila* (9) and the suppression of conidiation in *Glomerella cingulata* by filtrates from perithecial cultures (309) are also very probably hormone effects. There is no evidence as yet for hormonal mechanisms in basidiomycetes.

The literature on the chemical stimulus for the aggregation of myxamoebae in the Acrasidae has been thoroughly reviewed by K. B. Raper (238) and by Sussman (279). The substance responsible has been termed acrasin (34) and has now been separated from the cells; it was found to consist of two components and to be destroyed by some system, probably an enzyme of the cell (260, 280). It is especially interesting that acrasin not only attracts scattered myxamoebae but also stimulates them in turn to produce acrasin; the effect is that of a chain reaction which extends the range of the chemotactic system much beyond any possible range based only on diffusion from a single point.

11 METABOLISM AND REPRODUCTION

Throughout this chapter we have stressed the concept that the major stimulus to reproduction is exhaustion of the medium. However many different metabolic changes occur at the same time in a culture and the possibility must be kept in mind that one or more of these changes is the primary stimulus to spore formation. In particular the mycelium begins to autolyze (Chapter 1). In yeast colonies on agar ascospores appear in that part of the colony which is undergoing autolysis (191). Now it is possible that some factor in normal cells is

released by their death and reaches a threshold concentration high enough to induce sporulation. This possibility also relates to the stimulation of reproduction by ultraviolet radiation. It has been suggested (286) that cells killed by the radiation release into the medium substances which favor perithecial development. This line of investigation deserves more attention than it has so far received.

Other metabolic changes are often coincident with the onset of reproduction, e.g. changes in respiratory rate (118), appearance of tyrosinase (153) or other enzymes (100) and formation of carotenoids (57a, 150a, 288, 289). The correlation between tyrosinase formation and protoperithecium development in *Neurospora crassa* is the most striking in that a wide variety of environmental factors suppress both enzyme activity and reproductive capacity, but it cannot yet be said that the relationship is causal (304).

One obvious physiological necessity in sporulation is the transport of materials—presumably in a soluble form—from the mycelium to the developing spores. In *Aspergillus flavus* the appearance of conidia is coincident with a marked fall in mycelial nitrogen, easily interpreted as resulting from transport (224). Phosphate transfer is even more dramatic in *A. niger* during spore formation; mycelial phosphorus declines as much as 80 per cent and at least some of this phosphorus appears in the spores (12).

The transport of materials into fruiting structures may be more complete than it is usually thought to be. Thus microscopic observation of *Ascochyta pin* in an infected leaf shows the mycelium apparently devoid of protoplasm at the time of formation of pycnidia (43).

Blastocladiella pringsheimii in its natural habitat forms two types of sporangia: thin-walled zoosporangia and thick-walled resting sporangia. In pure culture, resting sporangia form only at very high carbon dioxide pressure, specifically in 99.5 per cent CO_2 at pH 5.5 (63). This discovery initiated an important series of investigations by Cantino and collaborators of members of the related genus *Blastocladiella*. These investigations have been reviewed by Cantino (64) and a brief summary is in order at this point.

Blastocladiella emersonii zoospores develop into a two-celled thallus consisting of a basal rhizoidal cell and an apical sporangium. The sporangium is either a thin-walled papillate zoosporangium or a thick-walled resistant sporangium. In pure culture the type of sporangium formed is controlled by bicarbonate: virtually all thalli develop resistant sporangia if 0.01 M bicarbonate is supplied to the medium. The problem thus is not that of initiation of reproduction but of the type

of reproduction which occurs. The explanation proposed by Cantano for the effect of bicarbonate is that the ion in some way interferes with the oxidative decarboxylation of α -ketoglutaric acid. Concurrently several other enzyme activities of the plant disappear and a new polyphenol oxidase becomes evident. The upshot of these changes is a reorientation of metabolism resulting in the formation of the resistant sporangium; this involves necessarily the synthesis of the chitin, melanin and γ -carotene characteristic of the resistant sporangium.

These conclusions have been buttressed by study of a mutant strain of *B. emersonii* which does not even with bicarbonate form resistant sporangia. Crude cell free extracts from the mutant strain are devoid of aconitase and of the enzyme system oxidizing α -ketoglutarate via the cytochromes (65). Precisely these enzymes are postulated to be required in the normal strain for the expression of the bicarbonate effect.

The rather elaborate metabolic hypotheses which have been erected in the course of this study of reproduction in *Blastocladiella* are somewhat provisional and speculative; further investigation is clearly necessary. But however the hypothesis may have to be modified later, it should be recognized as a bold and imaginative attempt to correlate biochemical events and patterns of morphogenesis.

12. THE SPORULATION OF PLANT PATHOGENS

Reproductive processes in fungi pathogenic to or deriving their food from higher plants are difficult to study but pose at least two interesting questions: the influence of the host on sporulation in general, and specific relations between the host and particular stages in the life cycle of the pathogen.

It is a commonplace that plant pathogens often lose their capacity to sporulate during continuous artificial culture. Of course this may only be an example of the very common phenomenon, not at all restricted to plant pathogens, of the selection in culture of non-sporulating variants (77, 127, 128, 138, 207, 213, 223, 244, 310, 311). However there is still the question of how contact with the host—often only in a single passage—acts to select for restoration of reproductive capacity.

It seems probable that specific features of the food plant promote sporulation over and above any selective process. We have already noted the important observation that the host tissue replaces a light requirement for the sporulation of *Helminthosporium graminum*. Experiments with autoclaved plant extracts are of little real help in working out the problem, since labile materials are probably involved. Cold sterilization of natural substrates by, for example,

propylene oxide appears much more promising as a technique (129 265) Analogously the insect pathogen *Cordyceps militaris* forms perithecia on dead pupae but not on autoclaved pupae (261)

The relation of the host to sporulation has one other dimension the influence of the host on the intensity and on the type of sporulation This has been most thoroughly studied in relation to the appearance of the telial stage of rust fungi From the work of Morgenthaler (210) and Waters (301) we may list factors that encourage or hasten the development of the telial stage

- 1 Exposure of the host to low temperature or to slow drying
- 2 Starvation of the host by dark incubation
- 3 Growth of the fungus on a partially resistant host
- 4 Wounding of the inoculated leaf
- 5 Transfer of detached leaves from sucrose to water

Obviously effects of this type must be indirect for the obligate parasites but the pattern seems clear that telia form under conditions of privation the analogy to Klebs conclusions is close The factor that is dominant in nature is not known it is most probable that drying out in autumn is dominant (91 108) but simple starvation cannot be dismissed

Perithecial formation in *Erysiphe graminis*—also an obligate parasite—is more rapid if the relatively resistant older leaves of the host are infected (114) Here again we may speculate that resistance of the host accelerates the reproduction of the pathogen by depriving it of some essential factor

Host influences on spore size merit brief mention Uredospores of rust fungi are smaller on a resistant host than on a susceptible one and they are smaller if the host is held under conditions unfavorable to its growth (187 188 268) Similar findings on other fungi have been summarized by Fischer and Gaumann (98)

13 CONCLUSIONS

The major barrier to generalizing about reproduction in the fungi is the repeated failure of experimental work to discriminate adequately between growth and sporulation too often only one is measured quantitatively Consequently many of our hypotheses on reproduction must leave open the possibility that the factor concerned may act on growth rather than specifically on the formation of spores

The most inclusive generalization is that reproduction is initiated by factors which check the growth of an established mycelium without

too drastically poisoning its metabolism. Studies on the effect of light especially ultraviolet light are the most important evidence in view of the well-established lethal action of radiation. Gross mutilation by scraping or cutting induces sporulation in a few fungi (195: 206-233) in scraped cultures of *Alternaria solani* Kunkel (179) observed that conidiophores arise from broken hyphae. Initiation of sporulation as the result of temperature changes, exhaustion of nutrients, and various treatments of the host plant of a pathogen are all consistent with the injury hypothesis.

Granted that a check in growth initiates sporulation, we can only speculate on the intimate mechanism. Perhaps the most promising suggestion is that injured or moribund cells release substances which act on surviving cells and divert them into a new developmental path.

It must be noted that a contrary hypothesis—that sporulation requires specific positive stimuli rather than the negative stimulation of injury—could easily be built up. In particular, the ready sporulation of many fungi on the living host plant or on cold sterilized plant materials argues for nutritional factors which must be present in some threshold amount to permit sporulation or which must be synthesized by the fungus before reproductive activity can begin.

Less inclusive but important and more firmly based generalizations may be listed briefly in conclusion.

1. Reproduction occurs in general over a narrower range of environmental conditions than does growth; this is especially clear in findings on temperature, acidity, mineral and oxygen requirements. The optimum for reproduction may be the same as or different from that of mycelial growth.

2. Klebs' principle—that reproduction is favored by depriving an established mycelium of one or more nutrients—is valid in general, especially for carbon and nitrogen sources. It must be qualified only to take account of the fact that for some nutrients—certain vitamins and minerals—the absolute amount required for sporulation is higher than that which supports growth. This qualification essentially only changes the definition of the "established mycelium" to indicate that it must have certain materials in high concentration.

3. There is no conclusive evidence for a specific nutritional factor required for sporulation and not required at all for growth. Specific endogenous hormones are, however, established as vital to reproductive activity or at least to the orderly development of reproductive structures once these are initiated.

4. The requirements for sexual and asexual reproduction in a given

propylene oxide appears much more promising as a technique (129 265) Analogously the insect pathogen *Cordyceps militaris* forms perithecia on dead pupae but not on autoclaved pupae (261)

The relation of the host to sporulation has one other dimension the influence of the host on the intensity and on the type of sporulation This has been most thoroughly studied in relation to the appearance of the telial stage of rust fungi From the work of Morgenthaler (910) and Waters (301) we may list factors that encourage or hasten the development of the telial stage

- 1 Exposure of the host to low temperature or to slow drying
- 2 Starvation of the host by dark incubation
- 3 Growth of the fungus on a partially resistant host
- 4 Wounding of the inoculated leaf
- 5 Transfer of detached leaves from sucrose to water

Obviously effects of this type must be indirect for the obligate parasites but the pattern seems clear that telia form under conditions of privation the analogy to Klebs conclusions is close The factor that is dominant in nature is not known it is most probable that drying out in autumn is dominant (91 108) but simple starvation cannot be dismissed

Perithecial formation in *Erysiphe graminis*—also an obligate parasite—is more rapid if the relatively resistant older leaves of the host are infected (114) Here again we may speculate that resistance of the host accelerates the reproduction of the pathogen by depriving it of some essential factor

Host influences on spore size merit brief mention Uredospores of rust fungi are smaller on a resistant host than on a susceptible one and they are smaller if the host is held under conditions unfavorable to its growth (187 188 268) Similar findings on other fungi have been summarized by Fischer and Gaumann (98)

13 CONCLUSIONS

The major barrier to generalizing about reproduction in the fungi is the repeated failure of experimental work to discriminate adequately between growth and sporulation too often only one is measured quantitatively Consequently many of our hypotheses on reproduction must leave open the possibility that the factor concerned may act on growth rather than specifically on the formation of spores

The most inclusive generalization is that reproduction is initiated by factors which check the growth of an established mycelium without

- 39 Bortels H 1927 *Biochem Z* 18^o 301-308
- 39 Bose S R 1943 *Mycologia* 3^o 33-46
- 40 Brandt W H 193 *Mycologia* 45 194-203
- 41 Brefeld O 1877 *Botan Ztg* 3^o 386
- 42 Bretzloff C W Jr 1934 *Am J Botany* 41 58-67
- 43 Brewer D and B H MacNeill 1953 *Can J Botany* 31 739-744
- 44 Brian P W and H G Hemming 1930 *Trans Brit Mycol Soc* 3 132-141
- 45 Brown W 1935 *Ann Botany* 39 373-408
- 46 Brucker W 1935 *Arch Protistenk* 100 339-350
- 47 Buchheim A 1928 *Ber deut botan Ges* 4^e 167-180
- 48 Buder J 1918 *Ber deut botan Ges* 36 103-104
- 49 Buller A H R 1909 *Researches on Fungi* Vol 1 London Longmans Green and Co pp 287
- 50 Buller A H R 1909 *Researches on Fungi* Vol 2 London Longmans Green and Co pp 49
- 51 Buller A H R 1914 *Researches on Fungi* Vol 3 London Longmans Green and Co pp 611
- 52 Buller A H R 1933 *Researches on Fungi* Vol 4 London Longmans Green and Co pp 416
- 53 Buller A H R 1934 *Researches on Fungi* Vol 6 London Longmans Green and Co pp 513
- 54 Bunning E 1937a *Planta* 6 719-736
- 54 Bunning F 1937b *Planta* 27 583-610
- 56 Burjeff H 1934 *Botan Abh Heft* 4 3-13
- 57 Burnet J H 1933 *New Phytologist* 52 58-64 86-89
- 57a Burnet J H 1934 *New Phytologist* 53 43-49
- 58 Burt F A 1897 *Bota Gaz* 24 73-92
- 59 Buston H W and S N Basu 1948 *J Gen Microbiol* 16^o-17^o
- 60 Buston H W and A H Khan 1956 *J Gen Microbiol* 14 635-660
- 61 Buston H W and B Rickard 1956 *J Gen Microbiol* 15 194-197
- 62 Camp W G 1937 *Am J Botany* 24 300-303
- 63 Cantino E C 1949 *Am J Botany* 36 93-112
- 64 Cantino E C 1956 *Mycologia* 48 37-40
- 65 Cantino E C and M T Hyatt 1953 *J Bacteriol* 66 71-70
- 66 Carll M J 1956 *J Gen Microbiol* 14 615-644
- 66a Carll M J 1957 *Nature (London)* 180 70
- 67 Carvajal F 1947 *Mycologia* 39 426-440
- 68 Castle E S 1939 *J Gen Physiol* 12 391-400
- 69 Castle E S 1931 *J Gen Physiol* 14 703-711
- 70 Castle E S 1937 *J Gen Physiol* 15 487-489
- 71 Castle E S 1933 *J Gen Physiol* 17 41-60
- 72 Castle E S 1935 *Cold Spring Harbor Symposia Quant Biol* 3 221-229
- 73 Cayley D M 1931 *J Genet* 21 1-63
- 74 Charlton K M 1933 *Trans Brit Mycol Soc* 36 348-350
- 75 Chaudhuri H 1933 *Ann Botany* 37 519-539
- 76 Cherech W J 1944 *Can J Research C* 2 5^o 86
- 77 Chilton S J P 1943 *Mycologia* 3^o 13-20
- 78 Chona B L 1937 *Trans Brit Mycol Soc* 17 221-228
- 79 Chowdhury S 1914 *J Indian Botan Soc* 23 91-107
- 80 Christenberry G A 1938 *J Elisha Mitchell Sci Soc* 54 297-310
- 81 Claussen P 1912 *Z Botan* 4 1-64
- 82 Coons G H 1916 *J Agr Research* 5 713-769
- 83 Cooper D C and C L Porter 1938 *Phytopathology* 18 881-899
- 84 Couch J N 1939 *Am J Botany* 26 119-130
- 85 Darby R T and G R Mandels 1957 *Plant Physiol* 30 360-366

species are usually not alike so that in a sense the two types compete. Apart from the relatively trivial generalization that the larger sexual structures must require more energy for their formation than the (usually) smaller asexual organs there is as yet no good evidence for a universal difference between sexual and asexual reproduction this possibility however should surely be explored. A few instances have been noted in which two different asexual structures have different requirements.

BIBLIOGRAPHY

- 1 Allington W B 1936 *Phytopathology* 26 831-844
- 2 Ames A 1915 *Phytopathology* 5 11-19
- 3 Aronescu A 1933 *Mycologia* 25 43-54
- 4 Aronescu A 1934 *Mycologia* 26 244-249
- 5 Aschan K 1934 *Physiol Plantarum* 7 571-591
- 6 Asthana R P and L E Hawker 1936 *Ann Botany* 50 325-343
- 7 Ayres T T 1911 *Mycologia* 33 148-187
- 8 Backus M P 1937 *Mycologia* 29 333-336
- 9 Backus M P 1939 *Bull Torrey Botanical Club* 66 63-76
- 10 Badcock E C 1943 *Trans Brit Mycol Soc* 26 127-137
- 11 Bailey A A 1932 *Botan Gaz* 94 225-271
- 12 Bajaj V S P Damle and F S Krishnan 1954 *Arch Biochem Biophys* 50 451-460
- 13 Baker R E D 1931 *New Phytologist* 30 303-316
- 14 Banbury G H 1952 *J Exp Botany* 3 77-85 86-94
- 15 Banbury G H 1955 *J Exp Botany* 6 235-244
- 16 Barnett H L and V G Lilly 1947a *Am J Botany* 34 196-204
- 17 Barnett H L and V G Lilly 1947b *Mycologia* 39 699-708
- 18 Barnett H L and V G Lilly 1950 *Phytopathology* 40 80-89
- 19 Barnett H L and V G Lilly 1955 *Mycologia* 47 26-29
- 20 Barnett H L and V G Lilly 1956 *Mycologia* 48 617-627
- 21 Basu S N 1951 *J Gen Microbiol* 5 231-238
- 22 Basu S N 1952 *J Gen Microbiol* 6 199-204
- 23 Baumann G 1953 *Kuhn Arch* 67 305-383
- 24 Benedek T 1943 *Mycologia* 35 222-242
- 25 Bennett C W 1921 *Mich State Univ Agr Exp Sta Tech Bull* 53 1-40
- 26 Bertrand D and A De Wolf 1955 *Compt rend (Paris)* 241 1877-1880
- 27 Biffen R H 1898 *J Linnean Soc London Botany* 51 147-174
- 28 Bille Hansen E 1953 *Physiol Plantarum* 6 523-528
- 29 Bisby G R 1923 *Mycologia* 17 89-97
- 30 Bishop H 1940 *Mycologia* 32 505-529
- 31 Buxis G 1956 *Am J Botany* 43 389-391
- 32 Blaauw A H 1914 *Z Botan* 6 641-653
- 33 Blodgett E W 1936 *Phytopathology* 26 115-152
- 34 Bonner J T 1947 *J Exp Zool* 106 1-26
- 35 Bonner J T, W W Clarke Jr, C L Neely Jr and M K Shukin 1950 *J Cellular Comp Physiol* 36 149-158
- 36 Bonner J T, K K Kane and R H Levey 1956 *Mycologia* 48 13-19
- 37 Borris H 1954 *Planta* 22 644-684

- 33 Bortels H 1927 *Biochem Z* 182 301-328
- 39 Bose S R 1943 *Mycologia* 32 33-46
- 40 Brandt W H 1923 *Mycologia* 45 194-208
- 41 Brefeld O 1877 *Botan Ztg* 32 386
- 42 Bretzlaff C W Jr 1954 *Am J Botany* 41 58-67
- 43 Brewer D and B H MacNeill 1953 *Can J Botany* 31 79-144
- 44 Brian P W and H G Hemming 1950 *Trans Brit Mycol Soc* 33 13-141
- 45 Brown W 1925 *Ann Botany* 39 373-408
- 46 Brucker W 1955 *Arch Protistenk* 100 39-50
- 47 Buchheim A 1928 *Ber deut botan Ges* 46 167-180
- 48 Buder J 1918 *Ber deut botan Ges* 36 103-10
- 49 Buller A H R 1909 *Researches on Fungi* Vol 1 London Longmans Green and Co pp 37
- 50 Buller A H R 1910 *Researches on Fungi* Vol 2 London Longmans Green and Co pp 492
- 51 Buller A H R 1914 *Researches on Fungi* Vol 3 London Longmans Green and Co pp 611
- 52 Buller A H R 1915 *Researches on Fungi* Vol 5 London Longmans Green and Co pp 416
- 53 Buller A H R 1934 *Researches on Fungi* Vol 6 London Longmans Green and Co pp 513
- 54 Bunn E E 1937a *Planta* 26 719-736
- 55 Bunning E 1937b *Planta* 27 583-610
- 56 Burgeff H 1904 *Botan Abh Heft* 4 3-13
- 57 Burnett J H 1913 *New Phytologist* 5 58-64 86-88
- 58a Burnett J H 1926 *New Phytologist* 55 42-49
- 58 Burt E A 1897 *Botan Gaz* 24 73-92
- 59 Buston H W and S N Basu 1948 *J Gen Microbiol* 2 162-171
- 60 Buston H W and A H Khan 1926 *J Gen Microbiol* 14 622-660
- 61 Buston H W and B Rickard 1916 *J Gen Microbiol* 12 194-197
- 62 Camp W C 1937 *Am J Botany* 24 300-303
- 63 Cantino E C 1949 *Am J Botany* 36 92-112
- 64 Cantino E C 1916 *Mycologia* 48 222-240
- 65 Cantino E C and M T Hyatt 1953 *J Bacteriol* 66 712-720
- 66 Carlile M J 1926 *J Gen Microbiol* 14 613-624
- 66a Carlile M J 1957 *Nature (London)* 180 202
- 67 Carvajal F 1947 *Mycologia* 39 426-440
- 68 Castle E S 1929 *J Gen Physiol* 12 391-400
- 69 Castle E S 1931 *J Gen Physiol* 14 703-711
- 70 Castle E S 1933 *J Gen Physiol* 15 481-489
- 71 Castle E S 1933 *J Gen Physiol* 15 41-6
- 72 Castle E S 1935 *Cold Spring Harbor Sympos Quant Biol* 3 21-29
- 73 Cayley D M 1931 *J Genet* 24 1-63
- 74 Charlton K M 1913 *Trans Brit Mycol Soc* 36 343-325
- 75 Chaudhuri H 1923 *Ann Botany* 37 519-539
- 76 Cherewick W J 1944 *Can J Research C* 2 52-82
- 77 Chilton S J P 1943 *Mycologia* 3 15-20
- 78 Chona B L 193 *Trans Brit Mycol Soc* 17 1-8
- 79 Chowdhury S 1914 *J Indian Bot n Soc* 23 91-106
- 80 Christenberry G A 1938 *J Elisha Mitchell Sci Soc* 54 297-310
- 81 Claussen P 1911 *Z Botan* 4 1-64
- 82 Coons G H 1916 *J Ag Research* 5 713-69
- 83 Cooper D C and C L Potter 1918 *Phytopathology* 18 881-899
- 84 Couch J N 1939 *Am J Botany* 6 119-130
- 85 Dauby R T and G R Mandels 1913 *Plant Physiol* 30 360-366

- 86 Deibrock M and W Reichardt 1956 In D Rudnick (ed) *Cellular Mechanisms in Differentiation and Growth* p 3-44 Princeton Princeton Univ Press
- 87 Denny F E 1933 *Contribs Boyce Thompson Inst* 5 90-107
- 88 Diener U L 1935 *Phytopathology* 45 141-143
- 89 Dillon Weston W & R 1936 *Trans Brit Mycol Soc* 40 112-115
- 90 Dimock A W and J H Osborn 1913 *Phytopathology* 33 317-331
- 91 Doran W L 1921 *Mass Agr Exp Sta Bull* 207 39-66
- 92 Driver C H and H E Wheeler 1955 *Mycologia* 47 311-316
- 93 Elarosi H 1936 *Nature (London)* 177 660-666
- 94 Ellis M 1931 *Trans Brit Mycol Soc* 16 102-114
- 95 Errera L 1885 *Mém Cour Acad Roy Belg* 37 (3) 1-50
- 96 Falck R 1901 *Beitr Biol Pflanz* 8 213-306
- 97 Fikry A 1932 *Ann Botany (London)* 46 29-70
- 98 Fischer E and E Gaumann 1929 *Biologie der pflanzenbewohnenden parasitischen Pilze* Jena Gustav Fischer pp 428
- 99 Flint L A 1949 *Am J Botany* 29 672-674
- 100 Flugge H 1939 *Lorratspflege u Lebensmittelforsch* 2 237-256 (*Zentr Bakteriell Parasitenk Abt II* 102 401 1910)
- 101 Foster J W 1939 *Botan Rev* 5 207-239
- 102 Foster J W L E McDaniel H B Woodruff and J L Stokes 1945 *J Bacteriol* 50 355-368
- 103 Fox A S and W O Gray 1930 *Proc Natl Acad U S* 36 538-546
- 104 Frank M C C T Calam and P H Gregory 1918 *J Gen Microbiol* 9 10-19
- 105 Fulkerson J F 1935 *Phytopathology* 45 22-23
- 106 Galleymore H B 1919 *Trans Brit Mycol Soc* 39 315-317
- 107 Galston A W 1930 *Botan Rev* 16 361-378
- 108 Gassner G and W Franke 1938 *Phytopathol Z* 11 517-570
- 109 Geach W L 1936 *J Council Sci Ind Research (Australia)* 9 77-87
- 110 Gehleno P M 1944 *Biodynamica* 4 359-368
- 110a Gibson I A S and J Trapnell 1957 *Trans Brit Mycol Soc* 40 213-220
- 111 Gubardt M 1932 *Flora N F* 39 477-525
- 112 Goldring D 1936 *Ann Missouri Botan Garden* 23 527-541
- 113 Gollmick F 1936 *Zentr Bakteriell Parasitenk Abt II* 93 421-419
- 114 Graf Marin A 1934 *Cornell Univ Agr Exp Sta Mem* 157 1-48
- 115 Grainger J 1946 *Trans Brit Mycol Soc* 29 52-61
- 116 Gray W D 1939 *Am J Botany* 26 709-711
- 117 Gray W D 1933 *Mycologia* 45 817-821
- 118 Gregg J H 1930 *J Exp Zool* 114 113-196
- 119 Gregory P H 1939 *Trans Brit Mycol Soc* 23 24-51
- 120 Greis H 1937 *Botan Arch* 38 113-151
- 121 Grossbard E 1954 *Phytopathology* 44 110-111
- 122 Grosser A H Kundtner Schwarzkopf and K Bernhauer 1950 *Arch Mikrobiol* 15 247-252
- 123 Hafiz A 1931 *Trans Brit Mycol Soc* 34 259-269
- 124 Hall M P 1933 *Ann Botany* 47 543-518
- 125 Hammarlund J C 1925 *Hereditas* 6 1-126
- 126 Hampson C R 1934 *J Bacteriol* 67 39-740
- 127 Hansen H N 1938 *Mycologia* 30 447-455
- 128 Hansen H N and W C Snyder 1911 *Science* 33 264-265
- 129 Hansen H N and W C Snyder 1917 *Phytopathology* 37 369-371
- 130 Harter L I 1939 *Am J Botany* 26 234-243
- 131 Hatch W R 1938 *Mycologia* 28 439-441
- 132 Hawker L E 1930 *Ann Botany* 50 699-717
- 133 Hawker L F 1932a *Ann Botany* 3 45-463

- 134 Hawker L E 1939b *J Botany* 3 651-675
- 135 Hawker L E 1942 *Ann Botany* 6 631-636
- 136 Hawker L E 1941 *Ann Botany* 8 19-30
- 137 Hawker L E 1947a *Ann Botany* 11 245-260
- 138 Hawker L E 1947b *Nature (London)* 159 136
- 139 Hawker L E 1948 *Ann Botany* 12 77-9
- 140 Hawker L E 1950 *Physiology of Fungi* London University of London Press Ltd pp 360
- 141 Hawker L E and S D Chaudhuri 1946 *Ann Botany* 10 185-194
- 142 Hawn E J and T C Vanterpool 1953 *Can J Botany* 31 699-710
- 143 Hazen E L 1947 *Mycologia* 39 200-209
- 144 Hazen E L 1951 *Mycologia* 43 284-296
- 145 Hedgecock G C 1906 *Ann Rept Missouri Botanical Garden* 17 115-117
- 146 Heim I 1930 *Am J Botany* 17 143-151
- 147 Heintzeler I 1939 *Arch Mikrobiol* 10 90-113
- 148 Henriksson L E and J F Morgan Jones 1951 *Svensk Botanisk Tidskrift* 4 618-65
- 149 Henry B W and A L Anderson 1948 *Phytopathology* 38 65-8
- 150 Hesselstine C W and P Anderson 1956 *Am J Botany* 43 696-703
- 150a Hesselstine C W and R F Anderson 1957 *Mycologia* 49 419-459
- 151 Heyn A N J 1953 *Proc Koninklijke Akademie Wetenschap Amsterdam* 58 1014-1081
- 152 Hickey R J and H D Tresner 1959 *J Bacteriol* 64 891-897
- 153 Hirsch H M 1954 *Physiol Plantarum* 7 2-97
- 154 Hodapp E L 1949 *Biodynamica* 4 33-46
- 155 Hofmann E H Scheck and K Saffert 1950 *Biochem Z* 50 196-198
- 156 Hopkins E F 1959 *Phytopathology* 49 390-393
- 157 Horne A S and J H Mitter 1957 *Ann Botany* 41 519-547
- 158 Horsfall J G and S R Ch 1955 *Trans N Y Acad Sci* 18 69-80
- 159 Houston B R and J W Oswald 1946 *Phytopathology* 36 1049-1055
- 160 Hutchinson A H and M R Ashton 1930 *Can J Research* 3 181-198
- 161 Ingold C T 1933 *New Phytologist* 32 118-196
- 162 Ingold C T 1946 *Trans Bot Mycol Soc* 29 43-1
- 163 Ingold C T 1953 *Dispersal in Fungi* Oxford Clarendon Press pp 197
- 164 Jacobs S E and A W Marsden 1948 *Ann Appl Biol* 35 18-24
- 165 Javille M 1913 *Ann Inst Pasteur* 97 1091-1039
- 166 Jump J A 1954 *Am J Botany* 41 561-6
- 167 Kehl H 1937 *Arch Mikrobiol* 8 319-406
- 168 Kerl I 1931 *Z Botan* 31 109-174
- 169 Kilpatrick R A and H W Johnson 1956 *Phytopathology* 46 180-181
- 170 Kirchhoff H 1959 *Centr Bakteriol Parasitenk Abt II* 77 310-369
- 171 Klebs G 1896 *Ueber die Fortpflanzungsphysiologie der niederen Organismen der Protobionten Die Bedingungen der Fortpflanzung bei einzelligen und Pilzen* Jena Gustav Fischer pp 543
- 172 Klebs G 1898 *Jahrb wiss Botan* 32 1-10
- 173 Klebs G 1899 *Jahrb wiss Botan* 33 513-93
- 174 Klebs G 1900 *Jahrb wiss Botan* 35 80-103
- 175 Klein D T 1948 *Botany* 36 110 139-141
- 176 Kohler F 193 *Planta* 93 38-377
- 177 Krafczyk H 1931 *Ber deut bot Ges* 49 141-146
- 178 Krafczyk H 1935 *Beitr Biol Pflanz* 23 349-396
- 179 Kunkel L O 1918 *Brooklyn Botanical Garden Mem* 1 306-312
- 180 Kusano S 1930 *Japan J Botany* 5 35-13
- 181 Lambert E B 1933 *J Agr Research* 47 599-608
- 182 Lambert E B 1938 *Botan Rev* 4 397-466

- 183 Lehman S G 1923 *Ann Missouri Botanical Garden* 10 111-118
- 184 Leonian L H 1923 *Phytopathology* 13 207-212
- 185 Leonian L H 1924 *Am J Botany* 11 19-50
- 186 Leonian L H and V G Lilly 1937 *Am J Botany* 24 700-709
- 187 Levine M N 1923 *J Agr Research* 21 539-567
- 188 Levine M N 1928 *Phytopathology* 18 7-123
- 189 Lieth H 1934 *Ber deut botan Ges* 67 323-325
- 190 Lieth H 1936 *Arch Mikrobiol* 24 91-104
- 191 Lindegren C C and E Hamilton 1914 *Botan Ga* 10 316-371
- 192 Lockwood L B 1937 *Mycologia* 29 280-290
- 193 Lohmus M P 1911 *Antonie van Leeuwenhoek J Microbiol Serol* 10 100-122
- 194 Longtree K 1939 *Cornell Univ Agr Exp Sta Mem* 223 1-43
- 195 McCallan S E A and S Y Chan 1944 *Contribs Boyce Thompson Inst* 13 323-331
- 196 McVickar D L 1942 *Am J Botany* 29 312-330
- 197 Madelin M F 1936a *Ann Botany* 20 307-330
- 198 Madelin M F 1936b *Ann Botany* 20 461-480
- 199 Mader E O 1943 *Phytopathology* 33 1134-1145
- 200 Magie R O 1935 *Phytopathology* 25 131-159
- 201 Mann M L 1932 *Bull Torrey Botanical Club* 59 443-490
- 202 Markert C L 1919 *Am Naturalist* 83 227-231
- 203 Marsh P B and K Bollenbacher 1916 *Am J Botany* 33 240-249
- 204 Mathur R S H L Barnett and V G Lilly 1930 *Phytopathology* 40 104-114
- 205 Metz O 1930 *Arch Mikrobiol* 1 197-251
- 206 Milburn T 1904 *Centr Bakteriell Parasitenk Abt II* 13 129-138 257-216
- 207 Miller J J 1915 *Can J Research C* 23 16-43
- 208 Mix A J 1933 *Phytopathology* 23 503-504
- 209 Molliard M 1920 *Compt rend (Paris)* 170 919-921
- 210 Morgenthau O 1910 *Centr Bakteriell Parasitenk Abt II* 27 73-90
- 211 Mulder E G 1939 *Arch Mikrobiol* 10 12-96
- 212 Munk M 1912 *Centr Bakteriell Parasitenk Abt II* 32 353-375
- 213 Nagel C M 1934 *Phytopathology* 24 1101-1110
- 214 Nickerson W J and Z Mankowski 1933a *Am J Botany* 40 584-591
- 215 Nickerson W J and Z Mankowski 1933b *J Infectious Diseases* 5 20-25
- 216 Nickerson W J and K V Thumann 1941 *Am J Botany* 28 617-621
- 217 Nickerson W J and K V Thumann 1913 *Am J Botany* 30 91-101
- 218 Nitumarga N M 1937 *Ann Botany* 49 19-40
- 219 Obel P 1910 *Ann Mycol* 8 421-443
- 220 Pady S M 1918 *Mycologia* 40 21-33
- 221 Page R M 1936 *Mycologia* 48 206-224
- 222 Parr R 1918 *Ann Botany* 30 177-205
- 223 Perlman D R B Greenfield Jr and E O'Brien 1934 *Appl Microbiol* 2 199-202
- 224 Pillai N C and K S Srinivasan 1936 *J Gen Microbiol* 14 248-251
- 225 Plimpel M 1937 *Arch Mikrobiol* 26 151-174
- 226 Plunkett B E 1933 *Ann Botany* 17 193-217
- 227 Plunkett B E 1936 *Ann Botany* 20 533-586
- 228 Powlesland R 1934 *Trans Brit Mycol Soc* 37 362-371
- 229 Prince A E 1913 *Farfousa* 1 79-93
- 230 Rabinowitz Sereis D 1933 *Bol R Stat Feg* 13 203-206 (*Biol Abstr* 10 1 992 1936)
- 231 Ram C S V 1939 *Nature (London)* 170 839
- 232 Ramsey G B and A A Bailey 1930 *Botan Ga* 89 115-136

- 233 Rands R D 1917 *Phytopathology* 7 316-317
- 234 Raper J R 1939 *Am J Botany* 6 639-650
- 235 Raper J R 1940 *Bot: Ga* 112 1-74
- 236 Raper J R 195 *Botan Rev* 18 447-545
- 237 Raper J R and A J Haaen Smit 1942 *J Biol Chem* 143 311-370
- 238 Raper K B 1956 *Mycologia* 48 169-05
- 239 Raper K B and C Thom 1944 *Mycologia* 36 555-575
- 240 Reidemeister W 1909 *Ann Mycol* 7 19-44
- 41 Reinert J 1957 *Naturwiss* 31 47-48
- 242 Richards M C 1949 *Phytopathology* 39 20
- 43 Robak H 1941 *Nyt Mag Naturv* 81 105-116 (*Rev Appl Mycol* 25 331 1946)
- 244 Robbins W J and I McVeigh 1949 *Mycologia* 41 141-170
- 45 Robbins W J and M B Schmitt 1949 *Am J Botany* 32 3 0-3 6
- 246 Roberg M 19 8 *Centr Bakt Parasitenk Abt II* 74 333-370
- 47 Robinson R 1926 *Trans Brit Mycol Soc* 10 307-314
- 248 Robinson W 19 6 *Ann Botany* 40 245-279
- 249 Roers C H 1933 *J Agr Research* 58 701-709
- 50 Ronsdorf L 1931 *Planta* 14 48 -514
- 250a Sagromsky M 1952 *Flora N F* 39 300-313 560-564
- 251 Salvin S B 1942 *Am J Botany* 29 97-104
- 252 Sanford G B 1946 *Phytopathology* 46 281-284
- 253 Schenck E 1919 *Beih Botan Centr* 36 Abt I 3 5-413
- 254 Schmidle A 1951 *Arch Mikrobiol* 16 80-100
- 255 Schneiderhohn G 1955 *Arch Mikrobiol* 21 230-236
- 256 Schonborn W 1955 *Arch Mikrobiol* 408-431
- 257 Schopfer W H 1928 *Bull soc botan Genève* 20 149-323
- 258 Schopfer W H 1935 *Compt rend soc phys et hist nat Genève* 50 152-154
- 259 Schopfer W H and S Blumer 1940 *Protoplasma* 34 574-532
- 260 Shaffer B M 1956 *Science* 123 117 -1173
- 261 Shanor L 1936 *J Elisha Mitchell Sc Soc* 57 99 104
- 26 Smith E C 1935 *Bull Torrey Botan Club* 67 151-164
- 263 Smith E C 1936 In B M Dugar (ed) *Biological Effects of Radiation* Vol 9 p 889-918 New York McGraw Hill Book Co
- 64 Snyder W C and H N Hansen 1941 *Mycologia* 33 580-591
- 465 Snyder W C and H N Hansen 1947 *Phytopathology* 37 470-421
- 66 Sorgel G 1953 *Arch Mikrobiol* 19 372-397
- 267 Sost H 1955 *Arch Protistenk* 100 541-564
- 268 Stakman E C and M N Levine 1919 *J Agr Research* 16 43-77
- 269 Steinberg R A 1935 *Bull Torrey Botan Club* 67 81-90
- 70 Steinberg R A 1936 *Am J Botany* 23 277-231
- 271 Stevens F L 19 8 *Botan Gaz* 86 210-2 5
- 72 Stevens F L 1930a *Am J Botany* 17 8 0-881
- 273 Stevens F L 1930b *Zentr Bakteriell Parasitenk Abt II* 82 161-174
- 274 Stevens F L 1931 *Mycologia* 23 134-139
- 475 Stiefel S 1957 *Planta* 40 301-312
- 276 Stockdale P M 1943 *J Gen Mikrobiol* 8 434-441
- 277 Stoll A 1936 *Zentr Bakteriell Parasitenk Abt II* 93 296-298
- 278 Streeter S G 1909 *Botan Gaz* 48 414-476
- 479 Susan M 1946 *Ann Rev Microbiol* 10 71-50
- 280 Sussman M F Lee and N S Kerr 1946 *Science* 103 1171-1172
- 281 Sweet H R 1941 *Am J Botany* 28 150-161
- 28 Takemaru T 1944 *Bot n Mag (Tokyo)* 67 82-86
- 283 Ternetx C 1900 *Jahrb wiss Botan* 35 273-312

- 284 Thren R 1937 *Z Botan* 31 331-391
- 285 Timmick M B H L Barnett and V G Lilly 1952 *Mycologia* 44 141-149
- 286 Timmick M B V G Lilly and H L Barnett 1951 *Phytopathology* 41 327-336
- 287 Tisdale W B 1923 *J Agr Research* 21 5, 86
- 288 Turian G 1952 *Experientia* 8 309
- 289 Turian G and F T Haxo 1954 *Botan Gaz* 115 254-260
- 290 Tyner L E and G B Sanford 1935 *Sci Agr* 16 197-207
- 291 Uebelmesser E R 1954 *Arch Mikrobiol* 20 1-33
- 292 Utiger H 1953 *Naturwiss* 40 292-293
- 293 Vandendries R 1931 *Compt rend (Paris)* 198 193-195
- 294 Voderberg K 1950 *Planta* 37 612-625
- 295 Wagner F 1955 *Arch Mikrobiol* 22 313-323
- 296 Wakefield E M 1910 *Naturw Z Land u Forstwirtschaft* 7 521-551
- 297 Wald G 1943 *Vitamins and Hormones* 1 195-297
- 298 Walker L B 1927 *J Elisha Mitchell Sci Soc* 42 151-178
- 299 Walker L B and E N Andersen 1925 *Mycologia* 17 154-159
- 300 Walter H 1921 *Z Botan* 13 673-718
- 301 Waters C W 1928 *Phytopathology* 18 157-213
- 302 Webster J 1952 *New Phytologist* 51 229-235
- 303 Weimer J L and L L Harter 1923 *J Agr Research* 24 1-10
- 304 Westergaard M and H M Hirsch 1954 *Proc Symposium Colston Research Soc* 7 p 171-181 New York Academic Press
- 305 Westergaard M and H K Mitchell 1947 *Am J Botany* 34 573-577
- 306 Weston W H Jr 1923 *J Agr Research* 23 239-248
- 307 Weston W H Jr 1924 *J Agr Research* 27 771-784
- 308 Wheeler H E 1954 *Phytopathology* 44 312-315
- 309 Wheeler H E 1956 *Mycologia* 48 349-354
- 310 Wilhelm S 1947 *Mycologia* 39 716-724
- 311 Williams A M and E McCoy 1953 *Appl Microbiol* 1 307-313
- 312 Witsch H von and F Wagner 1955 *Arch Mikrobiol* 22 307-317
- 313 Wolff L K and A Emmerie 1950 *Biochem Z* 298 413-450
- 314 Yarwood C E 1936 *J Agr Research* 52 645-657
- 315 Yarwood C E 1937 *J Agr Research* 54 365-373
- 316 Yarwood C E 1941 *Am J Botany* 28 353-357
- 317 Yarwood C E 1943 *Hilgardia* 14 595-691
- 318 Yarwood C E 1956 *Plant Disease Repr* 40 318-321
- 319 Yu C C 1951 *Am J Botany* 41 21-30
- 320 Zachariah A T H N Hansen and W C Snyder 1956 *Mycologia* 48 49-467
- 321 Zalekar M 1954 *Arch Biochem Biophys* 50 71-80
- 322 Zentmyer G A 1952 *Phytopathology* 42 24

12. Spore Germination

Spores—specialized self contained microscopic structures capable of initiating new growth—are the principal agent of dispersal of fungi. Consequently any consideration of ecology or of the spread of economically important fungi must take spore germination into account. The control of plant diseases by protectant fungicides is in essence the problem of inhibiting the germination of spores. And the transformation of a relatively inactive structure into an actively growing thallus poses some fundamental problems of interest to biology as a whole.

In this chapter the several cytological and morphological types of spore have been distinguished in the discussion only if some important physiological aspect is specific to the type. In the long run we may suppose such a practice—treating the rust fungus teliospore and the myxomycete swarmspore alike—will prove indefensible. For the present however the available information does not permit separate treatment of the different spore forms except in such special cases as the dormant resting spores.

It will be seen that as usual in the fungi the corpus of knowledge is seriously out of balance. We find literally hundreds of studies of temperature and water relations of the economically important fungi but such groups as the aquatic fungi the fleshy basidiomycetes and the actinomycetes are virtually unknown in this dimension. More serious still concentration on well known and easily determined environmental factors has meant relative neglect of the fundamental metabolic

processes which control germination and of the physiological and biochemical changes consequent upon germination. It is reasonable to expect that investigations into these and related topics would in turn throw light on some of the more obscure environmental and ecological problems in spore physiology.

1 MEASUREMENT

The most commonly used criterion of spore germination is the fraction of spores that in a given time forms a germ tube. The time period is usually chosen so as to allow all viable spores to germinate. Spores are sown on water on agar or on nutrient media incubated under controlled conditions and counted microscopically or the spores may be germinated in an aerated or shaken liquid medium (65). Self-inhibition and other associative effects (p. 413) make it imperative that the concentration of spores be held constant and be known. It is usually convenient to bring the spore suspension to a known turbidity measured optically and previously standardized against a direct microscopic count (240). Spore concentration can also be estimated by capillary volume (192-289). Clumping of spores introduces error and must therefore be minimized (130).

The rate of germination may be determined by periodic counts during the experiment and often provides valuable supplementary information.

A second major criterion is the latent period of germination, the time required for germination to begin or to reach some specified low value (25-318-287). Limited evidence suggests that this is fundamentally different from total germination, in that temperature affects it differently (p. 410) and toxicants may exert greater or less effect on the latent period than on total germination (287).

The rate of increase in the length of the germ tube is again a different criterion of spore activity. In general it is more sensitive to temperature than is total germination, responding more like mycelial growth; similarly the pH range for optimum germ tube growth in *Alternaria solani* is rather narrower than that for best total germination (159). The rate of elongation can be expressed as a velocity constant (126).

No one of these criteria can be said to be best. The method used must take account of the problem which is under investigation. Fundamental studies on the germination process, for example, might make use of the rate of germination and the total germination. On the other hand, an ecological or epidemiological problem might be served better

by measurements of germ tube growth. Infection of a plant by a pathogen is dependent usually on at least some extension of the germ tube; if this process is more sensitive to unfavorable conditions than is total germination, a false picture of the potentialities of the organism will be deduced from observations on total germination alone. Unfortunately, germ tube measurements are rather more laborious and are perhaps avoided for that reason; a rapid semi-automatic method of determining germ tube lengths should not be beyond the capacity of modern instrumentation and would be a real contribution to the study of fungi.

2 THE GERMINATION PROCESS

Just as we have had to lump together the various morphological types of fungus spore, it is also necessary, with some exceptions, to consider the several germination types as if they were all alike. Of these, the most common and the most frequently studied is formation of one or more germ tubes which grow indeterminately and develop into a mycelium in culture or in a parasitized plant or animal. Among the phycomyces, several types of spore germinate by the formation of smaller endogenous spores; i.e., the parent structure is functionally a sporangium. In *Phytophthora infestans* the same structure—perhaps best called a conidiosporangium—germinates either by a tube or by formation of swarmspores, depending upon the environment, particularly temperature, and perhaps on the age and previous history of the structure (19, 63, 279, 291). The chlamydospores of the smut fungi germinate somewhat differently under different external conditions of temperature (140), concentration (132), and nutrients (264). Basidio-spores of *Rhizoctonia solani* form secondary spores on water agar, germ tubes on a nutrient agar (131). The macroconidium of *Thielaviopsis paradoxa* similarly germinates very differently on water and on nutrient agar (149).

Multicellular spores often germinate by the simultaneous formation of two or more germ tubes (179) in *Pestalotia* spp.; however, the basal cell normally gives rise to the germ tube (300), and germination in *Helminthosporium sativum* may be either polar or bipolar (137).

Conidia of the higher fungi may have one, two, or several nuclei at the time of their formation; a few examples may be cited:

1. Predominantly uninucleate: *Aspergillus* spp. (8, 12), *Colletotrichum destructivum* (49), *Frichoderma viride* (12), *Pteridillum albo-atrum* (128).

2 Predominantly multinucleate *Aspergillus echinulatus* (8) *Botrytis cinerea* (128) *Helminthosporium sativum* (34 137) Sporangiospores of *Rhizopus nigricans* are uninucleate (12) those of *Mucor disseisus* multinucleate (68)

Nuclear condition is of particular importance in genetic studies since a heterokaryotic mycelium which produces uninucleate conidia—more accurately conidia from uninucleate conidiophores—essentially regenerates the parent types in the absence of gene exchange (112) By extension of this principle it is possible to infer from genetic evidence in *Streptomyces* spp. that the conidia are uninucleate or arise from uninucleate cells (28)

The number of nuclei in macroconidia of *Neurospora crassa* is reduced by cultivation in a minimal medium (138) or in a sorbose medium (5) The nuclei of the higher fungi are of course normally haploid Roper (236) has however succeeded in inducing the formation of diploid conidia in *Aspergillus nidulans* and camphor treatment of *Neurospora crassa* induces the formation of possibly diploid cells (243)




Figure 1 The time course of spore germination (probit scale) in *Monilinia fructicola* (curves 1) and *Alternaria solani* (curves 2) Arithmetic function of time plotted in 1 reciprocal function in P Redrawn from Wellman and McCallan (299) by permission of the Poyce Thompson Institute for Plant Research Inc

A given population of spores germinates over a period of time if the population is standardized as to age there is good evidence that random variability is the cause of the spread in time of germination. As shown in Figure 1 a plot of per cent germination on a probit scale against the reciprocal of time is linear (provided a correction is made for non viable spores) i.e. time of germination is normally distributed. Similar data have been reported in studies of *Neurospora crassa* (240). The over all rapidity of germination is a species characteristic. *Melampsora lini* begins germination in less than one hour (129) but *Phoma apicola* spores require 24-48 hours (13). Other data on rapidity of germination are reviewed by Gottlieb (113) and by Doran (73).

The three distinguishing morphological events of typical spore germination are nuclear division (8-12) swelling of the spore and emergence of the germ tube. Swelling with one significant exception appears to be universal in fungi and actinomycetes having been described in sporangiospores (41) ascospores (121) basidiospores (32-135) and conidia (25-114-125-192-255-260a-306). The exception—apart from the fact that observations on *Sclerotinia fructicola* conflict (192-306)—is found among the powdery mildew fungi: conidia of these do not swell prior to germination (306). The significance of swelling in relation to water requirements will be considered later.

There is scattered evidence that physiological changes also occur during the germination process although the data cannot be generalized to all fungi. Anion toxicity is more marked in the early stage of germination of the resistant sporangium of *Blastocladiella* sp. (45) than in a later stage: oxygen uptake by ascospores of *Neurospora tetrasperma* becomes more sensitive to iodoacetamide after appearance of the germ tube (109) and in the same material cation permeability appears to change at the beginning of germination (274). Other evidence of separable physiological states during the germination process appears in studies of oxygen uptake and of temperature responses.

Germ tube growth presumably follows the same rules as any other mycelial growth. Extension occurs at the tip only (259-263). For at least some time growth is exponential i.e. the logarithm of germ tube length increases linearly with time (220-259) germ tube growth in *Ustilina vulgaris* however is reported to follow a different course (301).

Germ tube tropisms toward water and toward nutrients have been reported but the evidence is conflicting and inconclusive (9-55-100-117-203). It is not in particular necessary to postulate a chemotropic reaction in the penetration of plant pathogens into host tissue (39-40).

62) Here an unspecified contact stimulus thigmotropism is supposed to operate. The possibility of chemotropism has been reopened more recently in studies on *Saprolegnia ferax* (87).

One type of germ tube tropism is well documented—the negative tropism of germ tubes toward each other (55 100 117). As described by Stadler (263) germ tubes of *Rhizopus nigricans* form on the side of the spore distant from other spores and in addition actually bend away from other germ tubes during their growth. The stimulus is almost certainly chemical and it is tempting to consider an auxinlike growth factor in explanation of the bending phenomenon. This possibility is also suggested by the finding that germ tubes of rust fungi are longer when germinating spores are crowded than when they are more dispersed (83 314). However Stadler (263) presents some evidence against an explanation based on growth factors; instead he postulates an unstable inhibitor which has the effect of thickening or strengthening the wall of the spore or germ tube. The last word on this problem has clearly not been spoken.

Germ tubes of sporidia and of uredospores of *Puccinia* spp. react negatively to strong light; studies with filters indicate that the active wavelength is in the blue region of the visible spectrum (90 98 235 266). It will be recalled from Chapter 11 that it is these shorter wavelengths which are most active in phototropic and inductive effects on reproductive structures. Negative phototropism is also shown by conidia of powdery mildew fungi (204 305).

3 SPORE METABOLISM

Water, minerals, carbohydrates, and other constituents of fungus spores have been mentioned incidentally in Chapter 2. In relation to spore germination the important materials are water—considered elsewhere in this chapter—and probably lipids. It should also be recalled that spores of *Aspergillus niger* are relatively high in total phosphorus, a part of which is in organic combination (7 154).

Although the fat content of most spores is not unusually high (248 271), three bits of evidence suggest that fats provide energy for the germination of spores which do not require an exogenous source of carbon. First, microscopic observation shows that stainable lipids disappear during germination (82 150). Second, the respiratory quotient of *Neurospora sitophila* microconidia is about 0.7 (208), a value characteristic of, but of course not sufficient evidence for, fat oxidation. Finally, the data of Table 1 show that at least in high fat spores the principal organic material disappearing during germination is the

Table 1 Chemical Changes during Germination of Uredospores of *Puccinia graminis tritici**

Fraction	Amount Per Cent of Dry Weight	
	Before Germination	After Germination
Carbohydrate	21.9	22.4
Chitin	1.4	3.4
Protein	25.9	22.3
Lipid	19.7	7.8

* From Shu Tanner and Ledingham (254) by permission of the *Canadian Journal of Botany*

lipid fraction. The conidia of *Erysiphe graminis* like the rust spores of Table 1 appear to be relatively high in fat (310).

During the germination of spores of *Aspergillus nidulans* the most pronounced changes which occur in the first few hours are decline in polymetaphosphate and in organically bound phosphate, disappearance of part of the free amino acids of the spore and a progressive fall in total lipid (253).

Several hydrolytic enzymes—acting on starch, oligosaccharides, glucosides, pectic substances, proteins and urea—occur in crude spore preparations; the early literature is reviewed by Mandels and Norton (195). Spores of *Myrothecium verrucaria* have been found to possess an enzymatic system acting on ascorbic acid (185, 186), another system oxidizing sulphydryl compounds to disulfides (191) and one or possibly two enzymes acting on sucrose (184, 187, 188). The invertase of *M. verrucaria* is synthesized rapidly during germination and appears in the medium almost immediately (190). These enzymes are inactivated by an acid treatment which does not kill the spore or damage its respiration; from this fact it has been proposed that they are located at or near the spore surface (184, 185). The diphosphopyridine nucleotidase of *Neurospora crassa* is produced primarily or entirely by the conidia and is so easily washed out of the cell that it too may have a surface location (309).

Although there is as yet no reason to think that spore respiration differs fundamentally from that of mycelium, studies of the general problem should contribute to our understanding of both spore germination and fungicidal action. Data available at present are too fragmentary to support any firm theoretical structure or even generalized description; the work summarized here should not therefore be taken as necessarily applying to all fungi.

The endogenous respiration of ascospores of *Neurospora tetrasperma* is shown in Figure 2. The curve has an upward break coincident with

62) Here an unspecified contact stimulus thigmotropism is supposed to operate. The possibility of chemotropism has been reopened more recently in studies on *Saprolegnia ferax* (87).

One type of germ tube tropism is well documented—the negative tropism of germ tubes toward each other (55 100 117). As described by Stadler (263) germ tubes of *Rhizopus nigricans* form on the side of the spore distant from other spores and in addition actually bend away from other germ tubes during their growth. The stimulus is almost certainly chemical and it is tempting to consider an auxinlike growth factor in explanation of the bending phenomenon. This possibility is also suggested by the finding that germ tubes of rust fungi are longer when germinating spores are crowded than when they are more dispersed (83 314). However Stadler (263) presents some evidence against an explanation based on growth factors; instead he postulates an unstable inhibitor which has the effect of thickening or strengthening the wall of the spore or germ tube. The last word on this problem has clearly not been spoken.

Germ tubes of sporidia and of uredospores of *Puccinia* spp. react negatively to strong light; studies with filters indicate that the active wavelength is in the blue region of the visible spectrum (90 98 235 266). It will be recalled from Chapter 11 that it is these shorter wavelengths which are most active in phototropic and inductive effects on reproductive structures. Negative phototropism is also shown by conidia of powdery mildew fungi (204 305).

3 SPORE METABOLISM

Water, minerals, carbohydrates, and other constituents of fungus spores have been mentioned incidentally in Chapter 2. In relation to spore germination the important materials are water—considered elsewhere in this chapter—and probably lipids. It should also be recalled that spores of *Aspergillus niger* are relatively high in total phosphorus, a part of which is in organic combination (7 154).

Although the fat content of most spores is not unusually high (218 271), three bits of evidence suggest that fats provide energy for the germination of spores which do not require an exogenous source of carbon. First, microscopic observation shows that stainable lipids disappear during germination (82 150). Second, the respiratory quotient of *Neurospora sitophila* macroconidia is about 0.7 (208), a value characteristic of but of course not sufficient evidence for fat oxidation. Finally, the data of Table 1 show that at least in high fat spores the principal organic material disappearing during germination is the

Table 2 Endogenous and Exogenous Oxygen Uptake by Conidia of Fungi*

Species	Oxygen Uptake†		
	No Substrate (endogenous)	Sucrose 1 per cent	Exogenous‡ Endogenous
<i>Neurospora sitophila</i>	8.6	68.6	7.0
<i>Monilinia fructicola</i>	5.1	54.2	9.6
<i>Cephalosporium acremonium</i>	2.7	42.8	14.8
<i>Mycrothecium verrucaria</i>	8.8	22.1	1.5
<i>Stemphylium sarcinariforme</i>	2.0	21.7	9.8
<i>Alternaria oleracea</i>	7.1	17.4	1.5
<i>Aspergillus niger</i>	12.8	17.7	0.4
<i>Ph. opus nigricans</i>	8.5	9.2	0.1

* Drawn from more extensive data of McCallan, Miller, and Weed (173) by permission of the Boyce Thompson Institute for Plant Research, Inc.

† In $\mu\text{l}/5 \text{ hr}/\text{mg}$ fresh spores.

‡ Exogenous = uptake with sucrose less endogenous uptake.

difficult (b4 110 254). In general, but with some exceptions, germination is more sensitive than spore respiration to both inorganic and organic fungicides (Chapter 14).

4. MATURATION AND DORMANCY

It is a common observation that morphologically normal and apparently fully developed spores often do not germinate until after a period of time. If this period is relatively short, we speak of maturation; if it is on the contrary a matter of weeks or months, the spore is said to have a dormant stage or resting period. There is as yet no evidence which enables us to say whether these two phenomena are different or the same.

The failure of newly formed spores to germinate normally is best documented in studies on the uredospores of the rust fungi (73 200 245 322), the behavior of which is exemplified by the data of Figure 3. As shown, immature spores have both a slower rate of germination and lower total germination than have mature spores. In addition, immature spores are more sensitive to unfavorable temperatures (57 73). It appears from most work that immature uredospores can ripen after removal from the sorus (11 57 178). The very long period of poor germinability of aeciospores of *Gymnosporangium* spp. has been noted repeatedly but has not as yet been analyzed (20 99 178 222 282).

Both asexual and sexual spores of ascomycetes germinate poorly




Figure 2 Oxygen consumption of activated (curve 1) and dormant (curve 2) ascospores of *Neurospora tetrasperma*. The arrow marks the time of appearance of germ tubes. Redrawn from Coddard and Smith (110) by permission of Plant Physiology.

the appearance of the germ tube this particular type of curve is not universal in the fungi and would not necessarily occur under different conditions of nutrient or oxygen supply. Endogenous respiration in *Memnoniella echinata* and *Aspergillus luchuensis* is affected by ammonium ion and by the previous cultural history of the spore population (64). Oxygen pressure also affects the endogenous rate (110, 194).

Spores generally, but not always, respond to added substrates by an increase in respiratory rate. Increases may also be induced by provision of nitrogen or of complex nutrients in addition to a carbon source (173, 193, 254, 255). The data of Table 2 are fairly typical in that the range of response is from zero to high, since however the substrate was sucrose the results reflect hydrolytic as well as respiratory capacities.

Just as mature mycelium spores of a given species may or may not ferment glucose anaerobically, Uredospores of *Puccinia graminis tritici* do not form carbon dioxide anaerobically (254). At the other extreme both activated ascospores and conidia of *Neurospora* spp. ferment vigorously and have demonstrable glycolytic enzymes (59, 110, 209).

Not much can be said at this time of detailed respiratory pathways in spores. Acids of the tricarboxylic cycle are oxidized by conidia of *Neurospora sitophila* are found in conidial homogenates and appear in the medium during germination (174, 208, 293). Poisons which affect glycolysis and the cytochrome system usually inhibit spore respiration but several anomalies in the results make interpretation dif-

Table 2 Endogenous and Exogenous Oxygen Uptake by *Conidia* of Fungi*

Species	Oxygen Uptake†		
	No Substrate (endogenous)	Sucrose 1 per cent	Exogenous‡ Endogenous
<i>Neurospora sitophila</i>	8.6	68.6	7.0
<i>Monilia fructicola</i>	5.1	54.2	9.6
<i>Cephalosporium acremonium</i>	2.7	42.8	14.8
<i>Myrothecium verrucaria</i>	8.8	22.1	1.5
<i>Stemphylium sarcinaeforme</i>	2.0	21.7	9.8
<i>Alternaria oleracea</i>	7.1	17.4	1.5
<i>Aspergillus niger</i>	12.8	17.7	0.4
<i>Rhizopus nigricans</i>	8.5	9.2	0.1

* Drawn from more extensive data of McCallan, Miller, and Weed (173) by permission of the Boyce Thompson Institute for Plant Research, Inc.

† In $\mu\text{l}/5 \text{ hr}/\text{mg}$ fresh spores

‡ Exogenous = uptake with sucrose less endogenous uptake

ficult (64, 110, 254). In general, but with some exceptions, germination is more sensitive than spore respiration to both inorganic and organic fungicides (Chapter 14).

4. MATURATION AND DORMANCY

It is a common observation that morphologically normal and apparently fully developed spores often do not germinate until after a period of time. If this period is relatively short, we speak of maturation; if it is on the contrary a matter of weeks or months, the spore is said to have a dormant stage or resting period. There is as yet no evidence which enables us to say whether these two phenomena are different or the same.

The failure of newly formed spores to germinate normally is best documented in studies on the uredospores of the rust fungi (73, 200, 245, 322), the behavior of which is exemplified by the data of Figure 3. As shown, immature spores have both a slower rate of germination and lower total germination than have mature spores. In addition, immature spores are more sensitive to unfavorable temperatures (57, 73). It appears from most work that immature uredospores can ripen after removal from the sorus (11, 57, 178). The very long period of poor germinability of aeciospores of *Gymnosporangium* spp. has been noted repeatedly but has not as yet been analyzed (20, 99, 178, 222, 282).

Both asexual and sexual spores of ascomycetes germinate poorly




Figure 3 The rate of germination of mature (curve 1) and immature (curve 2) urectospores of *Phragmidium mucronatum*. Redrawn from Cochrane (57) by permission of the Cornell University Agricultural Experiment Station

when newly formed (42 145 240). Conidia of powdery mildew fungi do not germinate until detached from the chain in which they are formed possibly because they are impermeable to gases until exposed (33) or because the parent mycelium forms a diffusible germination inhibitor (70) this is not therefore comparable to immaturity.

Many phycomycetes form thick walled resting spores these are probably always the site of meiosis and at least part of their apparent resting period represents the time required for cytological processes (81). Whether there is in addition a further period of purely physiological dormancy is still uncertain. There can be no doubt that the environment especially temperature influences the length of the resting period in some members of the Saprolegniales (216) and Peronosporales (3 17 18). However it is not yet possible to define the locus of this effect. Many oospores it should be noted germinate immediately after their formation (157 321) as do also the resistant sporangia of *Blastocladiella* sp. (45).

The rest period of resistant sporangia of *Allomyces* spp. is shortened dramatically by certain cultural conditions and by indoleacetic acid (176 177). In this and related genera the strength of the sporangium wall may be one of the decisive factors in germination (176 270)—sporangia grown under conditions in which the wall is thin germinate without a rest period.

The ascospores of many ascomycetes particularly but not exclusively coprophilic forms exhibit a pronounced dormancy which is broken by brief exposure to high temperature (69 120 121 122 123 139 251). Rather intensive study of ascospore dormancy in *Neurospora* spp. has revealed several features of the phenomenon

1 Activation—breaking of dormancy—is reversible i.e. activated ascospores become dormant again if prevented from germinating (107)

2 Activated ascospores decarboxylate pyruvic acid and respire vigorously dormant spores respire slowly (Figure 2) and do not decarboxylate pyruvate (108 110)

3 Furfural and related unsaturated heterocyclic compounds activate fresh ascospores (79 272 273). Aged dormant spores are activated by furfural only if they are first heated (275)

Emerson (80) proposes that furfural duplicates the action of a natural catalyst which is normally induced to form or to become active by the heat treatment. It is of course equally possible that a natural inhibitor is present in fresh spores and is inactivated by heat or by those chemicals which induce germination

Neither permeability to water nor to gases can be invoked as a factor in ascospore dormancy in *Neurospora tetrasperma* (107 170). However the activating effect of alkali on ascospores of *Ascobolus* spp. (318) may implicate permeability to water and dormant ascospores of *N. tetrasperma* are relatively impermeable to cations (169a)

Sporangiospores of *Phycomyces blakesleeanus* are activated by a heat treatment e.g. 15 minutes at 50°C (232). The effect is duplicated by hypoxanthine

Teliospores of the rust fungi normally do not germinate until several months after their formation. Although chemical and moisture treatments have been reported to hasten germination (142 181 196 221 280) the same treatments fail in other studies (129 156). Until systematic analysis has been accomplished we must assume a long obligate rest period as characteristic in these spores

In the life cycle of the smut fungi the chlamydospore is analogous to the teliospore of the rust fungi. Periods of 200 days or more are required for germination to begin in certain races of *Ustilago striaeformis* (66 152). The germination of these spores is often much accelerated by chemical treatments (47 66 165 203 206) by low temperature soaking (136) by high temperature cultivation of the infected host (152) or by variations in moisture (281). Mechanical abrasion also shortens the dormant period in *Ustilago striaeformis* (153)

It is not yet possible to envision a single explanation for all types of

long obligatory rest periods. If we define dormancy as a rest period after the spore has completed both its morphological and its cytological development and if we assume that at least part of the rest period in the phycomycete and basidiomycete spores just considered is such true dormancy then the most promising hypothesis is that some factor associated with the heavy spore wall—possibly impermeability possibly simple mechanical strength—is determinative. It is not likely that the same explanation will fit ascospore dormancy in *Neurospora* in which an internal biochemical mechanism seems more probable. However ascospore dormancy in *Onygena equina* develops only after the heavy spore wall is laid down—the immature spore germinates without a dormant period (30).

5 SPORE LONGEVITY

Several fungi form asexual spores whose survival under the most favorable conditions is measured in days only: examples include the conidia of the powdery mildew fungi (317), the sporangium of *Peronospora tabacina* (302) and the sporidia (basidiospores) of some rust fungi (178, 222, 262). The short life of the powdery mildew conidia is probably related to the high water content of the spores (317).

Aeciospores and uredospores of the rust fungi are intermediate in longevity. Although in a few species these spores may under exceptionally favorable conditions survive for as long as a year (57, 77, 89, 214), their lives under most conditions of storage are weeks or months only (6, 20, 72, 73, 74, 221) and are even less under uncontrolled natural conditions (57, 237). Early work on spore survival in the rust fungi is reviewed by Maneval (193) and by Zimmerman (322).

Types of spore characterized by a long or very long life include the chlamydospores of the Ustilaginales (88, 171, 278), basidiospores of Hymenomycetes (95, 127) and at least some ascospores (46, 127). Spores with long periods of inability to germinate—resting spores of phycomycetes, teliospores of the rust fungi and certain ascospores—are *ipso facto* long lived.

Although the rule in the fungi is that asexual spores are relatively short lived, the conidia of certain imperfect fungi and of actinomycetes may survive very long periods (15, 175, 212, 234, 297). Ecologically it appears that the survival function typical of sexual spores is taken over in these forms which lack a sexual stage by the conidia. Dried swarm spores of the Myxomycetes similarly are reported to survive many years (78, 258).

Techniques of preservation of spores are briefly reviewed in Chapter I

Temperature is one of the major factors in the longevity of spores. Almost without exception survival is greatest at low temperature i.e. near 0 C. This is particularly well documented for uredospores of rust fungi (57, 77, 214, 215, 237) and conidia of powdery mildew fungi (317). Presumably spore metabolism is accelerated at high temperature reserves are exhausted and death (inability to germinate) ensues. The effect of subzero temperatures on survival is considered later (Chapter 13).

Relative humidity is the second major factor in spore longevity and in nature is probably more often limiting than is temperature. Here however the relation is not so simple and we must for the present at least distinguish three types of response to atmospheric moisture. Conidia of *Sphaerotheca pannosa* (167) sporidia of rust fungi (222, 262) and conidiosporangia of *Phytophthora* spp. (63, 141) survive longest at saturation and are less viable at any relative humidity less than 100 per cent. Typical data appear in Figure 4. It should be noted that precisely these spores have just been described as short lived in general; their susceptibility to desiccation is almost certainly the reason for their failure to survive.

Uredospores of the rust fungi exhibit a different and characteristic response to humidity. Virtually all studies agree that survival is great




Figure 4. The survival of sporangia of *Phytophthora infestans* at different relative humidities. Redrawn from Croser (63) by permission of the Cornell University Agricultural Experiment Station.

long obligatory rest periods. If we define dormancy as a rest period after the spore has completed both its morphological and its cytological development and if we assume that at least part of the rest period in the phycomycete and basidiomycete spores just considered is such true dormancy then the most promising hypothesis is that some factor associated with the heavy spore wall—possibly impermeability possibly simple mechanical strength—is determinative. It is not likely that the same explanation will fit ascospore dormancy in *Neurospora* in which an internal biochemical mechanism seems more probable. However ascospore dormancy in *Onygena equina* develops only after the heavy spore wall is laid down—the immature spore germinates without a dormant period (30).

5 SPORE LONGEVITY

Several fungi form asexual spores whose survival under the most favorable conditions is measured in days only. Examples include the conidia of the powdery mildew fungi (317) the sporangium of *Peronospora tabacina* (302) and the sporidia (basidiospores) of some rust fungi (178 222 262). The short life of the powdery mildew conidia is probably related to the high water content of the spores (317).

Aeciospores and uredospores of the rust fungi are intermediate in longevity. Although in a few species these spores may under exceptionally favorable conditions survive for as long as a year (57 77 89 214) their lives under most conditions of storage are weeks or months only (6 20 72 73 74 221) and are even less under uncontrolled natural conditions (57 237). Early work on spore survival in the rust fungi is reviewed by Maneval (193) and by Zimmerman (322).

Types of spore characterized by a long or very long life include the chlamydospores of the Ustilaginales (88 171 278) basidiospores of Hymenomycetes (95 127) and at least some ascospores (46 127). Spores with long periods of inability to germinate—resting spores of phycomycetes teliospores of the rust fungi and certain ascospores—are *ipso facto* long lived.

Although the rule in the fungi is that asexual spores are relatively short lived the conidia of certain imperfect fungi and of actinomycetes may survive very long periods (15 175 212 234 297). Ecologically it appears that the survival function typical of sexual spores is taken over in these forms which lack a sexual stage by the conidia. Dried swarm spores of the Myxomycetes similarly are reported to survive many years (78 258).

Techniques of preservation of spores are briefly reviewed in Chapter 1

Temperature is one of the major factors in the longevity of spores. Almost without exception survival is greatest at low temperature i.e. near 0 C. This is particularly well documented for uredospores of rust fungi (57, 77, 214, 215, 237) and conidia of powdery mildew fungi (317). Presumably spore metabolism is accelerated at high temperature reserves are exhausted and death (inability to germinate) ensues. The effect of subzero temperatures on survival is considered later (Chapter 13).

Relative humidity is the second major factor in spore longevity and in nature is probably more often limiting than is temperature. Here however the relation is not so simple and we must for the present at least distinguish three types of response to atmospheric moisture. Conidia of *Sphaerotheca pannosa* (167), sporidia of rust fungi (222, 262) and conidiosporangia of *Phytophthora* spp. (63, 141) survive longest at saturation and are less viable at any relative humidity less than 100 per cent. Typical data appear in Figure 4. It should be noted that precisely these spores have just been described as short lived in general; their susceptibility to desiccation is almost certainly the reason for their failure to survive.

Uredospores of the rust fungi exhibit a different and characteristic response to humidity. Virtually all studies agree that survival is great




Figure 4 The survival of sporangia of *Phytophthora infestans* at different relative humidities. Redrawn from Croser (63) by permission of the Cornell University Agricultural Experiment Station.

est at intermediate relative humidity and is less at either extreme (6 57 129 214 215 221 237) Chlamydospores of *Urocystis tritici* respond in the same way (206) Damage at low moisture levels is probably simple water loss consideration of the possible effects of high humidity leads us to the third and final category of humidity response

This last type of response is exemplified by both ascospores and endoconidia of *Endoconidiophora fagacearum* (201) and by conidia of *Helminthosporium oryzae* (211) these remain viable longest at low humidity especially if the temperature is unfavorably high for survival It has been suggested that at high humidity the metabolism of this type of spore is accelerated with a consequent reduction in longevity (201) the mechanism of such an effect is obscure but this is at present the most attractive hypothesis and may be extended to the problem of uredospore survival at high humidity

Spores in nature are often protected from rapid loss of water by a gelatinous matrix by the structure in which they are formed or by host plant tissues several such phenomena are reviewed by Gottlieb (113)

Spores of *Myrothecium verrucaria* are strongly affected by the history of the culture in particular by the composition of the medium (64) Spores from a mycelium grown on filter paper survive more than twice as long as those from any of three agar media this greater viability is not however correlated with the rate of loss of dry weight The growth medium also exerts a significant effect on the dry weight nitrogen content and respiratory behavior of spores of *M. verrucaria* Conidia of nutritionally deficient mutants of *Ophiostoma multiannulatum* die less rapidly in a starvation medium than do conidia of the parent non deficient stock it has been suggested that the deficient spores metabolize less and hence survive longer (96)

Physiologically senescent spores are characterized by poor germinability by a slower rate of germination (36 57 194) and by lower respiratory activity (64)

In *Streptomyces* sp ageing of the spore population at low temperature increases the recovery of mutant types the mechanism is obscure (294)

6 NUTRITION AND GERMINATION

Germination may be considered as a growth process in which some or all the necessary nutrients are present in the spore *ab initio* We may therefore expect two extreme types of relation to external nutrients—complete independence in some fungi and complete dependence

on one or more nutrients in others—and various degrees of partial dependence on the environment.

Experimentally nutritional independence of the spore is difficult to prove it is essential that washed spores free of materials carried over from the growth medium be used or that spores be collected in such a way as to prevent contamination with nutrient materials (163). Washing of course must be as brief as possible in order to minimize any loss of materials from the interior of the spore. Washed ascospores of *Neurospora tetrasperma* germinate in distilled water (274) and are therefore to be classed as nutritionally independent. The uniformly good germination of rust uredospores and aeciospores which have been collected by shaking from the host plant suggests that these too are independent of the environment it has been noted earlier (Table 1) that uredospores are unusually high in reserve lipids. Although it is perhaps justifiable to assume that the larger resting spores of fungi are nutritionally independent critical evidence is lacking. Conidia of the powdery mildew fungi collected like the rust spores by shaking germinate well on water although sugars and several natural materials improve germination (167 307 309 317).

Nutritional dependence is more easily demonstrated and is probably more common than independence. Many fungus spores germinate very poorly or not at all without added nutrients examples include basidio spores of *Psalliotia bispora* and other Hymenomycetes (91 95 105) ascospores of *Sordaria fimicola* (29 43) and conidia of several imperfect fungi (75 194). Other fungi too numerous to mention individually germinate fairly well in water but do so more completely or more rapidly in nutrient media (5 53 75 85 121 202) or in contact with complex biological products (35 36 38 60 101 113 147 151 158). Since unwashed spores have been used in many of these studies nutritional deficiencies are probably more acute than they appear to be.

The simplest view is that the three reactions just described—complete independence partial dependence and complete dependence—reflect the adequacy or availability of the reserve substances of the spore. Critical studies on washed spores show that requirements may be simple or complex. Washed spores of *Sclerotinia fructicola* require a carbon source (163) those of *Fusarium roseum* need both carbon and nitrogen (255) and spores of *Glomerella cingulata* must be supplied carbon nitrogen phosphorus and sulfur (161). Responses to minerals e.g. magnesium (163) are not necessarily nutritional. Spores are so sensitive to ions that minerals such as magnesium may act primarily as antidoting agents.

Mutants of *Neurospora crassa* which are deficient for amino acids




Figure 5 The response to graded doses of lysine of macroconidia of a lysine requiring mutant of *Neurospora crassa*. Redrawn from Ryan (240) by permission of the *American Journal of Botany*

produce macroconidia which require the amino acid for prompt germination although the spores will germinate in time without the compound (239-240). The response of one such mutant to lysine is shown in Figure 5. The assay is complicated by the fact that amino acids are somewhat inhibitory to spore germination in both mutant and normal *Neurospora crassa*. Parathiotrophic mutants of *Aspergillus nidulans* respond in the same way to sulfur compounds (252).

Hypoxanthine partially replaced by guanine accelerates both spore germination and early growth of *Phycomyces blakesleeana* (226-227, 228-229, 230-231, 233-238). The effect of organic acids and of various physical and chemical treatments in increasing spore germination in this species has tentatively been ascribed to their influence on the availability of hypoxanthine (232).

Although as mentioned above complex biological materials improve germination in many species relatively few vitamin requirements for spore germination have been worked out. *Memnoniella eclanata* requires biotin (216). *Myrothecium verrucaria* also needs biotin but only in the phase following germ tube emergence (189) and *Colletotrichum gloeosporioides* is reported without data to respond to riboflavin and other vitamins (61). As discussed briefly in Chapter 10 successful use of the differential germination method for the detection of vitamin deficient mutants implies that these mutants require the factor for spore germination as well as for growth. Promotion of basidiospore germination by co-culture with *Torulopsis sanguinea* is at least suggestive of a vitamin effect (94-95) as is also the observation that the parent mycelium itself stimulates spore germination in *Psalliota campestris* (76).

Germination of uredospores of *Melampsora lini* is increased by methyl *p*-hydroxybenzoate and by aureomycin (288) there is however

no evidence of a nutritional mechanism in this as yet unexplained phenomenon

Conidia of *Erysiphe graminis* respond slightly to nutrients but the response is strongly conditioned by temperature and by the time of collection of the spores i.e. by their previous history (316-317)

Ammonium salts often a utilizable nutrient are toxic to some species of *Mycena* but not to all mycelial growth is much less sensitive or possibly insensitive to concentrations of ammonium tartrate which completely inhibit basidiospore germination (97) Chlamydospore germination in *Tilletia* spp. has repeatedly been found to be inhibited by complex biological products which are nutritive for other organisms (171)

Ecologically a nutritional requirement for spore germination may be an advantage helping to insure that germination will not occur in environments devoid of the materials needed for continued growth

7 pH AND GERMINATION

It has been mentioned earlier in Chapter 1 that spore germination usually has a narrower pH range than growth presumably because there is less time for metabolic products to modify an unfavorable medium. In the same chapter some of the ambiguities of acidity data were considered the same ambiguities attach to germination experiments in particular to the finding that several fungi show a double pH optimum under some conditions (90-283-295)

Analysis of spore germination data reveals several complicating factors (1) strains of a given species differ in response (43) (2) the type of buffer—and a buffer is essential in pH studies—affects the shape and limits of the curve (274) (3) nutrient materials often modify the pH response profoundly (43-295) and (4) the previous history of the spore population influences its behavior (2). These considerations suggest a certain caution in the interpretation of the rather extensive literature on pH response curves to which we now turn

Spores of most fungi germinate best at pH 4.5-6.5 with limits at about pH 3 and pH 8. These rough figures apply to for example uredospores of several rust fungi (2-90-266) ascospores (14-43-301) chlamydospores of *Urocystis tritici* (206) and spores of myxomycetes (257)

Variations from this general pattern are several a few fungi have more acid optima (283-295) some have optimum zones extending to about pH 8 (60-146-160-161-295) and many germinate over a somewhat wider range than the typical (53-198-257-295-301)

It is probable that only in rather unusual environments eg very acid soils is pH a natural limiting factor for spore germination. Insofar as acidity and alkalinity are ecological influences their effects on mycelial growth and spore formation are likely to be more decisive.

8 WATER AND GERMINATION

Available water is one of the primary determinants of spore germination. Consequently a considerable body of evidence has accumulated on the moisture responses of economically important fungi; several reviews of these data have been published (56, 71, 113, 256). The effect of humidity on spore survival has already been considered.

Certain problems of method have received less attention than they deserve. Most investigators have reported data in terms of relative humidity, although this practice is justifiable; a clearer picture of water relations emerges when the vapor pressure deficit is taken as the criterion (67, 262). The absolute humidity is not a useful value (224). The importance of accurate temperature control, especially in attempts to discriminate between free water and a saturated atmosphere with no free water, has not always been realized. Both static and moving air systems for the simultaneous control of temperature and humidity have been described (56, 57, 67); the static principle—incubation in a closed system over a humidity regulating solution—is both simpler and more accurate.

When germination is measured on solid substrates which take up water—agar, grain, textiles, etc.—the minimum humidity permitting germination is often lower than that in the absence of substrate (4, 22, 118). Thus we may reasonably infer means that the water absorbing capacity of these substrates is greater than that of the spore, but this explanation begs the question of how the spore is able to draw water from the substrate.

The response to moisture is usually conditioned by temperature. Thus conidia of *Uncinula necator* are adversely affected by low humidity at 28°C or above, not at 25° or 20° (67). The optimum humidity for germination of *Aspergillus niger* conidia is substantially higher at 40° than at 30°C (25). In *Alternaria citri* both germination time and germ tube length are less affected by relative humidity at the temperature optimum for germination than at non-optimal temperatures (286).

Excluding the powdery mildew and the rust fungi, to be taken up later, the fungi can be classified according to the minimum relative humidity at which detectable germination occurs. Several, including species of *Aspergillus*, *Penicillium*, and a few other genera, form spores

which germinate at relative humidities near 80 per cent (4 25 118 132 162 244 250 320). A somewhat larger group of species germinate at humidities of 90-95 per cent and do not require liquid water examples include *Alternaria brassicae* *Colletotrichum* spp (51) *Ustilago* spp (56) *Cladosporium* spp (51 119 224) *Botrytis cinerea* (224) *Fomes annosus* (225) *Septoria apu* *Fusarium* spp and *Verticillium albo-atrum* (247). Still higher humidity is required by such fungi as *Venturia inaequalis* (56) and *Magnusia brachytricha* (277). Most of or all the fungi mentioned in this paragraph germinate better at high humidity than at low regardless of the minimum value.

Finally the spores of many fungi are found to germinate only in liquid water in this group we may mention conidia and ascospores of *Endoconidiophora sagacearum* (60) asexual spores of the *Peronosporales* (73 141 302 308) and conidia of *Sclerotinia* spp (56 73).

Almost all studies of uredospore germination in rust fungi agree that liquid water is required and that germination is nil even at 100 per cent humidity in the absence of free water (1 10 57 111 129 133 134 267 308 322). It is not always realized that temperature control must be extremely accurate if a humidity of 100 per cent is to be maintained without condensation of liquid water. For this reason we must reject those reports in which accurate temperature control is not specified and germination in the absence of free water is claimed. This stricture does not however apply to the work of Clayton (56) who found that uredospores of *Puccinia coronata* and *P. graminis* germinate at 100 per cent relative humidity albeit less well than in the presence of free water.

Aeciospores teliospores and sporidia of *Gymnosporangium globosum* all require liquid water for germination (174).

Alone among the fungi the powdery mildew fungi form conidia which in the experience of most workers germinate at very low relative humidity even over desiccants (33 34 48 56 67 306). Exposure to liquid water in fact inhibits germination and may cause death of the spore by plasmolysis (67 167). As a result of these water relations the powdery mildew diseases are favored in nature by relatively dry conditions (26 48 311).

Yarwood has in a series of papers (306 310 312) provided evidence that the ability of these spores to germinate at low humidity is a function of their high water content. Two different methods of determining water content agree in assigning a value of about 70 per cent water to the conidia of *Erysiphe polygoni*. This is much higher than the water content of other fungus spores. Correspondingly although difficulties of measurement exist the density of the spores of powdery mildew conidia is less than that of most other spores. Finally it has

been shown that spore volume decreases during germination in dry air. Although somewhat conflicting data and a theory based on the hypothetical conversion of bound to free water have been reported (33-34) it seems that the high water content of conidia of the powdery mildew fungi explains their ability to germinate at low humidity. The occurrence of plasmolysis in pure water argues for a rapid entrance into the spore; this is still to be reconciled with the observation that the loss of water from ungerminated spores is very slow—approximately 0.08 per cent of the loss from an equivalent free water surface (312).

These studies on the powdery mildews have been generalized by Yarwood (310) in the proposition that fungus spores of the usual type, i.e. those with a low water content must absorb water to about the level of that in the powdery mildew conidia (70 per cent) before they can germinate. This hypothesis explains the swelling of these spores so frequently reported (p. 391) and in fact demands that all spores of low water content swell on germination. This appears to be the most promising general explanation of humidity relations.

As is to be expected, fungi tolerant of low humidity are also in general tolerant of high osmotic pressure in the germination medium. This relation is brought out in Figure 6.

Conidia of *Erysiphe graminis* germinate in concentrations of deuterium oxide up to 100 per cent, but the heavy water is markedly inhibitory to germ tube elongation (220). *Macrosporium sarcinaeforme* is unable to germinate in 60 per cent deuterium oxide (213).




Figure 6 The relation of the minimum relative humidity to the maximum tolerated concentration of sodium chloride for spore germination in several fungi (*Aspergillus* spp., *Penicillium* spp., *Fusarium moniliforme*). Each plotted point refers to one species. Drawn from data of Armolik and Dickson (4).

9 TEMPERATURE AND GERMINATION

The so-called cardinal temperatures for spore germination—the minimum optimum and maximum—have been determined for very many fungi most data on plant pathogens reported prior to 1936 are summarized by Togashi (285) other results by Wolf and Wolf (303) The student is immediately struck by the frequent and pronounced disagreements in this literature some of these are probably inherent in the nature of the material whereas others are traceable to inadequate methods of study

Some examples of reported cardinal temperatures are shown in Table 3 The total range of temperatures permitting at least some germina-

Table 3 Cardinal Temperatures of Spore Germination

Species	Minimum C	Optimum C	Maximum C
<i>Albugo candida</i> (199)	0	~10	25
<i>Peronospora effusa</i> (102)	~4	9	~27
<i>Mastigosporeum album</i> (24)	—	10	20-25
<i>Gonartium ribicola</i> uredospores (71)	8	14	25
<i>Puccinia malvacearum</i> teliospores (71)	3	14	30
<i>Phoma miedum mucronatum</i> aeciospores (57)	3-6	15-21	27-30
<i>Phytophthora infestans</i> direct (63)	6	24	27-30
<i>P. infestans</i> indirect (63)	15	12	21-24
<i>Phoma apicola</i> (13)	10	16-18	23
<i>Tranzschelia punctata</i> uredospores (111)	8	13-26	38
<i>Herpobasidium deformans</i> (115)	1	18-26	33
<i>Endoconidiophora fagacearum</i> (85)	3	21-32	33-36
<i>Colletotrichum indicum</i> (166)	16	20-32	—
<i>Erysiphe polygoni</i> (317)	5	25	35
<i>Allomyces arbuscula</i> meiosporangia (261)	14-16	20-26	32-34
<i>Glomerella cingulata</i> (299)	15	27	35+
<i>Magnusia nitida</i> (276)	15	~32	43
<i>Ustilago zeae</i> (144)	8	26-34	36-38
<i>Rhizopus nigricans</i> (298)	15	26-28	33
<i>R. chinensis</i> (298)	10	43-45	51

tion varies in these data from 13 centigrade degrees in *Phoma apicola* to 40 in *Rhizopus chinensis* and *Magnusia nitida* Optima range from a low of 9 C in *Peronospora effusa* to a high of 43-45° in *Rhizopus chinensis* In general it appears from these and other data and is to be expected on theoretical grounds that fungi with a high optimum have a broader total range than those with a low optimum that is fungi

with a high optimum have in a sense solved the problem of developing at high temperature without losing their capacity to germinate at low temperature. Spore germination in the truly thermophilic fungi and actinomycetes has not been studied quantitatively.

Primarily from the data collected by Togashi (285) we can estimate that the average optimum for plant pathogenic fungi is about 23° . No similar estimate can be made for saprophytic fungi probably the figure is somewhat higher.

Certain groups of fungi can be roughly characterized as a whole by their temperature preferences. Thus species of *Aspergillus* tend to have high optima (21-25) those of *Peronospora* to prefer rather low temperatures (84-285). Uredospores of most rust fungi germinate best at 22° or lower (44-57-72-183-266-268-285-322) although exceptions to this rule have been reported (111). Limited data on *Coprinus* spp suggest a rather high optimum $30-35^{\circ}$ (146).

Conidia of the powdery mildew fungi as a rule germinate best at a moderately low temperature about 21° . However, reported optima for different species range from 6° to 28° and within one species *Erysiphe cichoracearum* strains with optima of 15° and 28° occur (317). Similarly physiologic races of other parasitic fungi e.g. *Puccinia glumarum* (268) and *Tilletia* spp. (171) may differ considerably in their responses to temperature. Intrageneric variations in response are exemplified in *Rhizopus* individual species have optima as low as 27° and as high as 44° (298).

Figure 7 depicts representative types of curves of total spore germination versus temperature. The most frequent shape of curve is roughly symmetrical—or is slightly skewed to the right—with a wide optimum zone. This curve is usually found in studies of the rust fungi (44-57-266-268-269) and has been reported for such divergent types as *Fusarium lini* (31) *Ustilago* spp. (16) *Coccomyces hiemalis* (148) and several Hymenomycetes (260). An extreme example is found in the response of conidia of *Magnusia* spp. (276).

No one type of curve can however be said to characterize all fungi. The two most common modifications are (1) curves which are skewed to the right like most mycelial growth curves (53-115-167-317-318) and (2) curves with a sharp optimum rather than a broad optimal zone (71-93-129-143-296). More rarely a curve skewed to the left has been reported (48-84) from studies of fungi with low optimum temperatures.

The diversity of types of curve becomes less surprising when one considers the number of factors that have been shown to affect the shape of the curve. Of these time of observation is the most impor-




Figure 7 Idealized types of temperature response curves of spore germination. The distinctive features are exaggerated for emphasis. Curve 1 symmetrical with a broad optimum; curve 2 skewed to the right; curve 3 symmetrical with a sharp optimum.

tant early observation yields a curve with a sharply defined optimum but in time the germination at less favorable temperatures catches up and the optimum becomes much broader. This phenomenon appears to be general (13, 23, 84, 166, 219, 241, 260, 299).

Second in importance to the time effect as a factor in determining the shape of the temperature curve is the general principle that the limits of germination are narrower if some other factor is non-optimal. Thus conidia of *Sclerotinia fructicola* at an unfavorably acid pH germinate only at 13–29°C and the curve is sharply peaked at 21°C; at a more favorable reaction pH 2.4 both the total range and the optimum are much broader (283). The depressant effect of low temperature on *Erysiphe graminis* germination is virtually abolished in the presence of nutrients (317) and the minimum temperature of germination in *Colletotrichum lagenarium* is lower in a nutrient medium than in water (101). The principle applies also to relative humidity; i.e. the temperature curve becomes narrower and sharper as the humidity becomes more unfavorable to germination (118, 286).

Although different spore forms of the same species usually have at least similar temperature responses (148, 178, 285) the endoconidia and macroconidia of *Chalaropsis thielavioides* show optima at 9–30

and 12° respectively in this organism the age of endoconidia has a pronounced effect on the shape of the temperature germination curve (168)

So far we have considered the response of germination i.e. initiation of the germ tube to temperature. The further growth of the germ tube may be plotted against temperature. When such curves are compared to germination curves there is often perhaps usually a close correspondence (42 86 93 103 146 241). With many fungi the optimum is rather narrower than that for germination: examples include *Erysiphe graminis* (317) *Physalospora obtusa* (92) *Clasterosporium carpophilum* (103) and several rust fungi (57 20 266). In still other fungi the shape of the curve or the position of the optimum or both are different (84 102 116 148 260). In general it appears that germ tube growth is closer in its temperature response to mycelial growth than it is to spore germination: the data of Snell (260) on *Lenites septaria* are especially instructive in this connection. For this reason there are many problems especially in plant pathology in which the temperature characteristics of germ tube growth are more illuminating than those of germination.

The preceding discussion emphasizes the limitations of the conventional temperature study of spore germination. Results often depend upon the criterion used and upon such sometimes uncontrolled variables as age of spore, humidity and pH. The studies are not however to be dismissed as useless: thus it is often possible to correlate the ecological niche of a parasitic fungus with the response of its spores to temperature (44 57 317).

In this discussion we have tacitly assumed that temperature affects all processes up to the appearance of the germ tube in the same way. There is indirect evidence from two types of temperature reaction that this assumption is not valid: that the processes which initiate spore germination have a different sensitivity from later stages.

Although it might be expected *a priori* that at lower temperatures the rate of all processes is slower, the data of Figure 8 show a more complex situation: to be sure the rate of germination at suboptimal temperatures is somewhat less than the rate at the optimum (18) but the principal effect of low temperature is to delay the onset of germination i.e. to increase the latent period. Once germination has begun it is almost as rapid at low temperature as at high. Similar data may be found in studies of *Fusarium moniliforme* (241) *Sclerotinia fructicola* (299) and *Magnusia* spp. (276). Both rate and latent period are strongly affected by low temperature in *Phytophthora infestans* (63) and *Uncinula necator* (67). The latent period and by inference the metabolic




Figure 8 Temperature and the rate of germination of uredospores of *Phragmidium mucronatum*. Redrawn from Cochrane (57) by permission of the Cornell University Agricultural Experiment Station

reactions occurring in the very earliest phases of germination is uniformly extremely sensitive to low temperature. At supraoptimal temperatures there is again evident a differential sensitivity of these early reactions (284).

Sensitivity to low temperature of the processes concerned in the initiation of germination is paralleled by sensitivity of the same processes to a favorable temperature. The germination of uredospores of *Phragmidium mucronatum* at non-optimal temperatures is markedly accelerated by exposure for a short time to a moderate temperature: for example germination at 9°C is increased about 160 per cent by a pre-exposure for one hour at 18°C during which exposure there is no visible sign of germination (58). The same phenomenon in *Puccinia chrysanthemi* is suggested by the data of Campbell and Dimock (44). Similarly direct germination by germ tube formation of the conidiospores of *Phytophthora infestans* at 20°C is greatly increased by prior exposure for 5–10 minutes to 40°C (279).

It can be suggested therefore that early metabolic reactions in germination have a different and more sensitive response to temperature than do the later stages. Obviously the evidence is far from complete. It is perhaps analogous that the cracking of the sporangium wall in *Blastocladiella* sp. responds somewhat differently to temperature than do the later stages of germination (45).

Parenthetically it may be noted that these results have a methodo-

logical significance in studies of temperature the medium and its container must be adjusted to the experimental temperature before spores are sown. We may assume that this precaution has often been overlooked and deduce that many temperature germination curves are in error in that they exaggerate the extent of germination at sub- and supraoptimal temperatures.

As noted in Chapter 1 a given fungus usually has about the same temperature response whether mycelial growth or spore germination is the criterion (24-86); a few exceptions have been reported (84-298).

10 OXYGEN AND GERMINATION

It will be recalled that mycelial growth in fungi is often possible at very low oxygen pressures but that truly anaerobic growth is unlikely (Chapter 1). Virtually the same applies to spore germination in general with a few exceptions to be noted here.

Failure of submerged spores to germinate or to germinate normally has often been observed and credited without proof to oxygen deficit. As pointed out by Gottlieb (113) too many other factors are affected by submergence for this simple evidence to be acceptable.

Ideally experimental studies on the oxygen requirements for spore germination should include first demonstration of the response to graded oxygen pressures which are accurately measured and second proof that the spores which fail to germinate at a given pO_2 are still viable. Relatively few experiments meet this ideal standard. Uredospores of *Puccinia graminis tritici* (2) and chlamydospores of *Ustilago zeae* (217) fail to germinate at zero oxygen pressure and reach maximum or near maximum germination at 30-38 mm of oxygen. Conidia of *Botrytis cinerea* germinate as well in 5 per cent oxygen as in air (37) but spores of *Synchytrium endobioticum* are severely inhibited at 3 per cent oxygen (292).

Removal of oxygen by pyrogallol prevents germination in for example uredospores of rust fungi (6-129) and chlamydospores of *Ustilago avenae* (143). Although conidia of *Erysiphe graminis* are reported to germinate at a low level in an atmosphere of nitrogen (70) other studies on powdery mildew fungi indicate that germination is prevented by anaerobiosis (34-167). Ascospores of *Neurospora tetrasperma* also fail to germinate anaerobically (107).

Conidiosporangia of three species of *Phytophthora* can germinate by swarmspore formation—but not by germ tube—under anaerobiosis although other members of the Peronosporales fail completely to germinate without oxygen (291). In the Mucorales it appears that spores

of strongly fermentative species have a very low oxygen requirement and those of less strongly fermentative species a higher one all however probably require some oxygen. In this study of the Mucorales (304) it was also found that the swelling of the spore of *Mucor* spp. can occur without oxygen i.e. oxygen is required only at a later stage in development.

Pure oxygen is at least slowly toxic to spores of *Streptomyces scabies* (242) in short term experiments inhibition of germination in *Botrytis cinerea* by high oxygen pressure is measurable but small (37).

It appears to conclude that spore germination usually requires at least some oxygen but that the atmospheric level is more than adequate. In nature it seems unlikely that spore germination is prevented by oxygen deficits except under such extreme conditions as flooded soil or heavily polluted water.

11 CARBON DIOXIDE AND GERMINATION

From the observations cited in Chapter 7 that carbon dioxide often accelerates the early growth of fungi we might guess that spore germination is accelerated by carbon dioxide. This proves to be true in *Aspergillus niger* (223). Effects of high levels of carbon dioxide in increasing chlamydospore germination have been interpreted however as arising primarily from the effect of the dissolved gas on the acidity of the medium (218).

At high concentrations carbon dioxide is partially or even completely inhibitory to spore germination. Brown (37) determined that the spores of seven different fungi if sown in water are completely inhibited by an atmosphere of 20–30 per cent carbon dioxide sensitivity is markedly less if the spores are sown in a plant decoction. Other reports agree that carbon dioxide is inhibitory and that the level at which germination is affected may be low (180–266) or high (139–169). Inhibition at concentrations of 1 per cent or so is probably specific effects of the higher concentrations could be indirect e.g. on acidity or on oxygen supply.

12 BIOLOGICAL EFFECTS ON SPORE GERMINATION

It is commonly observed that excessive crowding of spores on a surface results in reduced germination the phenomenon is known in many fungi (27–37–70–73–148–160–172) and may be described by the general term self inhibition. In *Neurospora crassa* crowding of conidia reduces the germination rate but not total germination (240). These

interactions of spore populations are formally analogous to the self inhibition of growth reported in species of actinomycetes (207-290) and inhibition of spore germination in *Phycomyces blakesleeanus* by products of mycelial growth (249). However we may anticipate that these reactions of growing mycelium are more complex and more dependent on the environment than is self inhibition in spores. *Bacillus globigii* endospores produce a substance antagonized by alanine which inhibits germination of fresh endospores of the same species (265).

Self inhibition has been most clearly demonstrated in studies of rusts *Uromyces phaseoli* (313-314) *Puccinia graminis tritici* (2) and *P. helianthi* (315). The inhibitory effect of a dense concentration has been demonstrated *in vivo* on the host plant as well as *in vitro* (Figure 9). Most of our information indicates that the active material is volatile.

Allen (2) has determined several characteristics of the substance—possibly substances—responsible for self inhibition of *Puccinia graminis tritici* uredospores. The active material is stable to heat, is inactivated or removed from solution by glass surfaces, and is more effective at high pH than at low. The inhibitor is a spore metabolite produced under aerobic conditions. This differs somewhat from the situation




Figure 9 The effect of uredospore concentration on germination in *Puccinia graminis tritici*. Curve 1 spores in open vessels; curve 2 spores in sealed vessels. Redrawn from Allen (2) by permission of *Phytopathology*.

in *Aspergillus niger* (155) and *Coccomyces hiemalis* (148) in both of which limited evidence suggests that an inhibitor is present in the spore at the time of its formation and can be removed by washing

Preliminary evidence has been reported by Forsyth (91) that *Puccinia graminis tritici* uredospores produce 2 methyl butene 2 (trimethylethylene) and that this compound inhibits spore germination. Whether this is the effective substance remains to be seen

Self inhibition has at least one obvious ecological function: that germination of spores within the fructification is minimized. The ungerminated spore is much more resistant to the environment than the germinated and is of course the effective agent of dispersal. Although germination *in situ* has occasionally been reported, it appears to be the exception: this suggests that self inhibition is a general principle in spore physiology and that it is not limited to the few fungi so far studied

At the other extreme from self inhibition is the situation in *Ophiobolus graminis* (210): ascospores germinate slowly if isolated singly, more rapidly if many spores are present. Still a different effect occurs in certain myxomycetes (257) in which spore germination is promoted by the medium upon which a culture has grown

Plant materials, besides increasing spore germination through nutrient effects or carbon dioxide evolution, in some cases inhibit germination, e.g. of *Botrytis cinerea* (36, 38) and *Venturia inaequalis* (106). The mechanism of these effects, as that of the inhibition of spore germination in soil (50), is still unknown. Microbial products may also inhibit germination: these include, besides the fungitoxic antibiotics, the compound fusaric acid (Chapter 8) produced by several *Fusarium* spp. and active against *Ustilago zeae* (104).

BIBLIOGRAPHY

- 1 Abe T. 1933. *Ann. Phytopathol. Soc. Japan* 2: 50-512.
- 2 Allen P. J. 1955. *Phytopathology* 45: 9-266.
- 3 Arens K. 1909. *Jahrb. wiss. Botan.* 10: 57-9.
- 4 Armolik N. and J. G. Dickson. 1956. *Phytopathology* 46: 462-463.
- 5 Atwood K. C. and F. Mukai. 1955. *Genetics* 40: 438-443.
- 6 Bailey L. D. 1923. *Minn. Univ. Agr. Exp. Sta. Tech. Bull.* 16: 1-31.
- 7 Bajaj V. S. P., D. M. Damle and P. S. Krishnan. 1954. *Arch. Biochem. Biophys.* 50: 451-460.
- 8 Baker G. E. 1945. *Mycologia* 37: 582-600.
- 9 Balls W. L. 1905. *New Phytologist* 4: 18-19.
- 10 Beauverne J. 1904. *Compt. rend. (Paris)* 179: 993-996.
- 11 Becker J. 1929. *Kuhn. Arch.* 19: 33-411.
- 12 Bellinger H. 1956. *Zentr. Bakteriol. Parasitenk. Abt. II* 109: 13-16.

- 13 Bennett C W 1921 *Mich State Univ Agr Exp Sta Tech Bull* 53 1-40
- 14 Bennett F T 1937 *Ann Appl Biol* 21 236-257
- 15 Berestneff N M 1907 *Gentr Bakteriell Parasitenk Abt I Pfl* 40 99
- 16 Bever W M 1915 *J Agr Research* 71 41-59
- 17 Blackwell E 1940 *Trans Brit Mycol Soc* 24 68-86
- 18 Blackwell E 1913 *Trans Brit Mycol Soc* 26 93-103
- 19 Blackwell E and G M Waterhouse 1931 *Trans Brit Mycol Soc* 15 294-310
- 20 Bliss D E 1933 *Iowa State Coll Agr Exp Sta Research Bull* 166 339-399
- 21 Blochwitz A 1933 *Ann Mycol* 31 73-83
- 22 Block S S 1935 *Appl Microbiol* 1 28-293
- 23 Blodgett E C 1936 *Phytopathology* 26 115-152
- 24 Bollard E G 1930 *Trans Brit Mycol Soc* 33 2 0-264
- 25 Bonner J T 1918 *Mycologia* 40 728-738
- 26 Boughey A S 1949 *Trans Brit Mycol Soc* 32 179-187
- 27 Boyd A E W 1930 *Ann Appl Biol* 39 302-329
- 28 Bradley S G and J Lederberg 1936 *J Bacteriol* 72 219-220
- 29 Bretzlaff C W Jr 1934 *Am J Botany* 41 58-67
- 30 Brierley W B 1917 *Ann Botany* 31 127-132
- 31 Broadfoot W C 1926 *Phytopathology* 16 921-918
- 32 Brodie H J 1936 *Am J Botany* 23 309-327
- 33 Brodie H J 1945 *Can J Research C* 23 198-211
- 34 Brodie H J and C C Neufeld 1912 *Can J Research C* 90 41-61
- 35 Brown R 1946 *Nature (London)* 157 64-69
- 36 Brown W 1922a *Ann Botany* 36 101-119
- 37 Brown W 1922b *Ann Botany* 36 227-283
- 38 Brown W 1922c *Ann Botany* 36 283-300
- 39 Brown W 1936 *Botan Rev* 2 236-283
- 40 Brown W and C C Harvey 1927 *Ann Botany* 41 643-662
- 41 Buller A H R 1931 *Researches on Fungi Vol 6* London Longmans Green and Co pp 513
- 42 Burgert I A 1934 *Phytopathology* 24 384-396
- 43 Butler E E 1936 *Mycologia* 48 312-348
- 44 Campbell C E and A W Dimock 1935 *Phytopathology* 43 614-618
- 45 Canino L C 1931 *Antonie van Leeuwenhoek J Microbiol Serol* 17 59-96
- 46 Chamberlain D W and J L Allison 1913 *Phytopathology* 35 241-248
- 47 Cheo P C and J G Leach 1950 *Phytopathology* 40 581-589
- 48 Cherewick W J 1944 *Can J Research C* 22 52-87
- 49 Chilton S J P 1943 *Mycologia* 33 13-20
- 50 Chinn S H F 1953 *Can J Botany* 31 718-724
- 51 Chowdhury S 1937 *Indian J Agr Sci* 7 63-67
- 52 Chowdhury S 1944 *J Indian Botan Soc* 23 91-106
- 53 Chowdhury S 1946 *J Indian Botan Soc* 25 123-130
- 54 Christensen J J and F R Davies 1937 *Mycologia* 29 8-99
- 55 Clark J F 1907 *Botan Gaz* 33 26-48
- 56 Clayton C N 1942 *Phytopathology* 32 921-913
- 57 Cochrane V W 1945a *Cornell Univ Agr Exp Sta Mem* 268 1-30
- 58 Cochrane V W 1945b *Phytopathology* 35 361-366
- 59 Cochrane V W 1936 Unpublished data
- 60 Cole H Jr and C L Fergus 1936 *Phytopathology* 46 159-163
- 61 Cooper W C 1939 *Botan Gaz* 100 844-852
- 62 Corner E J H 1933 *New Phytologist* 34 180-200
- 63 Croser W 1934 *Cornell Univ Agr Exp Sta Mem* 133 1-40
- 64 Darby R T and G R Mandels 1935 *Plant Physiol* 50 360-366

- 65 Davies O L R B Duckworth and G C M Harris 1918 *Nature (London)* 161 612
- 66 Davis W H 1921 *Phytopathology* 14 251-267
- 67 Delp C J 1924 *Phytopathology* 44 615-626
- 68 Dimond A E and B M Duggar 1911 *Proc Natl Acad Sci U S* 27 429-468
- 69 Dodge B O 1912 *Bull Torrey Botan Club* 39 139-197
- 70 Domsch K H 1954 *Arch Mikrobiol* 20 163-175
- 71 Doran W L 1919 *Phytopathology* 9 391-40
- 72 Doran W L 1919 *Mass Agr Exp Sta Bull* 20 39-66
- 73 Doran W L 1920 *Bull Torrey Botan Club* 49 313-310
- 74 Duff G H 1918 *Phytopathology* 8 289-292
- 75 Duggar B M 1901 *Botan Gaz* 31 38-66
- 76 Duggar B M 1902 *U S Dept Agr Bur Plant Ind Bull* 85 1-60
- 77 Dunegan J C and C O Smith 1941 *Phytopathology* 31 189-191
- 78 Elliott E W 1949 *Mycologia* 41 141-170
- 79 Emerson M R 1948 *J Bacteriol* 55 327-330
- 80 Emerson M R 1924 *Plant Physiol* 29 418-428
- 81 Emerson R 1950 *Ann Rev Microbiol* 4 169-200
- 82 Evans M M and G Harrar 1930 *Phytopathology* 20 993-997
- 83 Ezekiel W N 1930 *Minn Univ Agr Exp Sta Tech Bull* 67 1-60
- 84 Felton M W and J C Walker 1946 *J Agr Research* 72 69-81
- 85 Fergus C L 1954 *Mycologia* 46 435-441
- 86 Fischer E and E Gaumann 1929 *Biologie der Pflanzenbewohnenden Parasitischen Pilze* Jena Gustav Fischer pp 4 8
- 87 Fischer E C and G Werner 1915 *Z Physiol Chem Hoppe Seyler's* 300 211-236
- 88 Fischer C W 1936 *Phytopathology* 26 1118-1117
- 89 Flor H H 1924 *Phytopathology* 44 469-471
- 90 Forles I L 1939 *Phytopathology* 29 659-684
- 91 Forsyth F R 1925 *Can J Botany* 33 363-373
- 92 Foster H M 1937 *Phytopathology* 27 803-803
- 93 Frick L 1913 *Phytopathol Z* 14 55-591
- 94 Fries N 1941 *Arch Mikrobiol* 12 266-284
- 95 Fries N 1943 *Symbolae Botan Upsalienses* 6 (4) 6-81
- 96 Fries N 1948 *Hereditas* 34 338-350
- 97 Fries N 1919 *Svensk Botan Tidskr* 43 316-31
- 98 Fromme F D 1915 *Am J Botany* 2 82-85
- 99 Fukushi T 1919 *J Coll Agr Hokkaido Imp Univ (Japan)* 15 269-307
- 100 Fulton H R 1906 *Botan Gaz* 41 81-108
- 101 Gardner M W 1918 *U S Dept Agr Bull* 727 1-68
- 102 Gardner M W 1924 *Proc India Acad Sci* 33 163-201
- 103 Gaumann E 1946 *Pflanzliche Infektionslehre* Basel Verlag Birkhauser pp 611
- 104 Gümman E S Naef Roth and H Kobel 1923 *Phytopathol Z* 20 1-38
- 105 Gehring F 1923 *Angew Bot* 27 97-105
- 106 Gilliver K 1947 *Ann Appl Biol* 34 136-143
- 107 Goddard D R 1932 *J Gen Physiol* 19 45-60
- 108 Goddard D R 1939 *Cold Spring Harbor Symposia Quant Biol* 7 362-376
- 109 Goddard D R 1948 *Growth* 12 (Suppl) 17-45
- 110 Goddard D R and P E Smith 1938 *Plant Physiol* 13 41-64
- 111 Goldsworthy M C and R E Smith 1931 *Phytopathology* 21 133-168
- 112 Gossop C H F Yuill and J L Yuill 1910 *Trans Brit Mycol Soc* 24 331-344
- 113 Gottlieb D 1950 *Botan Rev* 16 29-257

- 114 Gottlieb D 1953 *Intern Congr Microbiol 6th Congr* 5 122-136
- 115 Gould C J Jr 1945 *Iowa State Coll J Sci* 19 301-331
- 116 Graf Marin A 1934 *Cornell Univ Agr Exp Sta Mem* 157 1-48
- 117 Graves A H 1916 *Botan Ga* 62 337-369
- 118 Groom P and T Panisset 1933 *Ann Appl Biol* 20 633-660
- 119 Guba E F 1938 *Mass Agr Exp Sta Bull* 350 1-24
- 120 Gwynne Vaughan H C I 1937 *Ann Botany* 1 99-105
- 121 Gwynne Vaughan H C I and H S Williamson 1927 *Ann Botany* 41 489-495
- 122 Gwynne Vaughan H C I and H S Williamson 1933 *Trans Brit Mycol Soc* 18 127-134
- 123 Gwynne Vaughan H C I and H S Williamson 1934 *Ann Botany* 48 261-272
- 124 Hahne J 1925 *Kuhn Arch* 9 157-263
- 125 Haines R B 1930 *J Exp Biol* 8 379-388
- 126 Haines R B 1931 *J Exp Biol* 9 45-60
- 127 Hansen E C 1897 *Botan Ztg* 55 Abt I 111-131
- 128 Hansen H N 1938 *Mycologia* 30 442-455
- 129 Hart H 1926 *Phytopathology* 16 185-205
- 130 Hart M P and D M MacLeod 1955 *Can J Botany* 33 289-297
- 131 Hawn E J and T C Vanterpool 1953 *Can J Botany* 31 699-710
- 132 Heintzeler I 1939 *Arch Mikrobiol* 10 92-132
- 133 Hemmi T 1934 *Proc Pacific Sci Congr Pac Sci Assoc 5th Congr* 1933 4 3185-3194
- 134 Hemmi T and T Abe 1933 *Forsch Gebiet Pflanzenkrankheiten (Kyoto)* 2 1-9
- 135 Hesse R 1896 *Jahrb wiss Botan* 10 199-203
- 136 Holton C S 1943 *Phytopathology* 33 732-735
- 137 Hrushovetz S B 1956 *Can J Botany* 34 321-327
- 138 Huebschman C 1952 *Mycologia* 44 599-604
- 139 Hull R 1939 *Ann Appl Biol* 26 800-827
- 140 Huttig W 1931 *Z Botan* 24 529-577
- 141 Hyre R A and R S Cox 1953 *Phytopathology* 43 419-425
- 142 Johnson T 1941 *Phytopathology* 31 197-198
- 143 Jones E S 1923a *J Agr Research* 24 577-591
- 144 Jones E S 1923b *J Agr Research* 24 593-597
- 145 Jones F R 1919 *U S Dept Agr Bull* 759 1-58
- 146 Kaufmann F H O 1934 *Botan Gaz* 96 282-297
- 147 Kehl H 1943 *Planta* 33 731-737
- 148 Kett G W E C Blodgett E E Wilson and R O Magie 1937 *Wisconsin Univ Agr Exp Sta Research Bull* 137 1-117
- 149 Klotz L J and H S Fawcett 1932 *J Agr Research* 44 155-166
- 150 Kordes H 1923 *Botan Arch* 3 282-311
- 151 Kovacs A and E Szeoke 1956 *Phytopathol* 7 27 335-349
- 152 Krestlow K W 1943 *Phytopathology* 33 707-712 10 5-1063
- 153 Krestlow K W 1945 *Phytopathology* 35 152-158
- 154 Krishnan P S and V Bajaj 1953 *Arch Biochem Biophys* 47 39-55
- 155 Krishnan P S V Bajaj and S P Damle 1954 *Appl Microbiol* 2 303 308
- 156 Lambert E B 1929 *Phytopathology* 19 1-71
- 157 Latham D H 1935 *J Elisha Mitchell Sci Soc* 51 183-188
- 158 Leach J C 1923 *Minn Univ Agr Exp Sta Tech Bull* 14 1-41
- 159 Leben C 1954 *Phytopathology* 44 101-106
- 160 Lehman S G 1923 *Ann Missouri Botan Garden* 10 111-178
- 161 Le Roux P M and J G Dickson 1957 *Phytopathology* 47 101-107

- 162 Lesage P 1895 *Ann sci nat Botan Sér 3* 1 309-3 2
- 163 Lin C K 1940 *Cornell Univ Agr Exp Sta Mem* 233 1-33
- 164 Lin C K 1945 *Am J Botany* 32 296-298
- 165 Ling L 1940 *Phytopathology* 30 549-551
- 166 Ling L and J Y Yang 1944 *Ann Botany* 8 91-101
- 167 Longrée K 1939 *Cornell Univ Agr Exp Sta Mem* 223 1-43
- 168 Longrée K 1940 *Phytopathology* 30 793-807
- 169 Loptiore G 1895 *Jahrb wiss Botan* 28 531-6 6
- 169a Lowry R J A S Sussman and B von Böventer 1957 *Mycologia* 49 609-699
- 170 Lowry R J A S Sussman and B Heidenhain 1956 *Mycologia* 48 241-252
- 171 Lowther C V 1950 *Phytopathology* 40 590-603
- 172 McCallan S E A 1930 *Cornell Univ Agr Exp Sta Mem* 130 25-79
- 173 McCallan S E A L P Miller and R M Weed 194 *Contribs Boyce Thompson Inst* 18 39-68
- 174 McCallan S E A and F Wilcoxon 1936 *Contribs Boyce Thompson Inst* 8 151-165
- 175 McCrea A 193 *Science* 58 426
- 176 Machlis L and E Ossia 1953a *Am J Botany* 40 358-36
- 177 Machlis L and E Ossia 1953b *Am J Botany* 40 465-468
- 178 MacLachlan J D 1936 *J Arnold Arboretum (Harvard Univ)* 17 1-25
- 179 MacMillan H G and O A Plunkett 1947 *J Agr Research* 64 547-559
- 180 Magie R O 1935 *Phytopathology* 25 131-159
- 181 Mains E B 1915 *Ann Rept Mich Acad Sci* 17 136-140
- 182 Mains E B and H S Jackson 1916 *Phytopathology* 16 89-120
- 183 Mains E B and D Thompson 1918 *Phytopathology* 18 150
- 184 Mandels G R 1951 *Am J Botany* 38 213-221
- 185 Mandels G R 1953a *Arch Biochem Biophys* 42 164-173
- 186 Mandels G R 1953b *Arch Biochem Biophys* 44 367-377
- 187 Mandels G R 1953c *Exp Cell Research* 5 48-5
- 188 Mandels G R 1954 *Plant Physiol* 29 18-26
- 189 Mandels G R 1955 *Am J Botany* 42 91-929
- 190 Mandels G R 1956a *J Bacteriol* 71 681-688
- 191 Mandels G R 1956b *J Bacteriol* 72 230-234
- 192 Mandels G R and R T Darby 1953 *J Bacteriol* 6 16-26
- 193 Mandels G R H S Levinson and M T Hyatt 1956 *J Gen Physiol* 39 301-309
- 194 Mandels G R and A B Norton 1948 *Research Rept Quartermaster Gen Lab Microbiol Ser* 11 1-50
- 195 Mandels G R and A B Norton 1949 *Research Rept Quartermaster Gen Lab Microbiol Ser* 12 1-35
- 196 Maneval W E 1950 *Phytopathology* 40 471-487
- 197 Maneval W E 1954 *Phytopathology* 44 403-407
- 198 Marloth R H 1931 *Phytopathology* 21 169-198
- 199 Melhus I E 1911 *Wisconsin Univ Agr Exp Sta Research Bull* 15 1-91
- 200 Melhus I E and L W Durrell 1919 *Iowa State Coll Agr Exp Sta Research Bull* 49 115-144
- 201 Merek E L and C L Fergus 1951 *Phytopathology* 44 61-64
- 202 Miller H J 1949 *Phytopathology* 39 245-259
- 203 Miyoshi M 1895 *Jahrb wiss Botan* 28 69-989
- 204 Neger F W 1907 *Flora* 90 271 7
- 205 Noble R J 1953 *Phytopathology* 43 17-199
- 206 Noble R J 1954 *J Agr Research* 27 451-489
- 207 Oki N and T Hata 1952 *J Antibiotics (Japan)* 16-23

- 208 Owens R G 1955a *Contribs Boyce Thompson Inst* 18 125-141
- 209 Owens R G 1955b *Contribs Boyce Thompson Inst* 18 145-157
- 210 Padwick G W 1939 *Ann Appl Biol* 26 823-825
- 211 Page R M A F Sherf and T L Morgan 1947 *Mycologia* 39 158-164
- 212 Paley T 1936 *Arch Mikrobiol* 7 206-209
- 213 Parker Rhodes A F 1943 *Ann Appl Biol* 30 372-379
- 214 Peltier G L 1922 *Nebraska Univ Agr Exp Sta Research Bull* 20 1-15
- 215 Peltier G L 1925 *Nebraska Univ Agr Exp Sta Research Bull* 31 1-12
- 216 Perlman D 1951 *Am J Botany* 38 652-658
- 217 Platz G A 1928 *Iowa State Coll J Sci* 2 137-143
- 218 Platz G A L W Durrell and M F Howe 1927 *J Agr Research* 34 137-147
- 219 Powlesland R 1954 *Trans Brit Mycol Soc* 31 362-371
- 220 Pratt R 1956 *Am J Botany* 23 400-431
- 221 Raeder J M and W M Bever 1931 *Phytopathology* 21 167-189
- 222 Reed H S and C H Crabill 1915 *Virginia Agr Exp Sta Tech Bull* 9 1-106
- 223 Rippel A and H Bortels 1927 *Biochem Z* 181 237-244
- 224 Rippel A 1933 *Arch Mikrobiol* 4 530-547
- 225 Kishbeth H 1951 *Ann Botany* 15 1-21
- 226 Robbins W J 1940 *Am J Botany* 27 559-564
- 227 Robbins W J 1941 *Botan Gaz* 102 520-535
- 228 Robbins W J 1943 *Proc Natl Acad Sci U S* 29 201-209
- 229 Robbins W J and K C Hamner 1940 *Botan Gaz* 101 912-927
- 230 Robbins W J and F Kavanagh 1942a *Proc Natl Acad Sci U S* 28 4-8
- 231 Robbins W J and F Kavanagh 1942b *Proc Natl Acad Sci U S* 28 67-69
- 232 Robbins W J V W Kavanagh and F Kavanagh 1947 *Botan Gaz* 101 224-242
- 233 Robbins W J and M B Schmitt 1945 *Bull Torrey Botan Club* 70 16-85
- 234 Roberg M 1918 *Arch Mikrobiol* 14 1-11
- 235 Robinson W 1914 *Ann Botany* 28 331-340
- 236 Roper J A 1952 *Experientia* 8 14-15
- 237 Rosen H R and L M Weetman 1940 *Arkansas Univ Agr Exp Sta Bull* 391 1-20
- 238 Rother W 1954 *Arch Mikrobiol* 20 89-108
- 239 Ryan F J 1946 *Federation Proc* 5 366-369
- 240 Ryan F J 1948 *Am J Botany* 35 497-503
- 241 Saccas A 1951 *Rev path végétale et entomol agr France* 30 55-96
- 242 Sanford G B 1956 *Phytopathology* 16 505-517
- 243 Sansome E 1956 *Trans Brit Mycol Soc* 39 67-18
- 244 Sawyer W H Jr 1929 *Am J Botany* 16 87-121
- 245 Schaffnit E 1909 *Ann Mycol* 7 509-573
- 246 Schlosser L A 1959 *Planta* 8 529-570
- 247 Schneider R 1954 *Phytopathol Z* 21 63-18
- 248 Schonborn W 1955 *Arch Mikrobiol* 22 408-431
- 249 Schopfer W H 1935 *Compt rend soc phys et hist nat Geneve* 50 150-154
- 250 Semenuk G G M Nagel and J C Gilman 1947 *Iowa State Coll Agr Exp Sta Bull* 349 225-284
- 251 Shear C L and B O Dodge 1927 *J Agr Research* 31 1019-104
- 252 Shepherd C J 1956 *J Gen Microbiol* 15 29-38
- 253 Shepherd C J 1957 *J Gen Microbiol* 16 (1) 1

- 254 Shu P K G Tanner and G A Ledingham 1954 *Can J Botany* 32 16-23
- 255 Sisler H D and C E Cox 1954 *Am J Botany* 41 338-345
- 256 Sittler R G H 1951 *Microbial Decomposition of Cellulose* New York Reinhold Publ Corp pp 531
- 257 Smart R F 1937 *Am J Botany* 24 145-159
- 258 Smith E C 1959 *Mycologia* 21 321-323
- 259 Smith J H 1954 *New Phytologist* 23 65-78
- 260 Snell W H 1922 *U S Dept Agr Bull* 1053 1-47
- 260a Sorgel G 1956 *Phytopathol Z* 8 187-204
- 261 Sost H 1955 *Ach Potistenk* 100 541-564
- 262 Spaulding P and A Rathbun Gravatt 1956 *J Agr Research* 33 397-433
- 263 Stadler D R 1952 *J Cellular Comp Physiol* 39 449-474
- 264 Stakman E C 1915 *Minn Univ Agr Exp Sta Bull* 133 1-59
- 265 Stedman R L E Kravitz M Anmuth and J Harding 1956 *Science* 124 403-405
- 266 Stock F 1931 *Phytopathol Z* 3 231-249
- 267 Straub W 1940a *Zentr Bakteriell Parasitenk Abt II* 10 154-183
- 268 Straub W 1940b *Z Pflanzenkrankh* 50 579-585
- 269 Stroede W 1933 *Phytopathol Z* 5 613-614
- 270 Stuben H 1939 *Pflanz 30* 333-383
- 271 Sumi M 1958 *Botanica Z* 195 161-174
- 272 Sussman A S 1953a *Am J Botany* 40 401-401
- 273 Sussman A S 1953b *J Gen Microbiol* 8 211-211
- 274 Sussman A S 1954a *J Gen Physiol* 38 59-77
- 275 Sussman A S 1954b *Mycologia* 46 143-150
- 276 Sweet H R 1941 *Am J Botany* 8 150-161
- 277 Sweet H R 1942 *Am J Botany* 9 436-441
- 278 Tapke V F 1955 *Phytopathology* 45 3-18
- 279 Taylor C F et al 1955 *Phytopathology* 45 673-676
- 280 Thiel A F and F Weiss 1920 *Phytopathology* 10 448-459
- 281 Thumalachar M J and J G Dickson 1949 *Phytopathology* 39 333-339
- 282 Thomas H E and W D Mills 1959 *Cornell Univ Agr Exp Sta Mem* 13 1-71
- 283 Tilford P E 1956 *Ohio Agr Exp Sta Bull* 567 1-27
- 284 Tisdale W B 1950 *Phytopathology* 10 143-163
- 285 Togashi K 1919 *Biological Characters of Plant Pathogens Temperature Relations* Tokyo Meibundo pp 418
- 286 Tomkins R G 1959 *Proc Roy Soc (London) B* 10 375-401
- 287 Tomkins R G 1959 *Trans Brit Mycol Soc* 17 147-149
- 288 Turel F L M 1955 *Can J Microbiol* 1 293-298
- 289 Turner J N 1955 *Trans Brit Mycol Soc* 36 159-162
- 290 Uesaka I and Y Ito 1951 *J Antibiotics (Japan)* 4 418-424
- 291 Uppal B N 1956 *Phytopathology* 16 285-295
- 292 Vladimirova N N 1954 *Mikrobiologiya* 23 7-75
- 293 Wain R L and E H Wilkinson 1946 *Ann Appl Biol* 33 401-405
- 294 Wainwright L K 1956 *J Gen Microbiol* 14 533-544
- 295 Webb R 1951 *Ann Missouri Botanical Garden* 8 283-341
- 296 Weber G F 1959 *Phytopathology* 12 89-97
- 297 Wehmer C 1904 *Botanische Ges* 22 476-478
- 298 Werner J L and L L Hatter 1955 *J Agr Research* 4 1-40
- 299 Wellman R H and S E A McCallan 1949 *Contribs Boyce Thompson Inst* 1 431-440
- 300 Weston W H Jr 1951 *Trans N Y Acad Sci II* 13 305-308

- 301 Wilkins W H 1938 *Trans Brit Mycol Soc* 22 47-93
- 302 Wolf F A L F Dixon R McLean and F R Darkis 1934 *Phytopathology* 24 337-363
- 303 Wolf F A and F T Wolf 1917 *The Fungi* Vol 2 New York John Wiley and Sons pp 538
- 304 Wood Baker A 1935 *Trans Brit Mycol Soc* 38 291-297
- 305 Yarwood C E 1932 *Phytopathology* 22 31
- 306 Yarwood C E 1936 *Phytopathology* 26 845-859
- 307 Yarwood C E 1939a *Phytopathology* 29 828-829
- 308 Yarwood C E 1939b *Phytopathology* 29 933-945
- 309 Yarwood C E 1941 *Phytopathology* 31 863
- 310 Yarwood C E 1950a *Am J Botany* 37 636-639
- 311 Yarwood C E 1950b *Calif Agr* 4 7 12
- 312 Yarwood C E 1952 *Mycologia* 44 506-522
- 313 Yarwood C E 1954 *Proc Natl Acad Sci U S A* 40 374-377
- 314 Yarwood C E 1956a *Mycologia* 48 20-24
- 315 Yarwood C E 1956b *Phytopathology* 46 510-544
- 316 Yarwood C E and M Cohen 1949 *Science* 110 477-478
- 317 Yarwood C E S Sidky M Cohen and V Santilli 1954 *Hilgardia* 29 603-622
- 318 Yu C C 1954 *Am J Botany* 41 21-30
- 319 Zalokar M and V W Cochrane 1956 *Am J Botany* 43 107-110
- 320 Zeller S M 1920 *Ann Missouri Botan Garden* 7 51-74
- 321 Ziegler A W 1918 *J Elisha Mitchell Sci Soc* 64 13-40
- 322 Zimmerman A 1923 *Centr Bakteriell Parasitenk Abt II* 65 311-418

all cells are killed in 10 minutes has been determined for spores of several fungi as 40–60°C a value close to that usually reported for vegetative cells of mesophilic bacteria (1 19 21 24 88 128). However spores of some species are much more resistant known examples are the conidia of *Nocardia sebiivorans* (38) and ascospores of *Bysschlamys fulva* (70) *Neurospora crassa* (40) and *Penicillium* sp (153). Conidia of thermophilic actinomycetes appear to be heat resistant (37).

Ascospores are more resistant than conidia in *Neurospora crassa* (40 125) but are equally susceptible in *Endoconidiophora fagacearum* (24). Data on mycelium are too scanty for generalization. *Fusarium lini* is killed in 10 minutes at 55° but *Colletotrichum lini* survives 3 or 4 hours at this temperature (139). A culture of *Nocardia sebiivorans* withstands 10 minutes at 90° (38) and suspended sclerotia of *Penicillium* sp are completely killed at 82° only in about 16 hours (153). Spores of *Erysiphe* spp are somewhat more resistant to heat than mycelium (157) but comparative studies of spores and mycelium in other fungi have not been made.

Neither the thermal death point nor the related thermal death time has proved adequate in studies of bacteria (124) and there is no reason to hope that they will prove to be any more satisfactory for work with fungi. Development of a better index will however have to wait until we have more knowledge of the order of death. Characteristically the killing of vegetative cells of bacteria is exponential i.e. a plot of the logarithm of survivors against time is linear (118 124) although non-exponential orders of death by heat are known (77 119). Data on the killing of fungi by heat are few in number. Figure 1 shows a non-exponential order of heat killing of ascospores of *Bysschlamys fulva* ascospores of *Penicillium* sp respond similarly (153). At lower temperatures the curve becomes distinctly sigmoid i.e. short exposures have proportionately less effect (128). These few statements refer to studies of spores sclerotia of *Penicillium* sp respond to heat as one would expect in a multicellular structure (153).

Tentatively we may propose that the non-exponential order of death of fungus spores reflects genetic heterogeneity of the material but the data are too limited at present to justify a firm conclusion and in particular the occurrence of bi- or multinucleate cells may influence the reaction.

Death by heat is most drastically affected by moisture. Cells die much more rapidly at high temperatures if they are wet (22 37 74) or if the relative humidity is high (21 53). Heat resistance in *Bysschlamys fulva* is affected by the pH of the menstruum (70).

13 The Action of Physical Agents

It is convenient although probably not defensible in the final analysis to separate the physical and chemical agents that adversely affect fungi. Of the several agents of the first type attention must be focused upon temperature and radiation. Osmotic pressure has been touched upon earlier in Chapters 1 and 12. Hydrostatic pressure has hardly been studied in relation to the fungi. Very young infections by plant pathogens are prevented from developing by the application of pressure (156) and reduced pressure somewhat facilitates germination in *Erysiphe graminis* perhaps by accelerating carbon dioxide loss (12). It may be expected that studies of cellular reactions and of the action of fungicides under high pressures will contribute to our understanding of the fungi as similar studies have contributed to the knowledge of bacteria (76).

1 HIGH TEMPERATURE

The practical problem of heat sterilization is primarily that of killing bacterial endospores. Fungal structures with some exceptions are less resistant. Consequently there has been little interest in fundamental studies on the death of fungi at high temperature. However destruction of fungi within host tissues has proved practicable for a few pathogens which are present as mycelium in seeds or cuttings (148).

The thermal death point defined as the least temperature at which

all cells are killed in 10 minutes has been determined for spores of several fungi as 40–60°C a value close to that usually reported for vegetative cells of mesophilic bacteria (1 19 21 24 88 128). However spores of some species are much more resistant known examples are the conidia of *Nocardia sebiivorans* (38) and ascospores of *Byssochlamys fulva* (70) *Neurospora crassa* (40) and *Penicillium* sp (153). Conidia of thermophilic actinomycetes appear to be heat resistant (37).


Ascospores are more resistant than conidia in *Neurospora crassa* (40 125) but are equally susceptible in *Endoconidiophora fagacearum* (24). Data on mycelium are too scanty for generalization. *Fusarium lini* is killed in 10 minutes at 55° but *Colletotrichum lini* survives 3 or 4 hours at this temperature (139). A culture of *Nocardia sebiivorans* withstands 10 minutes at 90° (38) and suspended sclerotia of *Penicillium* sp are completely killed at 82° only in about 16 hours (153). Spores of *Erysiphe* spp are somewhat more resistant to heat than mycelium (157) but comparative studies of spores and mycelium in other fungi have not been made.

Neither the thermal death point nor the related thermal death time has proved adequate in studies of bacteria (121) and there is no reason to hope that they will prove to be any more satisfactory for work with fungi. Development of a better index will however have to wait until we have more knowledge of the order of death. Characteristically the killing of vegetative cells of bacteria is exponential i.e. a plot of the logarithm of survivors against time is linear (118 124) although non exponential orders of death by heat are known (77 119). Data on the killing of fungi by heat are few in number. Figure 1 shows a non-exponential order of heat killing of ascospores of *Byssochlamys fulva* ascospores of *Penicillium* sp respond similarly (153). At lower temperatures the curve becomes distinctly sigmoid i.e. short exposures have proportionately less effect (128). These few statements refer to studies of spores sclerotia of *Penicillium* sp respond to heat as one would expect in a multicellular structure (153).

Tentatively we may propose that the non exponential order of death of fungus spores reflects genetic heterogeneity of the material but the data are too limited at present to justify a firm conclusion and in particular the occurrence of bi or multinucleate cells may influence the reaction.

Death by heat is most drastically affected by moisture. Cells die much more rapidly at high temperatures if they are wet (22 37 74) or if the relative humidity is high (24 53). Heat resistance in *Byssochlamys fulva* is affected by the pH of the menstruum (70).

Figure 2 The influence of the rate of warming on viability (germinability) of conidia of *Aspergillus flavus* frozen at -70°C . Each curve represents a separate experiment. Re drawn from Mazur (103) by permission of the Rockefeller Institute for Medical Research.



the individual crystals (105). It follows that thawing must be equally rapid in order to prevent crystal growth. This is shown especially well by the data of Mazur (103) part of which is reproduced in Figure 2. The lethal effects of slow as compared to rapid warming were particularly evident in the range -20 to 0°C , i.e. the rate of subzero warming, not the rate of thawing of ice, is the critical factor.

The possibility of freezing material so rapidly as to prevent crystal formation entirely has been considered at length by Luyet and Gehenio (91). The experimental difficulties are formidable and there is some problem in determining whether or not success has been achieved (105). In any event it is clear that prevention of crystal formation is not essential to survival at low temperature.

We may conclude that fungi, especially vegetative mycelium and certain types of spores, are relatively susceptible to freezing injury, particularly to the slow freezing which occurs in nature. Repeated freezing and thawing are even more damaging (115). The principal factor is almost certainly the moisture content of the material; it follows that preservation in the frozen state will be most successful if applied to dry spores or other cells which can withstand a preliminary desiccation. Although it is unlikely that freezing can eliminate all cells of a given species, cold injury must be an important ecological factor at least for soil fungi and plant pathogens (22-39).

3 VISIBLE LIGHT

Light of the visible range $400-800\text{ m}\mu$, exerts undoubted but so far ill-defined influences on mycelial growth, spore germination and other phenomena. Effects on reproduction have been discussed in

assayed after thawing the material and as shown later (Figure 2) the rate of subzero warming is critical

Exposure to very low temperatures e.g. that of liquid air does not completely kill mycelium fruiting bodies or dry spores of most fungi (8 16 79 89). The mycelium of some fungi does not survive this treatment (89) and wet conidia of *Neurospora crassa* are killed almost immediately at -170° to -190°C (40).

Exposure to moderately low temperatures—from 0 to -40°C —is of immediate ecological interest and has frequently been investigated. The results however defy any neat generalization either fungi differ very widely in their tolerance or as suggested above methods of investigation are so different that comparison is not legitimate. Mycelium of *Phycomyces niteris* and *Botrytis cinerea* is killed by exposure to -15° to -22° (6) and neither mycelium nor sclerotia of *Phymatotrichum omnivorum* survives 21 hours at -13° (39). But mycelium of *Phacidium infestans* and *Collybia velutipes* is still viable after 138 days at -21° (114) and fruit bodies of *Schizophyllum commune* withstand exposure to -15 to -40° (17).

Similar differences are reported in the survival of spores at these moderately low temperatures. Uredospores of the rust fungi are usually found to survive no more than 2 months or so at -15° to -40° (22 42 101 149) but at least some uredospores of *Melampsora lini* are still viable after 1295 days at -10° (41) and uredospores of *Uromyces phaseoli* live as long as 2 years at -20° (59). Both spore forms of *Endoconidiophora fagacearum* are germinable after 83 days at -10° (24).

From these reports and from the partial list compiled by Luyet and Gehenio (91) it appears that spores are in general less damaged by low temperature than mycelium dry spores less than wet.

Most of the reported experiments on the true fungi are probably examples of relatively slow freezing in which the primary injurious factor is dehydration crystals of ice form extracellularly and withdraw water from the cell (105). Damage to dry cells is less than to wet (14 40) as would be expected. Furthermore it was early found that solutes protect the mycelium against freezing injury (6) hydrogen bonding substances—sugars glycols etc—essentially lower the amount of water lost to the crystallization process (105). Visual observation suggests that hyphal tips are more susceptible than basal mycelial cells and that aerial hyphae are more resistant than submerged (87) the water content may be a factor in these differences.

Rapid freezing in which crystal formation occurs throughout both cells and medium must obviously be so rapid as to limit the size of

4 ULTRAVIOLET RADIATION

The lethality and mutagenicity of ultraviolet radiation are problems of general biological interest studies on fungi have made significant contributions toward their solution The early and now largely obsolete literature is reviewed by Smith (127) more recent work by Pomper and Atwood (116) Techniques for the irradiation of spores and isolation of radiation induced mutants are well developed (7 120 138 152)

Virtually all studies of the action spectrum of killing and of mutagenesis in fungi as in other organisms agree that the most effective wavelength is in the region 260–265 $m\mu$ The action spectrum of lethality to *Trichophyton mentagrophytes* is shown in Figure 3 similar spectra prevail in *Neurospora crassa* (67 109) *Penicillium notatum* (69) *Ustilago zeae* (85) and *Aspergillus oryzae* (102) The primary peak is generally accepted to be consequent upon the specific absorption of this wavelength by nucleoproteins

McAulay and associates (92 93 94) have reported that biological activity for *Chaetomium globosum* is greatest at 280 $m\mu$ in the region of absorption by proteins These observations cannot be reconciled with any other data and merit re-examination

Aspergillus terreus conidia mutate after exposure to higher wavelengths (297 and 313 $m\mu$) or to sunlight (65) Efficiency however is




Figure 3 The action spectrum of killing of spores of *Trichophyton mentagrophytes* Re drawn from original of Figure 4 A Hollaender and C W Emmons *Journal of Cellular and Comparative Physiology* Vol 13 p 391 402 (1938) by permission of the Wistar Institute of Anatomy and Biology

Chapter 11 It must be recalled that exposure to white light may depending on the source involve exposure to some ultraviolet heating effects too cannot be disregarded unless specifically excluded

Several early observations, reviewed by Smith (127) and later data (20 58, 75 113 129) agree that light retards mycelial growth dry weight measurements confirm this for *Sclerotinia fructigena* (55) and *Marlingia rosea* (60) The growth of *Botrytis squamosa* on agar is completely inhibited by light (112)

Light stimulation of growth infrequently reported (60 127) has been most carefully analyzed in *Blastocladiella emersonii* (18) In this organism dry weight increases of up to 141 per cent at constant temperature are induced by illumination It is observed further that fixation of carbon dioxide is increased by light and it is proposed that light acts in some way to accelerate the reductive carboxylation of succinic acid and α -ketoglutaric acid


Vegetative mycelium like reproductive structures may be phototropic (122) and may show zonation in response to light and dark periods (25) Pigmentation is frequently dependent upon or enhanced by light (127) *Centrospora acerina* forms its red pigment after a very short exposure (107) Melanogenesis in *Neurospora crassa* is somewhat reduced by illumination (123) The effect of light on carotene formation has been mentioned elsewhere (Chapter 6)

It has been recognized at least since 1860 (61) that strong light inhibits spore germination Whether spores are more or less sensitive than mycelium is in doubt (127) Obviously any ultraviolet contamination is especially serious and there is no doubt that many of the lethal effects of light can be credited to a short wavelength component However in both *Erysiphe graminis* (117) and *Puccinia graminis* (30 31) experiments with filters implicate the longer wavelengths in the yellow red region and exclude ultraviolet Moist spores are more susceptible than dry (23)

A number of reports—too many to disregard—suggest that spore germination is increased by light or even requires light (17 48 71 106 130 155 160) The mechanism of such an effect is difficult to imagine and it can only be said at this time that further quantitative work under accurately controlled conditions is necessary

Photodynamic action has been demonstrated by Kaplan (78) in *Penicillium notatum* conidia vitally stained with erythrosin are sensitized so that visible light normally innocuous, becomes mutagenic Eosin stained cells of *Neurospora crassa* appear from limited data to respond similarly (34) Methylene blue intensifies the zonation response of *Penicillium notatum* to light (120a)

Figure 5 The survival of macroconidia (average number of nuclei 267 per conidium) of *Neurospora crassa* as a function of ultraviolet dosage. Redrawn from Atwood and Norman (5) by permission of the National Academy of Sciences.



species with dark spores tend to be more resistant (36-46) and a protective action of wall pigments in *Puccinia graminis* uredospores has been claimed (29). Pigmentation alone however cannot explain specific differences in sensitivity.

The curve generated by a plot of survival against dose is usually one of two types

1 The exponential curve in which the logarithm of the fraction surviving is a linear or approximately linear function of dosage. This curve is characteristic of microconidia of *Neurospora crassa* (Figure 4) and spores of *Penicillium* spp. (2-46), *Aspergillus melleus* (33) and *Streptomyces flaveolus* (80).

2 The sigmoidal curve (Figure 5) which is the more common type reported in studies of *Aspergillus* spp. (73-158), *Mucor dispersus* and *Rhizopus stolonis* (33), *Ustilago zeae* (85), *Glomerella cingulata* (95), *Fusarium eumartii* (126), *Trichophyton mentagrophytes* (64) and *Streptomyces griseus* (121).

The significance of these curve shapes has been debated vigorously and it may be too early for a categorical explanation. By itself the sigmoidal curve could be interpreted as expressive of variability in the spore population and Markert (95) has shown such variability in sensitivity in *Glomerella cingulata*. However this interpretation applied to the exponential curve requires a highly improbable distribution of sensitivity within the population. Studies on *Neurospora crassa*

very low and at still longer wavelengths most organisms survive very large doses (50) although bactericidal action can be detected at high intensity (62). The requirement for high intensity poses serious methodological problems in attempts to demonstrate the mutagenicity of the long ultraviolet (43). Very short ultraviolet—the Schumann region below $200\text{ m}\mu$ —has not been studied primarily for technical reasons; indications are that lethal action at least is very great (80, 111, 159).

The lethality of ultraviolet radiation is most easily expressed by the fraction of irradiated spores which fail to germinate. Doses that are not lethal delay germination or induce abnormalities (33, 61, 67, 85). Different fungi vary widely in susceptibility; thus spores of *Mucor dispersus* require almost 20 times the energy per spore that is sufficient for an equivalent degree of inactivation of *Aspergillus melleus* spores (32, 33). In surveys of fungal susceptibility it is often noted that




Figure 4 The survival of microconidia of three mutant strains (inositolless) of *Neurospora crassa* as a function of ultraviolet dosage. Redrawn from Gales (51) by permission of the Long Island Biological Association.

(152) Most important perhaps is the observation of McElroy (102) that postirradiation treatment with high hydrostatic pressure reduces the recovery of morphological mutants in *Neurospora crassa* and *Aspergillus terreus*. This observation suggests that the primary action of ultraviolet (and chemical mutagens) is to cause the formation of a semi stable intermediate which may decay to a new mutant state or may revert to its original normal state (103). The hypothetical intermediate may be the gene itself or a metabolite or metabolic system (143 146).

Peroxides and unknown products in irradiated media are mutagenic to *Neurospora crassa* (27 142). However results published so far fail to implicate products formed by ultraviolet irradiation of the medium at the wavelengths employed in mutation studies as important factors in the induction of mutations.

Photoreactivation was discovered in *Streptomyces griseus* (82) and has been observed in several of the true fungi (13 17a 52 81 109). As usually studied photoreactivation means that lethal effects of ultraviolet radiation are counteracted in part by subsequent exposure to light of longer wavelength 330 to 480 $m\mu$ however other changes induced by ultraviolet are also counteracted. The action spectrum of photoreactivation in *S. griseus* has a pronounced peak at 436 $m\mu$ differing significantly from the action spectrum of the same process in *Escherichia coli* (83). The theories and problems of photoreactivation are reviewed by Dulbecco (35). Wainwright and Nevill (144) present evidence that photoreactivation exerts independent effects on survival and mutagenesis in *Streptomyces* sp.

5 IONIZING RADIATIONS

Ionizing radiations as the name implies are radiations whose passage through a material causes the production of ion pairs. Most studies on fungi have employed X rays or the equivalent γ rays emitted by radium and other radioactive materials a few data are available on α -rays (from polonium but also produced by the cyclotron) β rays (from radioactive materials or from an ion source) and cyclotron produced neutrons. All these types of radiation we assume have the same mode of action but differ of course in energy ionization per unit path length and ability to penetrate (86). Consequently physically equivalent doses of different types of ionizing radiation are not biologically equivalent. However it is not yet clear how such differences in biological effect are to be explained. Thus in the killing of spores of *Aspergillus terreus* the densely ionizing radiations (protons and

(5 109 110) suggest instead that the shape of the dose response curve is a function of the number of nuclei per cell exponential for uninucleate spores sigmoidal for bi and multinucleate spores Death in a multinucleate spore requires that several targets must absorb radiation the sigmoidal curve follows from this requirement

Two mechanisms of lethality have been proposed (109) induction of recessive lethal mutations and non genetic damage to nuclei Obviously the first of these cannot be of much significance in the death of conidia having more than one nucleus

The induction of morphological and biochemical mutations by ultraviolet radiation responds rather differently as dosage is increased As shown in Figure 6 mutation frequency increases with dosage only to a point then levels off or declines at higher dosage Dose response curves of this general type have been reported from several studies in which all morphologically visible mutations are scored (63 66 67 73 100 135) However in one instance in which a known mutation at a single locus was scored the response curve was strikingly different (51) and the mutation frequency in *Streptomyces flaveolus* does not drop at high dosage (80)

Ultraviolet induced mutagenesis is affected by several modifying environmental factors Pretreatment before irradiation with infrared radiation 2 4-dinitrophenol or nitrogen mustard increases mutagenesis (133 134 135 136) Survival and mutation rate in *Streptomyces* sp are modified by treatments applied after irradiation (143 145 146) Respiratory poisons and anaerobiosis decrease both the lethal and the mutagenic effects of ultraviolet radiation on *Penicillium chrysogenum*




Figure 6 Morphological mutations in *Penicillium notatum* as a function of ultraviolet dosage Plotted from data of Hollaender (63)

Lethality and mutagenic action are not the only effect of irradiation. In particular irradiated spores may germinate but fail to make continued growth; this phenomenon reported in several fungi (11, 28, 90, 141) is analogous to the delayed effect of X irradiation on yeast (116) and to the microscopic mutant induced in *Chaetomium globosum* by ultraviolet or ionizing radiation (44, 93). The irradiation of *Aspergillus niger* spores with low (non lethal) intensities of α -particles results in the development of asporogenous colonies in an unspecified number (11). These delayed effects are perhaps best described as lethal mutations but something more than a descriptive analysis would surely be of interest.

Stimulation of spore germination by ionizing radiation has been reported (15, 154). Other less well-defined stimulatory effects claimed by earlier workers are reviewed by Smith (127).

The environmental factor most significant in modifying responses of fungi and bacteria to X irradiation is the ambient oxygen pressure (68, 131); both lethality and mutagenesis are less at low oxygen tension. Sensitivity of *Aspergillus terreus* to X irradiation is also increased by previous exposure to infrared radiation (133, 135) and by high spore moisture content (131). The lethality of X irradiation to *Streptomyces* sp. is decreased by various postirradiation treatments (147).

Radiation biology is undergoing rapid advances at this time and it is to be expected that generalizations will have a short life. In the fungi it appears that the primary effect of radiation is on the nucleus and this effect—which may be itself complex—is best visualized on the basis of the target theory so strongly urged by Lea (86). However there is good evidence—especially in the influence of environmental factors on response—that significant secondary reactions possibly in the cytoplasm must be postulated both for ultraviolet and for ionizing radiation (116, 159).

6 ELECTROMAGNETIC RADIATION OF LONG WAVELENGTH

Irradiation in the radio part of the electromagnetic spectrum may be inhibitory to fungi. Inhibition is however a heating effect and does not occur if the temperature is held constant (49, 97, 101, 140).

BIBLIOGRAPHY

- 1 Ames A. 1915 *Phytopathology* 5: 11-19
- 2 Arima K. 191a *J Antibiotics (Japan)* 4: 232-237
- 3 Arima K. 191b *J Antibiotics (Japan)* 4: 277-280

α rays) are more effective than the less densely ionizing λ rays and γ rays but the order of effectiveness is reversed if mutagenesis in *A. terreus* is the criterion (132). The killing of *A. niger* spores by α -radiation is similarly most effective at high ionization density (161). On the other hand lethality to vegetative cells of *Escherichia coli* is greatest with radiation of low ionization density (159).

Lethal and mutagenic effects of types of radiation other than λ rays have been reported viz neutrons (4 56 99) α particles from polonium and radium (9 10 11) γ rays from radioactive elements (26 137) and β particles from carbon 14 or an ion source (81 150 151). Earlier studies are reviewed by Smith (127).

The lethal effect of λ irradiation on spores of fungi and actinomycetes can usually be expressed as an exponential or approximately exponential dose response curve (3 72 80 121). Sigmoid survival curves have only occasionally been reported (45 90). It seems probable especially from preliminary work on yeasts that the form of the curve may be primarily determined by the number of nuclei per cell (116). This consideration cannot apply to the striking response of dry spores of *Aspergillus terreus*. λ irradiation follows the sigmoidal pattern but survival after irradiation with α particles is exponential (132 162).

Mutation frequency is linear with dose over a range as shown in Figure 7 some material shows a plateau in the curve at high dosage (54 80 108). Presumably this plateau will be reached with any organism provided the dose is high enough. Theoretical models for this saturation effect are considered by Harm and Stein (57).




Figure 7 Mutation frequency in *Streptomyces flavocolor* as a function of λ ray dosage. Each curve represents a separate experiment. Redrawn from Kellner (80) by permission of the Williams and Wilkins Company.

- 51 Giles N H Jr 1911 *Cold Spring Harbor Symposia Quant Biol* 16 283-313
- 52 Goodgal S H 1949 *Anat Rec* 105 496
- 53 Groom P and T Panisset 1933 *Ann Appl Biol* 20 633-660
- 54 Gwatkin R B and D Gottlieb 1916 *J Bacteriol* 71 328-332
- 55 Hall M P 1933 *Ann Botany* 47 543-578
- 56 Hanson H J W G Myers G L Stahly and J M Burkland 1946 *J Bacteriol* 51 9-18
- 57 Harm W and W Stein 1916 *Z Naturforsch* 11b 89-100
- 58 Harter L L 1939 *Am J Botany* 26 234-243
- 59 Harter L L and W J Zaumeyer 1941 *J Agr Research* 62 717-731
- 60 Haslins R H and W H Weston Jr 1950 *Am J Botany* 37 739-750
- 61 Hoffman H 1860 *Jahrb nuss Botan* 2 267-337
- 62 Hollaender A 1943 *J Bacteriol* 46 531-541
- 63 Hollaender A 1945 *Ann Missouri Botan Garden* 32 165-178
- 64 Hollaender A and C W Emmons 1939 *J Cellular Comp Physiol* 13 391-402
- 65 Hollaender A and C W Emmons 1946 *Cold Spring Harbor Symposia Quant Biol* 11 78-84
- 66 Hollaender A K B Raper and R D Coghill 1945 *Am J Botany* 32 160-160
- 67 Hollaender A E R Sansome E Zimmer and M Demerec 1945 *Am J Botany* 32 226-35
- 68 Hollaender A and G E Stapleton 1953 *Physiol Revs* 33 77-84
- 69 Hollaender A and E Zimmer 1945 *Genetics* 30 8
- 70 Hull R 1939 *Ann Appl Biol* 26 800-822
- 71 Hutchinson A H and M R Ashton 1930 *Can J Research* 3 187-198
- 72 Iguchi N 1911 *J Agr Chem Soc Japan* 24 81-84
- 73 Izuka H and T Yamaguchi 1952 *J Agr Chem Soc Japan* 26 71-14
- 74 Jensen H 1948 *Physiol Plantarum* 1 255-64
- 75 Jensen H L 1941 *Proc L nnean Soc N S Wal s* 66 216-86
- 76 Johnson F H H Eyring and M J Polissar 1954 *The Kinetic Basis of Molecular Biology* New York John Wiley and Sons pp 874
- 77 Jordan R C S E Jacobs and H E F Dav s 1947 *J Hyg* 40 136-143
- 78 Kaplan R 1950 *Planta* 38 1-11
- 79 Karcher H 1931 *Planta* 14 515-516
- 80 Kelner A 1948 *J Bacteriol* 56 457-460
- 81 Kelner A 1949a *J Bacteriol* 58 511-522
- 82 Kelner A 1949b *Pro Natl Acad Sci U S A* 30 73-79
- 83 Kelner A 1951 *J Gen Physiol* 34 835-852
- 84 Kern H and B D Sanwal 1954 *Phytopathol Z* 22 449-453
- 85 Landen E W 1939 *J Cellular Comp Physiol* 14 217-226
- 86 Lea D C 1955 *Actions of Radiation on Living Cells* 2nd ed Cambridge Cambridge Univ Press pp 416
- 87 Lindner J 1915 *Jahrb nuss Botan* 55 1-52
- 88 Ling L and E H Yu 1941 *Phytopathology* 31 64-210
- 89 L pman C B 1937 *Bull Torrey Botan Club* 64 537-546
- 90 Luyet B J 1932 *Radiology* 18 1019-1020
- 91 Luyet B J and P M Gehenno 1940 *Life and Death at Low Temperatures* Normandy Missouri Biodynamica pp 341
- 92 McAulay A L and J M Ford 1947 *Heredity (London)* 1 247-257
- 93 McAulay A L J M Ford and D L Dobie 1949 *Heredity (London)* 3 109-110
- 94 McAulay A L and M C Taylor 1939 *J Exp Biol* 16 474-482
- 95 McElroy W D 1951 *Science* 115 623-626

- 4 Atwood K C and F Mukai 1954 *Am Naturalist* 88 295-314
- 5 Atwood K C and A Norman 1949 *Proc Natl Acad Sci U S* 35 696-709
- 6 Bartetzko H 1910 *Jahrb wiss Botan* 47 51-98
- 7 Beadle G W and E L Tatum 1945 *Am J Botany* 32 678-686
- 8 Becquerel C 1910 *Compt rend (Paris)* 150 1437-1439
- 9 Berk S 1952a *Mycologia* 44 587-598
- 10 Berk S 1952b *Mycologia* 44 723-733
- 11 Berk S 1953 *Mycologia* 45 488-506
- 12 Brodie H J and J F Jones 1946 *Can J Research C* 24 318-329
- 13 Brown J S 1951 *J Bacteriol* 62 163-167
- 14 de Bruyn H L G 1926 *Phytopathology* 16 121-140
- 15 Buchwald C E and R M Whelden 1939 *Am J Botany* 26 718-731
- 16 Buller A H R 1912 *Trans Brit Mycol Soc* 4 106-112
- 17 Buller A H R and A T Cameron 1912 *Trans Roy Soc Can* 6 IV 378
- 17a Buxton E W F T Last and M A Nour 1957 *J Gen Microbiol* 15 764-773
- 18 Cantino E C and E A Horenstein 1956 *Mycologia* 48 777-799
- 19 Chaudhuri M 1923 *Ann Botany* 37 519-539
- 20 Chowdhury S 1944 *J Indian Bot Soc* 23 91-106
- 21 Cochran L C 1932 *Phytopathology* 22 891-812
- 22 Cochrane V W 1945a *Cornell Univ Agr Exp Sta Mem* 268 1-39
- 23 Cochrane V W 1945b *Phytopathology* 35 458-462
- 24 Cole H Jr and C L Fergus 1955 *Phytopathology* 45 159-163
- 25 Coons G H and F G Larmer 1930 *Papers Mich Acad Sci* 11 75-104
- 26 Davis R J V L Sheldon and S I Auerbach 1956 *J Bacteriol* 72 509-510
- 27 Dickey F H G H Cleland and C Lotz 1949 *Proc Natl Acad Sci U S* 35 581-586
- 28 Dickson H 1932 *Ann Botany* 46 389-403
- 29 Dillon Weston W A R 1931 *Sci Agr* 12 81-87
- 30 Dillon Weston W A R 1932a *Phytopathol Z* 4 229-246
- 31 Dillon Weston W A R 1932b *Sci Agr* 12 352-356
- 32 Diamond A E and B M Duggar 1940 *J Cellular Comp Physiol* 16 53-61
- 33 Diamond A E and B M Duggar 1941 *Proc Natl Acad Sci U S* 27 459-468
- 34 Doring H 1938 *Naturwiss* 26 819-820
- 35 Dulbecco R 1950 In A Hollaender (ed) *Radiation Biology* Vol 2 p 455-486 New York McGraw Hill Book Co
- 36 English H and F Gerhardt 1946 *Phytopathology* 36 100-111
- 37 Erikson D 1952 *J Gen Microbiol* 6 286-294
- 38 Erikson D 1955 *J Gen Microbiol* 15 127-135
- 39 Ezekiel W N 1945 *Phytopathology* 35 296-301
- 40 Faulk A F 1930 *Mycologia* 22 288-303
- 41 Flor H H 1954 *Phytopathology* 44 469-471
- 42 Forbes I L 1939 *Phytopathology* 29 659-684
- 43 Ford J M 1947 *J Gen Physiol* 30 211-216
- 44 Ford J M 1948 *Australian J Exp Biol Med Sci* 26 245-251
- 45 Ford J M and D P Karwan 1949 *J Gen Physiol* 32 647-653
- 46 Fulton H R and W W Coblenz 1929 *J Agr Research* 38 159-168
- 47 Gassner G and E Niemann 1954 *Phytopathol Z* 21 367-394
- 48 Gassner Hoechst G G 1954 *Phytopathol Z* 21 53-62
- 49 Gier L J 1938 *Trans Kansas Acad Sci* 40 55-57
- 50 Giese A C 1945 *Physiol Zool* 18 223-250

- 19 Tochizai Y 1926 *J Coll Agr Hokkaido Imp Univ (Japan)* 14 171-236
- 140 Tverskay D L 1937 *Plant Protect (U.S.S.R)* 1937 (13) 3-8 (*Rev Appl Mycol* 17 127-128 1938)
- 141 Uber F M and D R Goddard 1934 *J Gen Physiol* 17 577-590
- 142 Wagner R. P., C H Haddox R. Fuerst and W S Stone. 1950 *Genetics* 35 37-48
- 143 Wainwright S D and A. Nevill 1955a *Can J Microbiol* 1 416-426
- 144 Wainwright S D and A. Nevill 1955b *Can J Microbiol* 1 427-439
- 145 Wainwright S D and A. Nevill 1955c. *J Gen Microbiol* 12 1-12
- 146 Wainwright S D and A. Nevill 1955d *J Gen Microbiol* 12 13-4
- 147 Wainwright S D and A. Nevill 1955e *J Bacteriol* 70 547-551
- 148 Walker J C 1950 *Plant Pathology* New York McGraw Hill Book Co Inc. pp 699
- 149 Ward H M 1902 *Ann Botany* 16 233-315
- 150 Wheeler H E 1952 *Phytopathology* 42 431-435
- 151 Whelden R M 1940 *Mycologia* 32 630-643
- 152 Whittingham W F and J F Stauffer 1956 *Am J Botany* 43 54-60
- 153 Williams C. C. E. J Cameron and O B Williams 1941 *Food Research* 6 69-73
- 154 Woodward V W and C. M Clark 1955 *Science* 121 641-642
- 155 Yarwood C E 1939 *Phytopathology* 29 828-829
- 156 Yarwood C. E. 1953 *Phytopathology* 43 70-72
- 157 Yarwood C E S Sidky M Cohen and V Santilli 1954 *Hilgardia* 22 603-622
- 158 Zahl P A. L R Koller and C P Haskins 1939 *J Gen Physiol* 22 689-698
- 159 Zelle M R and A. Hollaender 1955 In A Hollaender (ed) *Radiation Biology* Vol 2 p 365-430 New York McGraw Hill Book Co
- 160 Ziegler A. W 1948 *J Elisha Mitchell Sci Soc* 64 13-40
- 161 Zirkle R. E 1940 *J Cellular Comp Physiol* 16 221-235
- 162 Zirkle R E D F Marchbank and K. D Luck. 1952 *J Cellular Comp Physiol* 9 Suppl 1 75-85

- 96 McElroy W D and C P Swanson 1951 *Quart Rev Biol* 26 348-363
- 97 Macdonald J A 1947 *Ann Appl Biol* 34 430-434
- 98 Machlis L and J M Crasemann 1956 *Am J Botany* 43 601-611
- 99 Markert C L 1952 *Genetics* 37 339-352
- 100 Markert C L 1953 *Exp Cell Research* 5 427-435
- 101 Matuo T and T Shiori 1951 *Research Repts Fac Textile Seric Shinsu Univ (Japan)* 1 47-52 (*Rev Appl Mycol* 34 234 1955)
- 102 Maxwell M E 1952 *Australian J Sci Research B* 5 56-63
- 103 Mazur P 1956 *J Gen Physiol* 39 869-888
- 104 Melander L W 1935 *J Agr Research* 50 861-880
- 105 Meryman H T 1956 *Science* 124 515-521
- 106 Neergaard P 1941 *Gartner Tidende* 57 319-322
- 107 Neergaard P and A G Newhall 1951 *Phytopathology* 41 1091-1093
- 108 Newcombe H B and J F McGregor 1954 *Genetics* 39 619-627
- 109 Norman A 1951 *Exp Cell Research* 2 454-473
- 110 Norman A 1954 *J Cellular Comp Physiol* 44 1-10
- 111 Oster R 1934 *J Gen Physiol* 18 71-88 249-250 251-254
- 112 Page O T 1956 *Can J Botany* 34 881-890
- 113 Page R M 1952 *Am J Botany* 39 731-739
- 114 Pehrson S O 1948 *Physiol Plantarum* 1 38-56
- 115 Piemeisel F J 1917 *Phytopathology* 7 294-307
- 116 Pomper S and K C Atwood 1955 In A Hollaender (ed) *Radiation Biology* Vol 2 p 431-453 New York McGraw Hill Book Co
- 117 Pratt R 1944 *Bull Torrey Botan Club* 71 131-143
- 118 Rahn O 1929 *J Gen Physiol* 13 19-205
- 119 Rahn O 1930 *J Gen Physiol* 13 395-407
- 120 Raper K B and D I Fennell 1946 *J Bacteriol* 51 161-177
- 120a Sagromsky H 1956 *Biol Zentr* 75 385-397
- 121 Savage G M 1949 *J Bacteriol* 57 429-441
- 122 Sawyer W H Jr 1929 *Am J Botany* 16 87-121
- 123 Schaeffer P 1953 *Arch Biochem Biophys* 47 359-379
- 124 Schmidt C F 1954 In G F Reddish (ed) *Antiseptics Disinfectants Fungicides and Chemical and Physical Sterilization* p 720-759 Philadelphia Lea and Febiger
- 125 Shear C L and B O Dodge 1927 *J Agr Research* 34 1019-1042
- 126 Smith E C 1935 *Bull Torrey Botan Club* 62 43-58
- 127 Smith E C 1936 In B M Duggar (ed) *Biological Effects of Radiation* p 889-918 New York McGraw Hill Book Co
- 128 Smith J H 1953 *Ann Appl Biol* 10 335-347
- 129 Snyder W C and H N Hansen 1941 *Mycologia* 33 580-591
- 130 Sorokine N 1916 *Ann sci nat Bot Ser* 6 3 46-52
- 131 Stapleton G E and A Hollaender 1952 *J Cellular Comp Physiol* 39 Suppl 1 101-113
- 132 Stapleton G E A Hollaender and F L Martin 1952 *J Cellular Comp Physiol* 39 Suppl 1 87-100
- 133 Swanson C P 1952 *J Cellular Comp Physiol* 39 Suppl 1 27-38
- 134 Swanson C P and S H Goodgal 1950 *Genetics* 35 695-696
- 135 Swanson C P A Hollaender and B N Kaufmann 1948 *Genetics* 33 499-457
- 136 Swanson C P W D McElroy and H Miller 1949 *Proc Natl Acad Sci U S* 35 515-518
- 137 Tarpley W J Harsky B Manowitz and R Horngan 1953 *J Bacteriol* 65 305-309
- 138 Tatum E L R W Barratt N Fries and D Bonner 1950 *Am J Botany* 37 38-46

1 THE RESPONSE TO TOXICANTS AND ITS MEASUREMENT

For practical reasons most of our information on the response of fungi to toxic agents relates to the inhibition of spore germination or of mycelial growth. However other effects are occasionally observed and may be briefly described. The most important aspect of response is of course its measurement since any investigation of mode of action presupposes an accurate reproducible assay of the biological effect of a fungicidal or fungistatic agent.

Morphological Responses Morphological abnormalities of growing hyphae are caused by many toxicants including such diverse agents as diphenyl (340) dyes (363) benzothiazole (142) solanin (2) and sodium tungstate (256). Other instances as well as data on abnormal development of germ tubes are tabulated by Horsfall (174). Thymol and carvacrol cause swelling of conidia of *Aspergillus niger* (450) as does phenol (420).

Griseofulvin, an antibiotic produced by *Penicillium* spp. (Chapter 6) causes stunting, branching, distortion and other abnormalities of growth in virtually all chitin-walled fungi tested, not in the yeasts or Oomycetes which have non-chitinous walls (Chapter 2). Brian (42) suggests that griseofulvin either acts as a growth regulator or interferes with a growth regulator controlling wall extension. Spore germination of *Botrytis allii* is not prevented by a concentration 500 times that which affects germ tube growth (43). Griseofulvin is taken up by green plants (74-403) and protects to a considerable degree against infection by *Alternaria solani* (44). A few fungi which probably have chitin walls are nonetheless insensitive (304).

Camphor causes the formation of large, possibly diploid conidia (361-362). Cholic acid and sodium taurocholate induce the formation of uninucleate cells in the dikaryon of several basidiomycetes, probably converting a clamp initial to a separate branch (286).

Mitotic Poisons Many fungitoxic substances are known to inhibit mitosis in other organisms (137-174). Exposure of basidia of *Gymnosporangium clavipes* to penicillin or streptomycin results in abnormal development and interference with meiosis (27). In yeast, penicillin inhibits cell division without inhibiting growth, so that filaments are formed. Reversal experiments suggest that sulfhydryl groups essential to cell division are blocked by the antibiotic (310).

The Response of Developmental Stages Specific inhibition of sporulation has both theoretical and practical importance. Simultaneous

14. The Action of Chemical Agents

Inhibition by chemical agents has probably been the single most active field of investigation of the fungi. Of necessity, however, much of this work has been directed to the practical problem of controlling the fungi pathogenic to plants and to man and other animals and the fungi that destroy economically valuable commodities. Theoretical problems have of course not been neglected; the important work has been summarized by Martin (278) and more recently by Horsfall (172-174).

In a study of the physiology of fungi such as this it seems appropriate to concentrate attention upon those materials regardless of their practical importance about which we have some information as to mode of action. This approach neglects many fascinating problems in the chemical inhibition of fungi: the role of antibiotic substances in soil (152-125) and the use of purified antibiotics (218-337-396-430); the expanding field of chemotherapy of plants (90-175-176-402); and many of the developments in the use of organic compounds as fungistatic or fungicidal agents. It is to be hoped that sufficient attention will be devoted to mode of action so that these topics can be treated theoretically in the near future.

The chapter has been organized so as first to consider some general problems and then to give detailed attention to a few types of compounds—heavy metals, sulfur, quinones, and the chelating agents—which have been most actively investigated.

under these conditions although other explanations are possible (20) Increase in oxygen consumption is also caused by poisons e.g. 2,4-dinitrophenol which interfere with the utilization of respiratory energy (380)

We may conclude that in general the activity of fungicides against respiration does not parallel their effects on growth and infer that effective fungicides usually act on systems much more sensitive than is the gross oxygen uptake We may speculate that a compound which acts on respiration and growth at approximately the same concentration—mercuric chloride is an example (247)—is a general enzyme poison It should also be clear that although a respiratory system may be inhibited by a fungistatic agent this fact alone is not sufficient evidence that the inhibition observed is the primary locus of toxicity

Sclerotinia laxa grown in contact with partially inhibitory concentrations of several fungicides is affected physiologically in its formation of respiratory enzymes some being increased and others decreased in amount by specific compounds (55)

Mutagenic Effects of Chemical Agents Cultural variants of undetermined genetic status are induced by high levels of zinc (89) and nitrite (Chapter 8) Mutation in fungi verified by genetic studies is caused by a wide variety of chemicals e.g. nitrogen and sulfur mustard and related compounds (16 286a 400 409) peroxides (87 188) epoxides (208) diazomethane (189) and caffeine (115) Some of these are sulfhydryl reagents others possibly generate free radicals but no single mode of action can be proposed at present

Fungistatic Action of Chemical Agents Both practically and theoretically the most important effect of toxicants on fungi is the inhibition of growth

This fungistatic action is the most frequently measured and should not be confused with killing fungicidal action *sensu stricto* We may distinguish two classes of fungistatic assay (1) determination of mycelial growth on a natural or artificial medium and (2) determination of spore germination In both methods the test fungus and the toxicant are in contact throughout the period of incubation

The early development of agar plate methods is described by Bateman (22) who established a satisfactory quantitative treatment of the data obtained with *Fomes annosus* This treatment has since been extended to other fungi (330 42b) After a more or less protracted lag period which is itself an index of toxicity the rate of growth across the agar surface is linear with time Comparison with a control inoculation on toxicant free agar yields the basic datum degree of inhibition




Figure 1 The response of germination (curve 1) and oxygen uptake (curve 2) of *Neurospora sitophila* conidia as a function of the concentration of cycloheximide. Respiration was measured in the presence of sucrose germination after removal of the toxicant. Redrawn from McCallan, Miller and Weed (214) by permission of the Boyce Thompson Institute for Plant Research Inc.

and fully quantitative determinations of both growth and sporulation are difficult preliminary data suggest that the high metal requirements of sporulation (Chapter 11) make the process especially sensitive to chelating agents e.g. dimethylglyoxime but many other types of toxicant also reduce reproductive activity (179 332 333 340 342 399 428 448 457)

It is generally believed that mycelial cells are less sensitive to toxicants than are spores this is borne out by direct comparisons for sulfanilamide (125 147) and cycloheximide (430a 431). The difference may however be related to uptake (246a)

Inhibition of Respiration Inhibition of respiratory activity as measured by oxygen uptake is often parallel to growth inhibition (106 269 309 334) and it has even been tentatively suggested that the principal activity of Captan¹ is on pyruvic carboxylase (160). However most studies on very different compounds show that in general endogenous and substrate respiration is much less sensitive to a wide range of fungicides than is spore germination or growth (83 125 127 132 203 224)

Figure 1 is drawn from an extensive study of fungicidal activity judged by killing of spores and spore respiration the figure shows that a compound in this case cycloheximide kills at concentrations which do not measurably reduce oxygen uptake. The data of Kloppe (203) show a similar relation between fungistatic action and inhibition of respiration. 8-hydroxyquinoline has no effect on respiration at a concentration 100 times the fungistatic concentration.

Low concentrations of toxicants often stimulate respiration (247). It seems likely that membrane permeability to substrate is increased

¹ (trichloromethylthio) 4-cyclohexene-1,2-dicarboximide

than they are against the growth or metabolism of mycelium (125 147 271 341 430a 431)

Fungistatic methods adapted for the assay of wood preservatives are reviewed by Cartwright and Findlay (61) The type of assay chosen has marked effects on the apparent efficacy of test compounds (110 223 348) Usually loss in weight of treated wood blocks is the best criterion of protection against decay a somewhat more rapid assay based on the static bending strength of a treated wood block has been introduced (126)

Special methods for fabric treatments are described by Siu (386) Soil fumigants are usually of interest insofar as they kill organisms but a fungistatic method of bioassay has been devised (101) For the practical testing of plant protectants assays employing living inoculated plants are well worked out (243 245 251 280)

The most progress has been made by the use of bioassays which are based on inhibition of spore germination This fundamental technique has been developed both practically and theoretically by McCallan and by Horsfall whose papers must be read for a thorough understanding of the method (172 173 174 241 242 243 252 254 258 259 436 44b) More or less standard procedures have been agreed upon (71 72)

A standardized spore suspension of a test fungus—*Monilinia fructicola* (= *Sclerotinia fructicola*) and *Stemphylium sarcinaeforme* (= *Macrosporium sarcinaeforme*) are the most often used but other sporulating fungi are suitable—is allowed to germinate in contact with the toxicant which may be either mixed with the spore suspension or sprayed on the slide (33) If washed or vacuum collected spores are used it is usually necessary to add nutrients and these materials especially complex plant products such as orange juice must be taken into account in any consideration of the mode of action orange juice possibly by reason of its content of citric acid which can complex metals reduces both the fungicidal and the fungistatic effect of copper on at least many fungi (91 250 287)

In the spore germination assay a simple plot of per cent inhibition (or germination) against toxicant concentration or the logarithm of toxicant concentration yields a sigmoid curve This can be converted to a straight line in the so called logarithmic probability system of coordinates The abscissa of this curve used in most studies is the logarithm of the dose the ordinate a calculated probit or normal equivalent deviation Use of this coordinate system is based on the general phenomenon that resistance in a population to a toxic agent is distributed normally with respect to the logarithm of the dose (30




Figure 2 The inhibition of *Aspergillus niger* by *p* hydroxy benzoic acid in an agar medium. Drawn from data of Vincent (426)

This is related to toxicant concentration by the equation

$$y = ax^b \quad (1)$$

where y stands for degree of inhibition x for concentration of the fungicide and a and b are constants. A plot of the logarithm of the degree of inhibition against the logarithm of the concentration yields curves of the type shown in Figure 2. Not all curves are as clearly linear as this one (426).

A variety of special methods are essentially similar to the assay of Bateman just described (26 102 108 109 114 151 203 271 283 296 297 317 318 359). Criteria used include the least concentration preventing all growth the colony diameter at a fixed time or the dry weight in liquid culture. A rather different approach is found in the use of filter paper discs impregnated with the toxicant (218 415) here of course, diffusion becomes an important factor.

These methods all have particular and limited value thus the medical mycologist has a practical interest in the inhibition of established infections rather than in prevention of spore germination. Fungi which do not sporulate in culture may have to be studied as mycelium. For theoretical work on mode of action however, the presence of nutrients and, in agar media, of undetermined contaminating substances severely restricts their value. It is almost always found that nutrient materials reduce the apparent toxicity of fungitoxic agents (214 352 404). In a qualitative way it is usually true that the agar plate method and the spore germination method agree but they may give divergent results for particular compounds (174). As a rule fungitoxic materials are more active against spore germination and spore respiration

difference between the concentration which inhibits the most sensitive spore and that required to prevent germination of the most resistant spore. Conversely a low slope or flat dosage response curve indicates that the difference in dose required for inhibition of the two extremes of resistance is large (446). Slope is most conveniently measured as the reciprocal of the difference between ED_{84} and ED_0 .

Although several factors of the assay influence slope the compound itself is the most significant (252). From this it has been argued that the slope is an indication of the fundamental mode of action (174, 241, 279, 346, 416). That is compounds with the same mode of action should generate dosage response curves of the same slope. Rich and Horsfall (346) show that homologues of 2-imidazoline yield curves of similar slope and different position: the ED_{50} varied from 0.0037 to 3.57 μ M per square centimeter but the slopes were essentially alike—9.37, 11.01 and 9.97. From this and other experience it is argued that position measures ability of the compound to reach its site of action and that slope indicates the mode of action once the site has been reached.

Attractive as this hypothesis is there are other factors that must give us pause. First as has been mentioned non linear curves are not uncommon. Second there is as yet no measure of mode of action which is truly independent and which could serve as a check on the hypothesis. Finally virtually all the available data are dosage response curves in which external concentration is the independent variable. The discovery that fungi accumulate very large amounts of toxicant (p. 450) does not by itself defeat the hypothesis but this fact does necessitate a re evaluation of the whole problem (246).

Spores of many fungi increase markedly in size during the first stages of germination (Chapter 12). Inhibition of this growth process in *Myrothecium verrucaria* has been adapted to the bioassay of fungistatic agents (268).

Fungicidal Action Fungicidal action the killing of fungi by a toxicant requires an assay method in which cells are exposed to a compound for a known period of time and are then removed from contact with the toxic agent and tested for viability. Such assays have been worked out for a variety of theoretical and practical problems (26, 69, 120, 128, 151, 246a, 250, 360, 463).

The plot of probit of per cent germination against logarithm of time is linear for the action of copper, silver, arsenate and crystal violet (250). The same plot of the data of Muller and Biedermann (302) shows however that the strict exponential relation is evident only at




Figure 3 Dosage response curves for the action of heavy metals on *Stemphylium sarciniforme*. Ordinate is probit of per cent inhibition of germination; abscissa is logarithm of concentration. Redrawn from more extensive data of McCallan, Wellman, and Wilcoxon (25^o) and selected to show different types of curve. Reproduced by permission of the Boyce Thompson Institute for Plant Research, Inc.

174 244 446) Representative curves of this type appear in Figure 3.

Three characteristics of the log probit curve are of greatest interest: position, shape, and slope. Lateral position is of course indicative of absolute toxicity; either the ED_{50} or the ED_{95} —the concentration required to inhibit germination by respectively 50 and 95 per cent—may be determined from the curve and used in the rating of fungicides. Position is affected by several factors in the assay—the presence of nutrients, the age of spores, the concentration of spores, extremes of temperature, the germination time, and the physical or chemical form in which a toxicant is supplied (19, 91, 144, 174, 252, 260, 265, 436).

Non-linear curves are not, however, uncommon; thus a detailed study of the action of copper sulfate on *Monilinia fructicola* and *Alternaria oleracea* shows for both fungi a J-shaped curve, concave upward (242). Parker Rhodes (323) proposed that this curvature represents the cumulative effect of more than one chemical or physical process in the inhibition reaction and also detailed an exponential transformation of the independent variable which converts the J curve to a linear form. Use of the transformation does not require acceptance of the multi-process theory of fungistatic action, some of the difficulties of which are discussed by McCallan (214). Non-linear curves of other types are frequently encountered (242, 244, 252).

The slope of the log probit curve is its most interesting feature. Since the ordinate is based on the distribution of sensitivity in the population of spores, a steep slope means that there is relatively little

general exclusion of cations from the cell As Horsfall (174) points out an organism producing metabolic acids may be less sensitive to a toxicant which must compete with hydrogen ion for its site of action

Fungi may also be trained by serial culture in media containing low doses of a toxicant to tolerate a normally inhibitory concentration. Such adaptation has been demonstrated to arsenite (157) borax (205) fluoride (222) and copper (186-195). Other reports are cited by Leben et al. (217) and Horsfall (174). Such adapted strains and strains which appear spontaneously in culture are usually unstable and lose their resistance on subculture in toxicant free media (158-195). Spontaneously arising sectors of *Fusarium caeruleum* appear in plates exposed to the vapor of 2,3,5,6-tetrachloronitrobenzene and these variants are somewhat less sensitive to the poison than is the parent strain (263); they are however almost equally sensitive—by rather inexact criteria—to 2,3,4,5-tetrachloronitrobenzene (47).

Genetic analysis has been performed only infrequently on resistant strains of fungi. Resistance to indole toxicity in *Neurospora crassa* (77) and to acriflavine in *Aspergillus nidulans* (355) is genetically controlled. This is in line with the usual experience in studies on bacteria and yeast (85-216-371). Copper adapted yeast is reported to contain a ribonucleic acid which can be transmitted to and which confers copper resistance on normal unadapted yeast cells (293).

Physiological mechanisms of acquired tolerance are possibly of diverse types. Work on the resistance of trypanosomes to arsenicals suggests that resistant strains are less permeable to the drug (38) possibly by virtue of an altered distribution of surface charges (366). Fluoride tolerance in clones of *Propionibacterium pentosaceum* appears also to be a permeability phenomenon (427). Other mechanisms are however possible: an antimycin resistant variant of *Venturia inaequalis* probably lacks the normal respiratory system sensitive to the antibiotic (219).

Detoxification of fungicidal agents has been observed and is of course a potential mechanism of resistance. Methylation of selenium tellurium and arsenic (Chapter 6) conceivably fits in this category. The effects of cycloheximide on both spore germination and respiration decline with time indicating an inactivation (431-440). Oxidative processes have been suggested as the basis for resistance to phenols and quinones (200-347). β -oxidation of a fatty acid side chain or hydroxylation of a ring in some instances causes detoxification (54-57). Other possible detoxification mechanisms are reviewed by Gottlieb (134); it seems likely that the most common is the complexing of toxic metals by chelate forming metabolites such as amino acids and hydroxy acids.




Figure 4 Dosage response curves for the action of uranium acetate on spores of *Alternaria solani*. Curve 1 fungistatic action curve 2 fungicidal action at 20 hours curve 3 fungicidal action at 10 minutes. Redrawn from McCallan and Wellman (20) by permission of the Boyce Thompson Institute for Plant Research Inc.

relatively high concentrations of copper—at lower concentrations there is no perceptible damage for a considerable period of time after which the exponential form of curve appears. Similarly the shape of the time inhibition curve for the action of phenol is affected by the concentration of the toxicant (391).

Killing is also a function of temperature: the fungicidal action of a number of relatively mild materials is sharply and non-linearly increased as the temperature is raised, with an especially sharp break as the temperature approaches its lethal value for spores in the absence of toxicant (183).

Although a strongly fungicidal agent is fungistatic the reverse is not necessarily true. In particular several compounds of high fungistatic activity—potassium dichromate, uranyl acetate, imidazolines and 8-hydroxyquinoline are examples—have only very weak fungicidal activity (250–437). In general dosage response curves for fungicidal action are much flatter than those for fungistatic, i.e. the range in resistance of the population is greater (Figure 4).

2. TOLERANCE AND RESISTANCE

Species of fungi differ widely in their sensitivity to toxicants. This has been especially noticed since the introduction of antibiotics and organic fungicides: differences in sensitivity of 50 to 100 fold are common (203–270–349–440). However strains tolerant to more general poisons are also encountered in studies on mercury (124–232–423), fluoride (227–423), arsenate (303) and creosote (65). Copper tolerance is often associated with tolerance to hydrogen ion or other cations (195–387–397). This suggests that the mechanism of copper tolerance is a

that spores of *Monilinia fructicola* exposed to copper accumulate the metal to a concentration 2000–4000 times its external concentration

Even more striking results are reported by Miller McCallan and Weed (290–291) using different fungi and such diverse materials as silver cerium 2 heptadecyl 2 imidazoline and ferric dimethyldithio carbamate. Several features of these data warrant emphasis

1 Uptake is extremely rapid spores of *Neurospora sitophila* take up their limit of the imidazoline within 15 seconds from a dilute solution within 5 minutes from a more concentrated solution. Uptake of dithiocarbamates by yeast is similarly rapid (127)

2 Fungicidal action is a linear function of the amount of toxicant taken up

3 Some toxicants are concentrated as much as 10 000 fold

4 A material which has low inherent toxicity on a spore weight basis may be so extensively concentrated by spores as to be an effective inhibitor. The application to bioassay is obvious—external concentration is not necessarily a measure of effective dose. Valko (419) distinguishes apparent toxicity a function of external concentration from specific toxicity a function of the amount of toxicant absorbed by the cell

5 Species differ in the amount taken up of a given toxicant.

6 Unrelated toxicants are taken up independently i.e. previous ex




Figure 5 The relation between uptake and antifungal activity (of the organism *Neurospora sitophila*) in chlorone derivatives of 4-dichloro-6-anilino-1,3,5-triazine. Curve 1 fungicidal action curve 2 fungistatic action. Compounds u = unsubstituted o = ortho-chloro m = meta-chloro p = para-chloro derivative. Drawn from data of Burchfield and Storrs (51)

3 SELECTIVE ACCUMULATION

We have customarily thought of the inhibiting of an unwanted organism as a problem in selective toxicity to find materials which are inherently more toxic to one form of life than to others. It has been assumed that the ultimate basis for such a selective action must be the existence of some sensitive protoplasmic reaction in the undesirable organism which does not exist or is dispensable in other species. This approach has of course had its successes notably in explaining the chemotherapeutic value of the sulfonamide drugs but the major effort of modern biology has been to show how similar are all forms of life in fundamental biochemical aspects and thus to render less and less likely the prospect of finding truly selective agents at the protoplasmic level of action.

It now appears that another selective principle may be equally important the differential accumulation of a toxicant by the unwanted organism. The term selective accumulation has been used by Yarwood (460) to describe the finding that spores of *Uromyces phaseoli* accumulate more sulfur from a fungicidal treatment with the vapor phase than do the leaves of the host plant. That is we may hope that some materials which are perhaps equally toxic to all organisms may be accumulated selectively by a fungus the result of course is equivalent to selective toxicity and is a form of it but does not depend on an inherent protoplasmic or enzymatic sensitivity. Selective accumulation does of course require that the sensitive organism differ physiologically from the insensitive but it does not require us to postulate for example that protein synthesis or energy metabolism of the susceptible species differs from that of the resistant. The problem becomes narrower and more manageable by what mechanism does the sensitive organism accumulate or the insensitive exclude the toxicant?

In the previous section some evidence has been cited that resistant strains of microorganisms owe their resistance to the failure of the toxicant to reach sensitive sites. This is in a sense the reverse of selective accumulation to be considered provisionally as selective exclusion.

The idea of selective accumulation is not of course new it has been known for some time that the lethal action of heavy metals on bacteria follows upon accumulation of the toxicant by cells in amounts far in excess of their concentration in the external medium (155). It is apparent that fungus spores can similarly remove from the medium very large amounts of many different toxicants. Marsh (273) found

4 SURFACE ACTION

It has been suggested immediately above that the first phase of toxicant action is an accumulation of the active material in a peripheral layer or zone outside the semi permeable membrane. We have further argued that, for most toxicants the primary inhibitory effect is not at this stage but later after some of the toxicant has desorbed from the outer zone and penetrated the protoplasm proper. However such a sharp separation of phases is only a convenience since accumulation of a foreign material is very likely to have some effects on the cell membrane. Hypothetically we may set up three possible classes: (1) effects of the sorbed toxicant are themselves lethal; (2) sorption contributes to lethality by damaging the osmotic barrier and thereby facilitating or making possible entrance of the toxicant into the protoplast; and (3) sorption has only minor effects.

A number of surface agents are known to be lethal to bacteria at least in part as the result of destruction of the semi permeable membrane (118-181). The antibiotic polymyxin has the same effect—disruption of the osmotic barrier—in bacteria (307). The contribution of lysis to lethality has however been questioned (398).

Surface toxicity need not in principle at least be confined to effects on permeability. If surface anabolic reactions are essential to growth for example inhibition of them by a non penetrating toxicant would be sufficient. Finholt et al. (110) interpret the different toxicity of aliphatic amines in malt agar culture vis a vis wood block tests as indicating that a toxicant could act externally by preventing the action of an extracellular hydrolytic enzyme essential to survival. This suggestion deserves further exploration perhaps the most promising possibility is that the extracellular enzymes are actually synthesized exteriorly to the osmotic barrier and that a toxicant might interfere with the synthesis. Such an effect would be missed completely in the usual assay which provides diffusible nutrients and thereby renders the fungus independent of extracellular enzymes.

The second possibility that sorption of a toxicant might contribute to lethality without being the primary event comes to mind particularly in relation to a few findings that toxicants at sublethal concentrations alter cell permeability. Cellular phosphorus is lost from spores treated with subfungistatic doses of silver (288) or of substituted triazines (51). Lysozyme treatment causes an increase in permeability to water and probably to furfural (235). In bacteria phenol increases permeability to glucose it is probable that one way effects on per-

posure to one material does not materially change the results with another. Closely related chemicals appear however to compete for a common receptor (246, 288).

7. Absorbed toxicants of some types are held firmly in or on the spore and can be released in quantity only by rather drastic treatment. Silver by contrast is easily removed by exchange (292).

The relation of uptake to effectiveness is shown in Figure 5 for a series of related s triazines. Fungitoxic action is seen to be correlated with the capacity of the spore to remove the toxicant from solution.

In general two mechanisms suggest themselves as alternative explanations of the phenomenon of rapid uptake against a concentration gradient. First it is of course known that active transport of nutrient materials into cells occurs commonly and results in accumulation against a gradient and Pramer (336) has shown that streptomycin is so accumulated by *Nitella clavata*. However to suggest active transport as the mechanism of entrance of a variety of toxicants raises the difficulty that active transport as usually understood necessarily involves the expenditure of metabolic energy and is prevented by many inhibitors. As discussed later in connection with the heavy metals it seems at present more promising to look toward a second mechanism, the adsorption of the toxicant by a solid phase in or on the cell. For both these alternative mechanisms it should be noted the same general relation between uptake and external concentration holds in both: there is some saturation value of the external concentration. Consequently whether or not the concentration uptake relation is described by the adsorption isotherm does not distinguish between active transport and adsorption. More important results will come from studies of the effect of metabolic state upon uptake.

Drawing primarily from studies on the action of heavy metals (see below) we may suggest that a toxic material which does not move freely across the semi permeable membrane undergoes a two stage process. In the first phase there is a rapid sorption of the toxicant outside the permeability barrier or perhaps both outside it and on it. In the second phase the material is released at a slower rate to the interior of the cell from the saturated peripheral sites; only in this phase, for most materials, does interference with growth occur.

This hypothesis has many difficulties which are however better considered in connection with heavy metal toxicity. It ignores for convenience the possibility that the sorbed toxicant may exert specific effects before it enters the cell at all; this possibility is treated as a separate topic.

The negative logarithm of the ionization constant of Equation 2 signified by pK_a , is numerically equal to the pH at which the acid is 50 per cent dissociated. The pK_a is characteristic of the molecule and one of the commonly overlooked effects of addition of substituents to an organic molecule is that the substituent may change pK_a . Exactly the same considerations of course apply to weak bases.

Acidic toxicants almost invariably are more effective at low pH than at high as discussed later the effect of pH is most noticeable near pK_a so that data on very strong and very weak acids will not normally show an effect of hydrogen ion concentration under the usual experimental conditions. Data on inhibition of fungi which fit the rule of greater activity at low pH are extremely numerous although they do not include spore germination results. As examples we may cite the action of fatty acids (161 350 354 456) aromatic acids (75 76 78 162 339 390) acid dyes (205) substituted phenols (41 187 383) and nitrous acid (Chapter 8).

The usual explanation for the effect of dissociation on toxicity is that only the undissociated molecule enters the cell. This reflects the general experience that ions penetrate the cell at least much more slowly than do neutral molecules (84). Several experiments with fungi have shown that the ester of a toxic acid is more effective than the free acid (24 62 140 178 301). The ester almost certainly penetrates the cell more rapidly than the partially ionized acid and is then split by esterases. More direct evidence would however be desirable since esterification of an acid may have other effects for example on reactivity of groups near to the carbonyl carbon.

It is often stated that one can conclude from these pH activity studies that it is only the undissociated molecule which is toxic i.e. action within the cell is visualized as caused solely by some combination of the neutral molecule of the acid or base with a sensitive element. For this conclusion to hold the biologically effective dose in terms of neutral molecules must be invariant with pH. Simon and Blackman (383) review several studies on echinoderm eggs in which this relation prevails for both weak acids and weak bases.

However studies on a wide variety of compounds—phenols cresols fluoride azide cyanide etc—show quite conclusively that the ion must also play a role in toxicity within the sensitive volume of the cell (382 383). Figure 6 exemplifies the data: it is seen that as the ionization increases with increasing pH the equi-effective concentration of undissociated molecules actually declines. The total drug—ions plus neutral molecules—rises of course but it is the fall in undissociated toxicant which is decisive.

meability occur a compound increasing the loss of cellular nitrogen and phosphorus but not increasing the entrance of substrates (189)

Finally we may expect relatively minor effects of sorbed materials especially on permeability and accumulation. We have referred earlier (Chapter 8) to the possibility that amino acid toxicity is exerted by the exclusion of one acid by another. Among other metabolic effects of β propiolactone on *Blastomyces dermatitidis* is the inhibition of amino acid uptake (119) although this could be secondary.


Reduction of the rate of solute movement is shown especially well by the studies of Rothstein (356) on the poisoning of yeast respiration by uranium. This appears to be a specific interference with the uptake of hexoses, endogenous respiration and the utilization of pyruvic acid or ethanol are very little affected. Rothstein suggests that uranyl ion complexes with groups at or near the cell surface which are essential to sugar uptake and suggests further that these groups are cellular polyphosphates which are also capable of binding other ions (357). Similar responses to uranium have been found in animal material (315).

5 HYDROGEN ION CONCENTRATION

In any survey of the literature on fungicides one is forcibly struck by the common failure to determine or to report the pH of the test solution. For practical purposes this may not be important at least if the pH is that of the projected use situation but conclusions on mode of action of ionizable materials are difficult indeed to draw if the acidity is unknown.


Some effects of pH are obvious: a fungicide may break down more or less rapidly and in so doing yield products of greater or less toxicity; thus thiosulfate forms a more toxic product (162) and dithiocarbamates form less toxic materials (432). A second possibility is that the toxicant itself changes the pH to a value unfavorable for growth or spore germination; this can of course be compensated for by specific controls. The chelating agents as mentioned later become less effective in binding metals as the pH is lowered: uranyl ion forms complexes with hydroxyl ion at pH values above 4.5 and becomes therefore unavailable (356). Competition between hydrogen ion and toxic cations is considered in connection with the heavy metals.

When weak acids or bases are dissolved in water the degree of ionization is pH dependent. This is evident by inspection of the equation for the dissociation of an acid



(2)

Figure 7 The general form of the relation between pH and the toxicity of a weak acid. Redrawn from Simon and Beevers (381) by permission of the American Association for the Advancement of Science



effect of structural changes on toxicity indicates that the acridines act at the surface and that they compete with other cations. Competition between toxicant ions and other cations is considered elsewhere in connection with the action of copper. It should only be mentioned here that this competition insofar as hydrogen ion is involved represents still another effect of pH on fungitoxicity.

Finally as suggested by Burchfield and Storrs (50) the internal pH of the cell could affect fungicidal action by its influence upon the ionization of cellular compounds which react with the toxicant. Thus reaction of amino groups with 2,4-dichloro 6-(*o*-chloroamino)-s triazine probably occurs only if they are unionized but sulphhydryl groups react only if ionized.

6. STRUCTURE AND ACTIVITY

Structurally Non Specific Inhibitors Specific chemical structure as a determinant of toxicity may be approached by considering first the structurally non specific poisons the narcotics. These are compounds the effect of which is freely reversible e.g. hydrocarbons chlorinated hydrocarbons aliphatic alcohols and phenols. The data on these are with few exceptions derived from studies on animal systems.

Probably no compound is absolutely without structural specificity. The structurally non specific inhibitors are however characterized by a relatively low specificity in the sense that changes in structure cause no apparent change in the mode of action. Historically the concept of structural non specificity arose from consideration of the effect of homologous series in general toxicity on a molar basis rises with

Two conclusions flow from this analysis. First inhibition of the sensitive cell element must be effected by both neutral molecules and ions although they need not be equally effective. A chemical model for this behavior is analyzed by Briggs (45). Second the pH at the site of action must be affected by changes in the external pH in order for the equeffective concentration of the penetrating species to decline as found (these are long time experiments in which we can assume equilibrium conditions). As mentioned in Chapter 1 the effect of external pH upon cytoplasmic pH is not known with any certainty or precision.

The significance of pK_a can now be assessed (Figure 7). The rise in required concentration for an equal effect begins at about pK_a . At lower pH (higher for a base) there is much less effect of acidity (383). It follows that in theoretical or comparative studies on weak acids or weak bases the effect of ionization can be minimized by assaying activity at a pH at least two units below pK_a (for an acid) or above pK_a (for a base). There are exceptions to this rule the sulfonamide antibacterials being the outstanding and best known example (6, 382).

The acridine antibacterials exemplify a rather different relation of toxicity to dissociation and pH (4, 6) one which has so far not come to light among organic fungitoxic materials. In this series only the cation is biologically effective. This fact and observations on the




Figure 6 The effect of pH on the concentration of 2,4-dinitrophenol ($pK_a \approx 4.0$) required to effect a 50 per cent reduction in the growth rate on agar of *Trichoderma viride*. Curve 1 total concentration curve 2 concentration of undissociated molecules. Redrawn from Simon and Blackman (383) by permission of the Cambridge University Press.




Figure 8 Distribution functions and toxicity in the aliphatic primary alcohols. Curve 1 solubility (μM per liter) curve 2 equitoxic concentration (μM per liter) for *Eberthella typhosa* curve 3 concentration (μM per liter) reducing the surface tension of water to 50 dynes per centimeter curve 4 vapor pressure at 25 ($\text{mm} \times 10^4$) curve 5 partition coefficient ($\times 10^4$) between water and cottonseed oil Redrawn from Ferguson (104) by permission of the Cambridge University Press

critical distribution or site of action. Nor do studies of this type offer the hope of determining the actual site of the toxic action.

It is true, however, that at present the distribution function most susceptible of experimental attack is permeability. That is, changes in toxicity coincident upon changes in thermodynamic activity must be investigated first to see whether the answer lies in the greater rate of entrance of the less water soluble compounds into the cell. In the aliphatic alcohols, permeability is almost certainly not the answer; toxicity increases rapidly as the molecular weight increases, but all members of the series penetrate very rapidly, and differences in rate of entrance do not explain differences in toxicity (79).

Toxicity does not, of course, increase indefinitely in a homologous series. Above a certain chain length, the exact point depending on the series and on the test organism, toxicity falls off sharply. This cutoff is a reflection, probably, of insolubility of and micelle formation by the higher members of the series (18, 53, 417).

The non-specific poisons are often referred to as physical toxicants.

molecular weight as the series is ascended but at some point the trend is reversed and higher members of the series are less toxic. All compounds of the series however appear to act in the same way on cells.

Early explanations of this homologous series effect were centered on permeability of the cell to toxicants. It was found that toxicity is correlated in such a series with an oil water partition coefficient. Since the plasma membrane is in general more permeable to non polar compounds than to polar compounds (84) it was natural to hypothesize that the more lipid soluble compounds are more toxic simply because they enter the cell more rapidly.

The classical paper of Ferguson (104) corrects and expands this limited hypothesis by introducing chemical potential as the index. Essentially Ferguson's principle states that the biological effectiveness of non specific poisons is a function of proportional concentration at the site of action. At equilibrium any compound is at the same proportion of saturation in all phases of the system. Consequently toxicity is a function not of bulk concentration in the external phase but of relative concentration or chemical potential. It follows that a substance only sparsely soluble in water will be more active mole for mole than a more soluble related compound. Consider compound A soluble in water to the extent of 1.0 M and compound B soluble only to 0.1 M. Ideally if an 0.05 M solution of B (all phases 50 per cent saturated) exerts a given non specific effect compound A will exert the same effect only at 0.5 M i.e. at 50 per cent of saturation.

Concrete experimental results are not of course so neat. It is a matter of some doubt for example whether the chemical activity required for a given biological response is constant or rises slowly as a homologous series is ascended (18-46).

Figure 8 taken directly from Ferguson (104) illustrates the principle for the series of aliphatic primary alcohols. As the series is ascended all the properties—solubility, surface tension lowering, vapor pressure and oil water partition coefficient—which depend on chemical potential change simultaneously. So does toxicity.

It must be emphasized that these considerations do not give specific information as to the site of action. All the constants of Figure 8 are distribution functions. From this we argue that toxicity depends upon a distribution between heterogeneous phases. We can not argue for any one distribution e.g. that between external solution and protoplasm. The governing partition could for example be that between the medium and the cell surface or that between the non polar portions of a protein chain and the ambient intracellular medium. Acceptance of Ferguson's principle does not of itself commit us to any particular

Table 1 Homologous Series and Fungistatic Activity

Series	Organism(s)	Chain Length of Maximum Activity
Saturated aliphatic alcohols (99)	<i>Penicillium glaucum</i>	10-11
Saturated aliphatic alcohols (110)	<i>Lentinus lepideus</i>	10-11
N alkylpyridines (230)	Four fungi	14
N alkyl α bromoacetamides (221)	<i>Aspergillus niger</i>	6-10
Fatty acids (295)	<i>Ophiobolus mayabeanus</i>	10-12
Fatty acids (350)	<i>Phymatotrichum omnivorum</i>	11
Fatty acids (456)	Two fungi	11
Aliphatic amines (110)	<i>Lentinus lepideus</i>	10-12
Nicotinic acid esters (177)	Two fungi	6-8
2 Alkyl 1 hydroxyethylimidazolines (437)	Four fungi	15-17
Omega (2 naphthyloxy) n alkyl carboxylic acids (55)	<i>Aspergillus niger</i>	2-3

ably more specific in their action. But in both types—specific and non specific—the homologous series effect is evident. Gross activity therefore depends in part on chemical potential and again we are dealing with a distribution function of some kind. Whether this distribution is between cell and environment i.e. is a function of permeability or whether some more subtle partition is decisive cannot at present be determined.

In general compounds of this type with branched chains are less toxic than would be anticipated from their molecular weight alone. Evidence for this generalization is reviewed by Horsfall (174).

Data on bacteria illustrate a quasi specificity (202) the carbon chain length for optimum toxicity depends to a considerable extent on the test organism.

Competitive Inhibition At the other extreme from structurally non specific inhibitors lies a group of poisons of extreme specificity. These may be detected operationally by the fact that over some range of concentration they compete with a cell metabolite. The theory and known examples have been considered extensively in the literature (277 353 454 455). Of the chemotherapeutic agents the sulfonamides are the classical example (450). Some at least of the sulfonamides are toxic to certain fungi (393 449) and the available data (40 92 98 147-149 365 393 410 449) indicate that as in bacteria sulfanilamide competes with *p* aminobenzoic acid for an essential cell reaction. Sulfanilamide is however not strongly fungitoxic (203).

Competitive inhibition in fungi is also demonstrable with metabolite

apparently because such factors as solubility are of more predictive value for gross biological action than are specific reactive groupings. The dichotomy does not seem justified either on theoretical or experimental grounds. Separation in theory of physical and chemical activities of molecules is artificial: the covalent bond and, for example, van der Waals forces may both be involved in the reactivity of the same molecule. Experimentally it is found that such narcotics as urethane, chloral hydrate, and ethyl ether have specific measurable effects on some cell activities at concentrations which do not affect other activities (138, 193, 194, 207, 261, 285). It has been proposed (190, 261) that the ultimate action of structurally non-specific poisons is the denaturation of one or more proteins of the cell. As Danielli (79) points out, this is not in conflict with the more general theory that these poisons act by virtue of their concentration in or on a lipid phase or non-polar structure of the cell.

The distinguishing feature of the structurally non-specific poisons from a chemical viewpoint is that they must form weak bonds with receptors of the cell. This follows from the fact that inhibition is within limits easily reversible by dilution alone. This also explains why activity is associated with chemical potential—a relatively high proportional concentration is necessary if such weak bonds are to occupy receptor sites enough of the time for perceptible inhibition to occur.

Most fungicides are not narcotics; i.e., structural specificity is the rule. The importance of the narcotics to the problem of structure and activity in the fungitoxic materials is that they may explain the effect of modifications of a basic toxicant molecule. Specifically, and following Horsfall (174), we may theorize that for an effective toxicant two requirements must be met. First, there must be one (possibly more) reactive atomic configuration that can combine with and inactivate essential materials of the cell. Second, the total structure of the toxicant molecule must allow concentration at the site of action. This second property includes, but emphatically is not restricted to, entrance into the cell. If, for example, the union between an enzyme and the combining region of a toxicant molecule is ionic or partially ionic, the formation of hydrogen or other weak bonds between the rest of the inhibitor and areas on the enzyme protein adjacent to the point of union would tend to stabilize the enzyme-inhibitor complex. (6)

Homologous Series and Fungicidal Action A selection of recent data on the homologous series effect is embodied in Table 1. Some of these compounds, e.g., the alcohols, are essentially narcotics as described in the previous section. The imidazolines, on the other hand, are prob-

endogenous respiration increased both processes generate an increase in oxygen uptake (52 67 145 331 446) In view of the generally non specific action of the phenols as a group (29) nitrophenols probably exert still other effects on cells

The reagents which inactivate sulfhydryl enzymes form a second category in which the specificity is toward the group of enzymes rather than toward any one although differences in sensitivity are common Free sulfhydryl groups on an enzyme or other protein can be inactivated by any of several reactions (21 94 95 306 316 455)

1 Alkylation by e.g. iodoacetic acid The product may be visualized as a thioether structure on the enzyme surface

2 Oxidation of two nearby sulfhydryl groups to a disulfide Ferri cyanide iodosobenzoic acid and presumably almost any mild oxidizing agent can effect this type of inactivation Thiols protect against this reaction

3 Mercaptide formation The characteristic agents of this reaction are heavy metals and trivalent arsenicals one of the most specific poisons of this type is *p* chloromercuribenzoic acid (316) Thiols protect also against these agents

4 Addition reactions These have been postulated particularly to explain enzyme inactivation by quinones and other $\alpha\beta$ unsaturated compounds

Although none of these four types of reaction is certainly known to be responsible for the action of fungitoxics it will be seen later that mercaptide formation can be invoked as a possible mechanism for the action of heavy metals against fungi and that the fungistatic quinones may act either by oxidation or by addition reactions to sulfhydryl groups

The third group is the most amorphous of all—compounds which are so generally reactive that they may be expected to inactivate a variety of enzymes possibly by more than one reaction As mentioned later the quinones may belong to this group rather than to the class of toxicants affecting sulfhydryl enzymes only The action of 2,4 dichloro 6 (*o* chloroamino) *s* triazine and its relatives (447) is presumably of this type Amino acids and peptides are alkylated by an exchange reaction for which the chlorine atoms attached to the heterocyclic ring are responsible and both sulfhydryl and amino groups are alkylated (50)

As discussed in the section on selective accumulation many fungicides are active only after a large amount of the toxicant is bound by the cell This suggests as McCallan (246) points out that most fungistatic

analogues of vitamins (Chapter 10) and amino acids (Chapter 8) Horsfall (174) lists in convenient form a number of examples

A few of the more active fungicides are conceivably metabolite analogues competing with normal substrates or coenzymes but in no instance has the action of such a fungicide been explained on this basis. Inhibition by 2,3-dichloro-1,4-naphthoquinone is reversed by vitamin K (453) but there is no evidence that the vitamin is essential to fungi. Simple reversal experiments, inconclusive in themselves, have been the bases of suggestions that 2-heptadecyl-2-imidazoline interferes competitively with purine biosynthesis (438) that ascocin is a fatty acid anti-metabolite (154) and that propionic acid interferes with acetate metabolism (155, 156, 326).

Inhibition of Enzymes Present-day biology relies in perhaps too great measure on the findings of enzymology. Toxicants to which no specific role can be assigned are often stated to affect some unspecified, enzyme or enzymes: such a statement is more an article of faith than an hypothesis.

However, there is no doubt that toxicants do affect enzymes, and we may distinguish three classes in relation to fungi: (1) agents which act specifically by inhibition of one essential enzyme system; (2) agents which act on sulphydryl enzymes as a group; and (3) agents whose demonstrable reactivity with proteins suggests that they act on enzymes in general. This classification may be considered either as based on decreasing specificity of the fungicide or on our increasing ignorance.

The first group—compounds which owe their toxicity to inhibition of a single enzyme system—is exemplified by the antimycins, antibiotics produced by *Streptomyces* sp. (96, 231). These interfere specifically with electron transport in the cytochrome system (63, 335, 343, 421). Antimycin may affect other electron transport systems also (305). Obviously, the specifically antifungal activity of the antimycins cannot be explained on this basis alone, since most organisms depend on the cytochrome system.

The action of the so-called uncoupling agents—2,4-dinitrophenol and 3,5-dinitro-*o*-cresol—are examples—cannot be specified so narrowly as that of the antimycins. It appears that they prevent the oxidative phosphorylations which are associated with the transfer of electrons from substrate to oxygen (233, 379, 380, 412). The effect is to liberate respiration from the control normally imposed upon it by phosphate storage compounds of the cell, and at the same time inhibit reactions which are driven by the hydrolysis of phosphorylated intermediates. Operationally, it is found that oxidative assimilation is suppressed or

endogenous respiration increased both processes generate an increase in oxygen uptake (52 67 145 331 446) In view of the generally non specific action of the phenols as a group (29) nitrophenols probably exert still other effects on cells

The reagents which inactivate sulphhydryl enzymes form a second category in which the specificity is toward the group of enzymes rather than toward any one although differences in sensitivity are common Free sulphhydryl groups on an enzyme or other protein can be inactivated by any of several reactions (21 94 95 306 316 455)

1 Alkylation by e.g. iodoacetic acid The product may be visualized as a *thioether structure on the enzyme surface*

2 Oxidation of two nearby sulphhydryl groups to a disulfide Ferricyanide iodosobenzoic acid and presumably almost any mild oxidizing agent can effect this type of inactivation Thiols protect against this reaction

3 Mercaptide formation The characteristic agents of this reaction are heavy metals and trivalent arsenicals one of the most specific poisons of this type is *p*-chloromercuribenzoic acid (316) Thiols protect also against these agents

4 Addition reactions These have been postulated particularly to explain enzyme inactivation by quinones and other $\alpha\beta$ unsaturated compounds

Although none of these four types of reaction is certainly known to be responsible for the action of fungitoxics it will be seen later that mercaptide formation can be invoked as a possible mechanism for the action of heavy metals against fungi and that the fungistatic quinones may act either by oxidation or by addition reactions to sulphhydryl groups

The third group is the most amorphous of all—compounds which are so generally reactive that they may be expected to inactivate a variety of enzymes possibly by more than one reaction As mentioned later the quinones may belong to this group rather than to the class of toxicants affecting sulphhydryl enzymes only The action of 2,4-dichloro-6-(*o*-chloroanilino)-s triazine and its relatives (447) is presumably of this type Amino acids and peptides are alkylated by an exchange reaction for which the chlorine atoms attached to the heterocyclic ring are responsible and both sulphhydryl and amino groups are alkylated (50)

As discussed in the section on selective accumulation many fungicides are active only after a large amount of the toxicant is bound by the cell This suggests as McCallan (246) points out that most fungistatic


agents now in use are general poisons rather than inhibitors of specific enzymes. The same conclusion arises from the finding (319) that several very different enzymes are inhibited by fungicides and is supported by the usual failure to obtain mutants resistant to organic fungicides (217).

Structural Modification and Fungitoxicity. The possibility of obtaining new fungicides or more useful modifications of older materials has motivated considerable research on the relation between structure and activity. Unfortunately much of this work is of little help in determining mode of action since such essential factors as dissociation of the toxicant, permeability of the cell, and the extent of accumulation are usually unknown.

Horsfall (174) and Sexton (372) review the general field of structure and activity, some especially important recent work has been published on the s triazines (51-367) derivatives of acetanilide (66) triphenyl methanes and other dyes (93) substituted pyrazoles (267) unsaturated hydrocarbons (262) ethylene oxide derivatives (329) and nitrophenols (374).

The phenols have been the most extensively investigated from this aspect. Phenol itself and many of its derivatives have been regarded as structurally non specific, i.e. as narcotics (29) but activity toward fungi and bacteria is markedly influenced by the number and type of substituents (372-408). As a rule alkylation increases activity and within limits the longer the alkyl side chain the greater the toxicity (29-141-377). This follows the rule for homologous series i.e. there is an optimum chain length for activity against a given organism (372). Halogenation also increases the toxicity of phenols (29-282-376) but the position of the halogen is important (29-375-408). The nitrophenols have already been mentioned as respiratory inhibitors.

The bisphenols—not necessarily resembling the phenols in their action—provide another example of the effect of structure on activity. In general both antifungal and antibacterial activities require that the compound be halogenated, that the linkage between the two rings be *ortho* to hydroxyl groups and that the two rings be not too far apart (58-274). Hexachlorophene or 2,2-methylenebis(3,4,6-trichlorophenol) illustrates these features.



Replacement of the methylene bridging group by sulfur yields active materials (328) and still other linkages are compatible with activity (73 295) Bisphenols have proved especially effective in the protection of textiles from fungal decay (386) and against fungi pathogenic to animals (58) The mode of action is not known hexachlorophene is inhibitory at high dilution to several respiratory enzymes (135 136)

In considering the effect of substitution on fungitoxicity data can not be evaluated properly on the basis solely of the usual two-dimensional visualization of an organic molecule Some of the often neglected factors may be listed very briefly

- 1 A substituent may alter the reactivity of nearby groups or of as large a component as an entire ring Substituents may be electron attracting or electron repelling may encourage or discourage hydrogen bonding and chelation

- 2 A change in structure may either reduce or facilitate stabilization by resonance

- 3 The addition of substituents may change the pK This is especially marked in the phenols (29) phenol itself has a pK of 10.0 p nitrophenol of 7.2 2,4-dinitrophenol of 4.0 (383) Thus at pH 7 phenol is less than 1 per cent ionized and 2,4-dinitrophenol is over 99 per cent ionized

- 4 The oxidation reduction potential of for example the naphthoquinones may be either raised or lowered by substituent groups

- 5 The shape of the molecule considered three dimensionally may be altered by substitution this has been especially thoroughly explored in the acridines the antibacterial action of which requires a minimum planar area (6)

- 6 Alterations in a molecule may make it more or less susceptible to biochemical attack by enzymes of the test organism as suggested by Owens (319) for substituted phenols

- 7 Lastly as previously discussed changes in the physical properties of a compound may render it more or less able to enter the cell or to accumulate at its site of action

7 THE ACTION OF HEAVY METALS

The major share of our information on the fungistatic activity of the heavy metals comes from studies on copper the active ingredient at least until the advent of organic fungicides of some of the most important plant and textile protectants Mercury compounds have been used less for fungi but more as bacteriostatic agents Silver is of little practical importance because of its cost although it is an

effective toxicant (256 311 312) Very little information is available on heavy metals other than these three chromate complexes have been studied recently (146 439)

Horsfall (174) summarizes the action of metallic cations measured by external concentration against fungi in the order $\text{Ag} > \text{Hg} > \text{Cu} > \text{Cd} > \text{Cr} > \text{Ni} > \text{Pb} > \text{Co} > \text{Zn} > \text{Fe} > \text{Ca}$ For any one organism the order may differ slightly (23, 256 300) Roughly the same order of effectiveness is found in experiments on animals (373)

Because of the reactivity of the heavy metals growth experiments in complete media are of dubious value the spore germination technique is to be preferred Proteins and peptones are especially active in decreasing fungitoxicity (153 214 458) but other materials present in culture media have the same effect (247)

Present information allows us to consider the three most toxic heavy metals—silver mercury and copper—together i.e. to assume provisionally that all have the same general mode of action silver may however differ We can also but perhaps with less justification consider that the final toxic effect of heavy metal compounds is to be ascribed to the metal itself However, some problems are raised by this second assumption Different inorganic salts of copper for example are not always of equal toxicity (28 279) when copper is supplied as cupric ammonium sulfate it is more firmly bound by spores than copper supplied as the sulfate (34) These differences cannot be explained as yet in detail but we may guess that the form in which copper is supplied affects its uptake by the spore Provision of silver as the iodide sharply reduces both uptake and toxicity as compared to those of other silver halides (288)

The toxicity of coordination complexes of copper deserves special attention For years the effectiveness of Bordeaux mixture as a plant protectant was a mystery inasmuch as the copper is present on the leaf in an insoluble complex A number of investigations reviewed by Martin (278) and by McCallan (244) established firmly the proposition that materials from the spore solubilize the copper The most important of these substances are the amino and hydroxy acids compounds which form soluble chelate complexes with copper Mono methylamine liberation by spores of *Tilletia caries* also results in solubilization of copper (327) and seeds of higher plants liberate copper from Bordeaux mixture (429)

This hypothesis requires that soluble copper complexes be toxic. Some data show that under specified conditions the glycine and malate action may not be so great (130) Concentration factors are critical complexes are as fungistatic as copper sulfate itself (257) fungicidal

if the ratio of complexing agent to metal is high—five to ten—toxicity disappears (28). This is consistent with the concept that cell and complex compete for copper: it is significant that ethylenediaminetetraacetate (EDTA) which forms very stable metal complexes abolishes toxicity at a 1:1 molar ratio to copper (28, 302). Unfortunately these same observations are consistent also with a second explanation: that the malate and glycine complexes enter the cell whereas the EDTA complex does not.

Organic mercurials are generally more toxic than inorganic, both to bacteria (39) and to fungi (174, 372). Part of the answer to their practical effectiveness lies in the appreciable volatility of some of them (17, 88, 220). There is some evidence, however, that the effective organic mercurials penetrate the cell as such, i.e. that they are more readily taken up than mercuric ions (35, 324). Against this may be cited limited evidence from pH studies in a complete growth medium (187) that phenylmercuric acetate is more toxic as the ion. In general both inorganic and organic mercurials affect the same enzymes and are equally antagonized by thiols, although occasional secondary effects complicate the picture as in the case of pancreatic amylase (319).

The extreme case of increase in toxicity by provision of a metal in an organic molecule is found in a study of the fungitoxicity of tin compounds (199). Stannous and stannic chlorides are essentially non-toxic but tri-*n*-butyltin acetate prevents the growth of fungi at 0.1–0.5 ppm.

McCallan and Wellman (267) find that copper, silver and mercury are both fungistatic and fungicidal. Fungicidal action is of course a function of time of exposure and of the method by which the spores are washed free of toxicant.

The action of heavy metals on fungi may be conveniently divided into two topics: the uptake of metals by spores and the metabolic or other effects of the metal once it has reached a sensitive site.

We have already mentioned that metals, in common with other fungistatic materials, are accumulated by fungus spores in very high concentration and very rapidly (288, 290, 291). Some further light is cast on this by earlier and technically less satisfactory studies on the effect of copper and mercury on *Tilletia tritici* (34, 35, 36). Copper uptake was found to follow the Freundlich adsorption isotherm, i.e. to be non-linearly related to external copper concentration. Copper supplied as an ionizable salt is removable by acid, but the copper of cupric ammonium sulfate, a coordination complex, is not so removed. Finally, ionic copper disappearing from solution is almost quantitatively accounted for by other cations— H^+ , Fe^{++} , Mg^{++} , etc.—liberated from the spore.




Figure 9 The effect of pH on the toxicity of copper chloride to spores of *Alternaria tenuis*. Curve 1 (solid line) control curve 2 (broken line) 0.0001 M CuCl₂. Drawn from data of Biedermann and Muller (28)

Divalent cations generally reduce both uptake and toxicity of the heavy metals (1 28 150 226 232 273 308). The uptake of silver by spores is reduced by copper and more completely by mercury (288). Competition by hydrogen ion is equally important; this is shown in Figure 9 and also by the fact that absorbed copper is removed from spores by acid (3 34 174 291).

There are at present two ways in which to interpret the data on uptake of metals by spores. First we may assume that the entire volume of the spore protoplasm accumulates the metal. This assumption requires, in view of the rapidity of the uptake, that the metal move relatively freely across the semi permeable membrane; such a rapid movement is suggested by the speed with which cellular zinc of *Aspergillus niger* exchanges with extracellular zinc (288) and by some data on the loss of potassium from bacteria (209). Staining methods show that copper does enter spores (129 418) but this approach has not been developed sufficiently to be of critical importance.

The second mode of interpretation is to consider metal uptake as an adsorption defined as a process that concentrates one component of a system at an interface and to limit the locus of this adsorption to peripheral regions of the cell, i.e. outside the permeability barrier.

This approach has the advantage that we need not postulate that ions are able to penetrate the cell rapidly although as just mentioned there is some evidence for rapid passage of metals through the membrane it is generally true considering all the organisms studied that cations especially di and trivalent cations do not move easily into cells by diffusion (84) The concept that the initial uptake is an adsorption or ion exchange process has the additional advantage of explaining both ion competition effects and the fact that uptake of copper is accompanied by roughly equal extrusion of other ions Nor does a surface adsorption require that the cell be living or even metabolizing normally it has been noted (291) that uptake of several materials continues long after enough toxicant has been absorbed completely to prevent spore germination

Against an adsorption hypothesis it may be urged that the uptake of some materials is so large that the classical monomolecular film could not accommodate the observed toxicant (290)

To test these two hypotheses the most direct approach is provided by model experiments on the uptake of methylene blue by bacteria (238-239) and by ascospores of *Neurospora tetrasperma* (407) The uptake reaction is independent of temperature a metabolically powered accumulation would have a high temperature coefficient Dead cells adsorb dye as rapidly and completely as living cells Different cations compete with each other for a limited number of sites as in yeast (356-357) it appears that the cell surface has a countable number of more or less strongly negative binding sites Exploration of the uptake-concentration relation does not appear so profitable any type of accumulation would show saturation effects as the external concentration is raised


If it proves possible to regard the initial rapid uptake of metals as a surface adsorption then the toxic action of heavy metals can be visualized as already mentioned as composed of a rapid stage of uptake at the surface followed by a slower penetration perhaps through a damaged membrane into the interior of the cell Alternatively the metal may enter as a coordination complex.

Effects of sorbed metals upon the cell surface must be considered as a possibility although it seems doubtful that the major site of toxic action is outside the permeability barrier Copper reduces the permeability of the erythrocyte to glycerol (84) and heavy metals especially mercury depress phosphate uptake by bacteria (294) Silver increases the loss of phosphorus from fungus spores although mercury does not (288) Interference with catabolic anabolic or carrier mechanisms outside of or on the plasma membrane is also a possibility as in the action

of uranyl ion on yeast (p 151) Increases in respiratory rate are induced by sublethal concentrations of metals (247) the explanation may lie in an increased permeability to exogenous substrate or less likely in inactivation of regulatory enzyme systems (20)

Present theories of the ultimate action of heavy metals visualize them as enzyme poisons and admit only the question of whether they are primarily effective against sulphydryl enzymes or generally inhibitory to enzymes of all types

The reaction of a divalent metal with a sulphydryl enzyme is depicted conventionally as mercaptide formation



The resemblance to a five membered chelate complex—there are two points of attachment on the enzyme—is obvious. A monovalent metal or an organic mercurial like *p*-chloromercuribenzoic acid reacting as such presumably forms a mercaptide with only a single sulphydryl group. The argument for the importance of this reaction is based primarily on the observation that heavy metal poisoning of both bacteria and fungi can be prevented or within certain time limits reversed by such thiol compounds as cysteine glutathione and 2,3 dimercaptopropanol (107 185). In addition it can be shown that mercaptide forming reagents react stoichiometrically with free sulphydryl groups of proteins (13 316)

Shaw (373) proposes a model for the inhibition of vital processes that is based on these considerations and the added assumption that the sulphydryl groups with which cations combine are part of the catalytically active site of the inhibited enzyme. The driving force of Equation 3 is regarded as proportional to the free energy of the simple reaction



That is toxicity should be correlated with the insolubility of the metal sulfides (MeS). The order of insolubility is in fact very much like the order of toxicity $\text{Hg} \approx \text{Ag} > \text{Cu} > \text{Pb} > \text{Cd} > \text{Zn}$. This is also the order of affinity of metals for the sulphydryl group of purified serum albumin (201)

Emphasis on reaction with sulphydryl groups may however divert attention from other possibilities. In particular the simple demonstration that thiol compounds antagonize the toxicity of metals to in

tact cells cannot by itself prove that the primary effect of metals *in vivo* is on sulphhydryl enzymes. Thiols form stable covalent compounds with metals and would be expected therefore to compete successfully with almost any bond formed between enzyme and metal.

That the sulphhydryl hypothesis may be too limited is indicated above all by the observation that many enzymes which are not apparently dependent upon sulphhydryl groups for their activity are inhibited *in vitro* by one or more of the heavy metals. These include sucrase, β glucosidase, arginase, asparaginase, carbonic anhydrase, catalase, and cytochrome oxidase (319, 388, 405). As already mentioned, inhibition of respiration and spore germination at approximately the same concentration by heavy metals is also suggestive of a general poisoning of many different enzymes.

If we grant that heavy metals are general enzyme poisons, we may inquire about the sites at which they might act on an enzyme. Obviously, as Horsfall (174) emphasizes, the order of toxicity closely resembles the order of stability of metal chelate complexes. From this it is apparent that the heavy metals might act by displacing from an enzyme a metal like magnesium which is essential to its activity and which is bound normally in a coordination complex. However, the order of stability of chelate complexes is, for obvious reasons, much the same as the order of insolubility of metal sulfides just described, and either series may with equal probability be considered the more important.

Several potential binding sites are available on proteins to react with transition metals like mercury, silver, and copper. These include carboxyl, phosphoric, imidazolium, ammonium, and phenolic hydroxyl groups (204). The primary carboxyls and the α amino groups of the amino acids are, of course, tied up in peptide bonds, and such side chains as the guanidine and amide groups cannot be expected to bind metals at biological pH values.

In principle, at least, inactivation of enzymes by metals is not limited as to location; the enzyme or enzymes may just as well be at the surface of the cell as in the interior. However, it seems more probable that the metals reach the interior of the cell and are not confined in their action at fungistatic concentrations to the surface.

This discussion of toxicity of the heavy metals makes it clear that there are numerous unsolved problems, especially with regard to the penetration of heavy metal ions into the cell. Some of the more important points may be listed in conclusion.

1. Differences in toxicity between various compounds of a given metal are ascribed to differences in the rate at which the toxic metal

reaches the site of action. The contrast between ionized salts and unionized lipid soluble organic compounds is especially marked.

2 Toxicity of the metals is affected by hydrogen ion and divalent cations and by the presence of natural substances—proteins amino acids hydroxy acids etc.—which bind metal ions.

3 Metals are accumulated by spores in large amounts from dilute solutions.

4 If metal ions cross a permeability barrier slowly it is possible that accumulation occurs by an adsorption process and is followed by a slower entrance of the ion or a complex of it into the protoplast. Ion antagonism according to this hypothesis is exerted in the adsorption phase. For the present however it must be admitted that this mechanism is speculative.

5 Although the primary effect of metals is probably on enzymes in the interior of the cell effects on the permeability barrier itself cannot be excluded entirely.

6 Heavy metals inactivate enzymes. Combination with essential sulfhydryl groups is the favored explanation but it is likely that other inactivation reactions involving the formation of electrostatic or coordinate bonds also contribute to enzyme inhibition.

8 THE ACTION OF SULFUR

The most important organic sulfur fungicides are the dithiocarbamates considered later as chelating agents. Many other organic sulfur compounds—dithiazolidines (12) thiourea (80 112) mercaptobenzo thiazole (100) xanthates (82) arylthioalkancarboxylic acids (102) phenothiazone (131) 3 pyridinethiol (394) benzothiazoles (142) compounds with the *N*-trichloromethylthio grouping (201) substituted thio cyanates (215 144) thio bisphenols (328) and such antibiotics as acti thiazic acid and gliotoxin—are more or less fungitoxic. However there is no reason to suppose that sulfur is necessarily directly involved in the reaction of all these diverse compounds with cellular constituents. In some organic molecules the presence or absence of sulfur has no influence on toxicity (57 82) in others it has (102). Some of the sulfur compounds probably act primarily as chelating agents (11 174). In others sulfur may exert an effect by modifying the reactivity of nearby functional groups or by affecting solubility.

Inorganic sulfur like copper has been in use as a fungicide since the beginning of serious efforts to control plant disease. The various forms used including liquid fine sulfur all act ultimately as elemental sulfur. Sulfur is of course virtually water insoluble and a great deal of effort

was earlier devoted to proving that some more soluble derivative is the active agent the history of these investigations has been reviewed elsewhere (174-244). It is now agreed that the form in which sulfur reaches the fungus spore is the vapor phase of elemental sulfur (246). Fungitoxicity in any particular situation of use is dependent upon temperature (459), the particle size of the sulfur deposit (370-442) and to some degree the allotropic form of the sulfur (103).

For some time following a series of critical studies the details of which are reviewed by McCallan (244-246) and Horsfall (174) elemental sulfur was considered to be reduced by the fungus spore and the effective toxic agent was taken to be the hydrogen sulfide formed by the biological reduction. Hydrogen sulfide is fungistatic (442-443) and somewhat fungicidal (248-249). It appears now however that the toxicity of elemental sulfur on a weight basis is greater than that of an equivalent amount of hydrogen sulfide as shown in Table 2. Conse-

Table 2 The Relative Toxicities of Sulfur and Hydrogen Sulfide*†

Species	Wettable Sulfur	Colloidal Sulfur	Hydrogen Sulfide
<i>Monilinia fructicola</i>	54	0.5	2.8‡
<i>Cephalosporium acremonium</i>	>1000	0.3	12‡
<i>Aspergillus niger</i>	>1000	0.3	15‡
<i>Glomerella cingulata</i>	>1000	0.4	20‡
<i>Neurospora sitophila</i>	>1000	1.0	38‡
<i>Rhizopus nigricans</i>	>1000	2.7	5.9
<i>Alternaria olivacea</i>	>1000	18	15
<i>Stemphylium sarcinaeforme</i>	>1000	31	8.8

* From Muller, McCallan, and Weed (789) by permission of the Boyce Thompson Institute for Plant Research, Inc.

† Toxicity is expressed as the dose in parts per million required to kill 50 per cent of spores in an exposure of 24 hours; concentrations are of the external solution or suspension.

‡ Highly significant difference between colloidal sulfur and hydrogen sulfide.

quently the second part of the older theory—that hydrogen sulfide is the ultimate toxicant—is untenable.

Attention has therefore been drawn to the process of sulfur reduction itself as a possible primary event in toxicity. The reduction of sulfur to hydrogen sulfide by fungus spores is a general phenomenon and has been shown to be brought about by spores of many different fungi under both anaerobic and partially aerobic conditions (253-289). The activity of species varies widely. *Cephalosporium acremonium* under anaerobic conditions forms 6.39 mg hydrogen sulfide per

gram of spores per hour almost 300 times the rate of the same process in *Rhizopus nigricans*. In these experiments there was no external hydrogen donor so that cell constituents must have been oxidized; it is found as would be expected that under such circumstances high rates of hydrogen sulfide evolution are not maintained for long. In aerobically the formation of hydrogen sulfide from sulfur is accompanied by an equivalent release of carbon dioxide; aerobically sulfur increases oxygen uptake (246b).

Mycelium of *Fusarium lini* is similarly capable of reducing elementary sulfur in the presence of isopropyl alcohol as hydrogen donor (369).

Fungi of different species differ very markedly in their sensitivity to sulfur under field or laboratory conditions. Of the common test fungi for example *Stemphylium sarcinaeforme* is about 60 times as resistant as *Monilia fructicola* (Table 2). However it has not yet been possible to correlate the enzymatic reduction of sulfur with its fungitoxicity—fungi which are relatively tolerant of sulfur are numbered among both those most active and those least active in reducing sulfur, whereas the more sulfur sensitive species tend to be intermediate in reducing capacity (289). However this type of study has only begun and it is not yet possible to exclude any hypothesis with finality.

The production of large amounts of hydrogen sulfide from elemental sulfur in the absence of external hydrogen donors suggests that some relation may in the future be established between the derangement of metabolism associated with this process and the toxicity of sulfur (259). It is clear that sulfur is competing with normal hydrogen acceptors and two speculative hypotheses are suggested by this conclusion. The first may be expressed rather crudely by the proposition that in the presence of very large amounts of a suitable hydrogen acceptor sulfur dehydrogenation reactions escape from the normal control imposed upon them under anaerobic conditions. This normal control may be either the limited supply of acceptors for high energy phosphate or a poisoning of metabolism by the accumulation of such fermentation products as ethanol; the first of these is the more likely. The physiological effect of such an escape from regulation would be the disappearance of the cell's reserves. According to this hypothesis sensitivity would be a function of the amount of utilizable reserve materials available to the spore and should be reduced by provision of an external hydrogen donor.

A related hypothesis to explain a deleterious effect of the reduction process itself is that utilization of reserves per se is less important than the fact that provision of sulfur results in a series of oxidations which are not coupled to the energy storage and energy utilization mechanisms.

of the cell. That is, sulfur diverts metabolism into non productive or inefficient pathways.

Anaerobically methylene blue prevents the reduction of sulfur by spores of *Neurospora sitophila* (289). It is known that methylene blue does not accept hydrogen directly from substrates or the pyridine nucleotide coenzymes but from hydrogen (electron) carriers such as the flavoproteins and possibly cytochrome b (116). If the sulfur reduction process is in itself damaging to the cell, it is at this level that we should look first. It may be noted that sulfur reduction is also inhibited by 2,3-dichloro-1,4-naphthoquinone, the oxidation-reduction potential of which is presumably close to that of methylene blue.

Selenium and tellurium are much less toxic than sulfur and their reduction to hydrides by yeast cells is correspondingly much less rapid (443). Selenium is accumulated in organic form by the mycelium of *Aspergillus niger* in competition with sulfur and selenate inhibition is partially or wholly counteracted by reduced sulfur compounds (435). These and similar data on *Chlorella vulgaris* (378) can be interpreted in terms of competitive inhibition by a metabolite analogue.

9 QUINONES AND OTHER UNSATURATED COMPOUNDS

A number of actively fungistatic materials are characterized chemically by a system of conjugated double bonds. Of these the most important are the quinonoid compounds but the α,β -unsaturated ketones are also of interest.

The most generally active quinonoid compounds are the halogenated *p*-benzoquinones (266-411) and 1,4-naphthoquinone and various derivatives of it (228-229, 280, 364-413) including synthetic vitamin K (2-methyl-1,4-naphthoquinone) (15, 70, 133). The triphenylmethane dyes, e.g., malachite green and crystal violet, are fungistatic and significantly the leuco form (benzenoid) is not active against either fungi (93) or bacteria (111).

The chromophoric group of the actinomycins is quinonoid (Chapter 8) and the antibacterial javanicin is a naphthoquinone (Chapter 6). The tropolones, some of which are produced by fungi (Chapter 6), have a modified quinonoid structure (325). Some of the thujaplicins present in the decay-resistant heartwood of *Thuja plicata* show moderate fungistatic activity (19, 314-345).

A few examples of quinonoid fungicides are shown in Figure 10.

It will be recalled that many fungi possess part or all of an enzymatic system which oxidizes phenols to quinones and further metabolizes the quinones to black melanin pigments (Chapter 6). It follows therefore




Figure 10 Fungistatic quinonoid compounds

that appropriate hydroquinones themselves non-toxic may be converted to toxic quinones. Thus 2,3-dichloro-1,4-naphthohydroquinone inhibits *Monilia laxa* presumably because it is oxidized by the fungus to the corresponding naphthoquinone; esters of the hydroquinone are first hydrolyzed then oxidized (56).

Detoxification of quinones would be expected of a fungus which forms melanin pigments. *Stemphylium sarcinaeforme* which is normally black pigmented appears to convert many quinones to non-toxic products (347).

The reactivity of quinones is so great that more than one mode of action may easily be envisaged. Although any quinone affects the oxidation-reduction potential it does not seem possible to correlate either bacteriostatic or fungistatic effects with this property (266, 321, 434). As α,β -unsaturated ketones, quinones can react with both sulphydryl and amino groups (372) and these possibilities have enlisted considerable interest.

Quinones react by addition with simple thiols (212, 292). Correspondingly the antimicrobial activity of quinones is often reversible by sulphydryl compounds (70, 121). However we have already noted that thiol reversal is not by itself sufficient proof of the nature of the *in vivo* reaction. Many of the enzymes known to depend upon sulphy

dryl groups for their activity or known to be inhibited by sulphhydryl reagents are it is true inhibited *in vitro* by quinones and naphthoquinones (113 159 211 213 228 319 320 364) and protected from this inhibition by thiols (319). However a series of investigations shows first that quinones inhibit enzymes other than sulphhydryl-dependent types and second that the order of effectiveness of different quinones as enzyme inhibitors is not correlated with their antimicrobial activity (163-171). It is also to be noted that quinone toxicity to Gram positive bacteria is not antagonized by thiols (121).

These considerations indicate that some other reaction with essential proteins should be considered possibly an inactivation of amino groups. For the present it is not possible to specify any particular ultimate site of action for the quinones and the evidence suggests that they may be general rather than specific enzyme inhibitors comparable to the heavy metals.

The relation of structure to fungistatic activity in the quinones is reviewed by McNew and Burchfield (266). Toxicity generally decreases in the series naphthoquinone > phenanthraquinone > benzoquinone > anthraquinone. Halogenation improves effectiveness the effect of other substituents depends upon specific type and position. Comparison of the effect of structure on inhibition of amylases reveals several instances in which a given structural feature affects fungitoxicity in one way enzyme inhibition in another (320). Among the naphthoquinone antimalarials activity is generally correlated with low water solubility that is not only a reactive grouping but also some non polar properties are essential (105).

Although some quinones are natural products of fungi and could in principle participate in terminal respiration (Chapter 6) there is as yet no firm evidence that fungitoxic quinones act as metabolite analogues although as mentioned earlier their action may be reversed by natural quinones.

Non quinonoid $\alpha\beta$ unsaturated ketones are not so active as the quinones against fungi many however are antibacterial. Several e.g. patulin and penicillic acid appeared first as antibiotics produced by fungi. These compounds inhibit sulphhydryl enzymes and their inhibition of both isolated enzymes and intact cells is reversed by thiols. Consequently it is believed that antibacterial action depends upon some reaction perhaps addition across the reactive double bond with sulphhydryl groups (122 123 351). Patulin however does not inhibit urease a sulphhydryl enzyme (168) and the action of dehydroacetic acid is not reversed by thioglycolate (451).

Studies of the action of unsaturated compounds on *Botrytis allii* and

groups and hence extracts easily into a non polar solvent. The 1:1 chelate is polar. A polar group not involved in the union with the metal would of course remain unchanged and reduce lipid solubility. The number of different complexes formed by a given metal with a ligand is a function of its coordination number.

Biological systems necessarily contain many different ions able to participate in chelate complexes and the nature of competition must be considered. In general most chelating agents form complexes of varying stability with metal ions in the order (from most to least stable) Cu^{++} Ni^{++} Co^{++} Zn^{++} Cd^{++} Fe^{++} Mn^{++} Mg^{++} (276). Thus copper will displace any of the other listed ions, magnesium none. However some biologically important chelating substances do not follow this rough rule. $\alpha\alpha$ dipyridyl and *o* phenanthroline for example have a specific affinity for iron.

Stability of metal chelates is usually expressed by the constant K , the formation (or dissociation) constant, a measure of the equilibrium



in which M = metal, A = ligand and x = the number of ligand molecules per metal atom in the complex. Ideally K is expressed by

$$F = \frac{(\text{MA})}{(\text{M})(\text{A})} \quad (7)$$

in which brackets indicate activities (276). Values of $\log K$ may be very high, e.g. 23 for the copper oxine complex; it is obvious that negligible amounts of free copper can exist in the presence of excess oxine.

Factors which fundamentally determine K are discussed by Calvin (59). Chabarek et al. (63) describe the use, under biological conditions, of a plot of the negative logarithm of metal concentration against pH for prediction of the behavior of chelate systems. The ligand may be considered as a Lewis base; at low pH more of it is bound by protons and consequently the metal chelates are less stable under acid conditions.

Many normal cell metabolites are chelating agents; examples of such metabolites include amino and hydroxy acids, porphyrins, peptides and polyphosphates. Some of these may owe their biological function to their chelating properties. Chelation is almost certainly necessary to the function of the manganese requiring peptidases (289) and quite likely it is essential to the action of other metalloenzymes and electron or oxygen transport systems (59-97).

We may approach the question of fungistatic action by first noting that chelating agents are often powerful inhibitors of metalloenzymes.

in cell free preparations. The poisoning of certain iron enzymes e.g. cytochrome oxidase by cyanide or azide presumably involves chelate bonds with the iron atom. Enzymes known to depend on metals for their activity are inhibited by chelating agents: polyphenoloxidases by diethyldithiocarbamate azide and other compounds able to complex with copper (184, 319, 405); zinc metalloenzymes by oxine, $\alpha\alpha$ -dipyridyl and *o*-phenanthroline (420). The specificity of chelating agents is however so low at biological pH levels that inhibition by them does not as a rule suffice to identify the metal active for a given enzyme or enzymatic system.

In the systems just mentioned it is not known whether the chelating agent combines with the metal *in situ* on an enzyme or coenzyme or competes with the catalyst for the limited amount of metal present. Inhibition of dialkylfluorophosphatase by 8-hydroxyquinoline and ethylenediaminetetraacetic acid (EDTA) appears to arise from removal of calcium from the enzyme (299); other enzymes dependent upon calcium for activity or stability are also inhibited by EDTA (86, 139).

Finally a few examples of *in vivo* effects of chelating agents may be mentioned: inhibition of synthesis of phenolic compounds by *Phycomyces blakesleeanus* (48); prevention of the transformation of *Brucella abortus* from the smooth to the non smooth form (68); and inhibition of glutamic acid assimilation by *Staphylococcus aureus* (117). In the two last mentioned instances it is believed that the biologically active metal is manganese but the locus of the manganese chelate—within the cell or outside it—cannot be determined. Several metal complexing agents including cyanide, sodium diethyldithiocarbamate and ethylenediaminetetraacetic acid cause the release of subcellular particulates from cells of *Candida albicans* (284).

Although it is usually found that in any series of chelating agents many compounds are toxic (174, 180) and many empirically discovered toxicants turn out to be chelating agents (7, 60, 433) the association is not uniform presumably because both the stability of the complex and its molecular architecture are factors in toxicity or in the ability of the molecule to reach the site of toxicity. Chelation alone in other words does not necessarily confer toxicity (8, 261). The best example of a close association between toxicity and chelation is in the series of compounds related to oxine: only those that complex metals are bacteriostatic (10, 11).

The association of toxicity and ability to chelate was at first thought to signify only that the chelating agent deprives the cell of essential metals (161, 462). Several subsequent developments however make it seem unlikely that metal starvation per se is the primary mechanism

of toxicity at minimally fungistatic doses. First copper oxine is as toxic as or even more toxic than oxine itself (281 313 424). The same relation holds for the metal complexes of other chelating agents (272 372 395 433). Second there is some evidence that toxicity to bacteria and fungi is less in a medium low in metals than in the usual medium (5 9 14 358). Finally successful use of chelating agents to supply metals both to fungi (Chapter 9) and to higher plants (143 401) suggests that cells in general can compete successfully for metals with even quite stable complexes.

Metal starvation cannot however be dismissed completely as a factor in toxicity. Chelating agents reduce the metal-dependent spore pigment of *Aspergillus niger* (338). EDTA prevents spore germination of *Neurospora tetrasperma* without apparently entering the cell (406) and citrate effects on enzymatic adaptation in *A. obacter* spp. can at least be interpreted as a depletion phenomenon (368). Inhibition of sporulation by chelating agents (179 428) suggests that a higher metal requirement makes sporulation especially sensitive to metal deprivation.


A number of observations covered in detail in the original literature (9 31 32) support a plausible working hypothesis for the action of oxine and its metal complexes. The hypothesis starts with the equilibria already described (Equation 5) of oxine: its charged 1:1 chelate and the neutral 1:2 chelate. It is believed that the 1:2 chelate perhaps because of its higher lipid solubility enters the cell more rapidly than the 1:1 chelate and in that sense is the active species. However within the cell it seems likely that the 1:1 chelate is the more toxic perhaps by reacting directly with essential metal binding sites or with sulphydryl groups essential to the cell. This hypothesis has the advantage of explaining the bimodal dose response curve of oxine (6 31 281) which is similar in principle to that of the dithiocarbamates (Figure 11) and the complex interactions of metal and oxine concentrations (32). Still to be demonstrated however are the roles of free metal ions and free oxine: the site of the proposed toxic reaction and the role of natural chelating substances within and on the cell. Also as with all permeability hypotheses the lack of direct evidence is distressing.

8-Hydroxyquinoline-5-sulfonic acid is a chelating agent but is not antibacterial (6). It fails to inhibit *Monilia fructicola* but is active against *Macrosporium sarcinaeforme* (177). It seems possible that the anionic group acts by preventing access of the molecule to the site of action (6) on this hypothesis *M. sarcinaeforme* must be more permeable to the compound than are bacteria and *M. fructicola*.

The action of the dithiocarbamates and their derivatives may be ex

tested as fungitoxics (174) but it is presumed that the active group is the same in all

The important reactions of the dithiocarbamates from the standpoint of mode of action may be summarized (127)



(Symbols are taken from Figure 11 TMTM = tetramethylthiuram monosulfide)

The primary problems of mode of action in this series are set for us by (1) the bimodal response curve and (2) the effect of metals on toxicity. Although it has been considered possible that carbon disulfide formed under acid conditions (234) or enzymatically (191, 192, 384) from DMDT is the active toxicant, both this and other decomposition products are of too low inherent toxicity to account for DMDT action (322, 432).

The best available working hypothesis for the action of DMDT is rather similar to that just discussed for oxine. It is proposed (127, 196) that the first peak in the dose response curve mirrors the formation of a toxic 1:1 complex of DMDT with copper present in the medium. It is suggested on the basis of work with yeast that this water-soluble charged 1:1 complex exerts its effect at or just beneath the cell membrane (127); this suggestion, supported by indirect evidence, avoids the problem of explaining penetration of a charged complex. At higher DMDT concentrations the insoluble and non-toxic 1:2 copper complex (2 moles DMDT per copper atom) forms and accounts for the trough in the dose response curve. At still higher concentrations of DMDT either the free anion becomes toxic (196) or some other toxic complex, e.g., with zinc, is formed (127). The first of these possibilities seems more probable: free dithiocarbamate inhibits enzymes *in vitro* (319).

It is to be noted that this hypothesis explains the first peak of the bimodal response curve as a function of copper in the system: in a medium low in copper and other metals there should be only a single peak in the curve. This is borne out by visual estimates of mycelial growth (196) but has not yet been shown by the more precise spore germination technique.




Figure 11 The effect of the concentration of tetramethylthiuram disulfide (TMTD) on spore germination of *Stemphylium sarcinaeforme*. Redrawn from Diamond Horsfall Heubarger and Stoddard (91) by permission of the Connecticut Agricultural Experiment Station New Haven Connecticut

plained in a similar way. Figure 11 illustrates the well known bimodal dose response curve characteristic of this family of toxicants; the general form of the curve shown has been confirmed in studies on spore germination of *Penicillium marneffei* (298) and on growth of *Aspergillus niger* (196, 198).

The structures of interest are shown in Figure 12 with code abbreviations. Other complexes can, of course, be prepared and have been






Figure 12 The structure of some dithiocarbamate fungicides

bimodal dose response curve does not prevail in the bisdithiocarbamates (174 246) Bisdithiocarbamates are unstable (234) and several breakdown products have been suggested as the active toxicant Disodium ethylenebis[dithiocarbamate] itself is completely without fungicidal action (247 432) Carbon disulfide one of the breakdown products is unlikely to be of primary importance (432) It appears that three derived compounds which arise by breakdown of the parent substance all contribute to toxicity ethylene diisothiocyanate ethylene thiuram monosulfide and polymers of the thiuram monosulfide (197 236 237 414) Reactivity with thiol groups has suggested that the primary toxic action is on a sulfhydryl enzyme (197)

11 CONCLUDING REMARKS

In his treatise on the principles of fungicidal action Horsfall (174) emphasizes the two requirements of a toxicant that it be able in some way to react with an essential cell constituent and that it be able to reach the site of this reaction It seems fair to say that for no fungicidal or fungistatic material can both these requirements be specified in terms of structure Nor does it seem likely to me that further empirical testing of ever-different organic compounds will by itself allow us to draw up accurate specifications as guides to both theory and practice Instead we need to know more about the toxicants we have especially in terms of the theory of chemical reactions On the biological side of the problem a frontal attack on the problems of the cell surface is surely an urgent necessity It must be realized that at the present time all explanations based on an assumed permeability or impermeability are no more than guesses in the absence of independent evidence i.e. evidence not based upon the toxic reaction itself

Although no line of investigation can of course be ruled out it does seem that the search in the fungi for specific enzymatic loci of toxicity is less promising Many of the most active materials may be general rather than specific poisons if this is true they will inhibit almost any enzyme the investigator happens to test but none of these will be the site of activity

BIBLIOGRAPHY

- 1 Abelson P H and E Aldous 1950 *J Biol Chem* 60 401-413
- 2 Agerbe g L S R Schick M Schmidt and R von Sengbusch 1933 *Z. Chem* 5 212-230
- 3 Akai S and S Itoi 1954 *Botan Mag (Tokyo)* 67 1-3

Liebermeister (225) has proposed that several drugs effective against *Mycobacterium tuberculosis* are also active by virtue of their formation of a toxic complex

Some of the detailed data on the action of the dithiocarbamates is inconsistent with or at least not explained by the theory of action just outlined. These data are thoroughly reviewed by Horsfall (174) and it should be clear that many problems remain. No other hypothesis however explains the bimodal dose response curve.

Imidazoles which are themselves complexing agents antagonize the action of DMDT in the concentration range of the first peak of toxicity (198). This suggests that the locus of toxicity of the 1:1 copper chelate is an enzyme which is normally coordinated to some metal and that the available coordination positions of the copper in the 1:1 complex replace the metal on the enzyme.

Dithiocarbamates at minimally fungistatic concentrations have very slight activity against endogenous or glucose respiration in fungi (81, 127, 203, 314). Acetate oxidation in yeast is considerably more sensitive (127). The respiratory enzyme triose phosphate dehydrogenase is inhibited in yeast and in *Fusarium roseum* by TMTD (384, 385). This may be explained on the basis that the dithiocarbamates react with glutathione (432) and that the reduction of TMTD to DMDT (Equation 8) is likely to be coupled to an oxidation of such labile cell materials as glutathione. Since glutathione is the prosthetic group of triose phosphate dehydrogenase (210) it follows that this enzyme will be especially sensitive. It must be concluded that we are at present unable to point to any specific system the inhibition of which is the primary effect of either free dithiocarbamates or their active complexes.

The present picture of the action of fungistatic chelating molecules is thus one of great complexity. A given compound exists as an anion or as an uncharged molecule and as complexes of various metal ligand ratios with one or more metals. One or more of these molecular species reaches the sensitive region of the cell at which point a new set of equilibrium conditions complicated by the presence of cellular chelating agents must be envisioned. Still other complications can be introduced by structural alteration of the chelating agent such structural changes influence ionization, water and lipid solubility and the stability of metal complexes.

Although the bisdithiocarbamates are superficially similar to the dithiocarbamates and are also chelating agents the evidence to date suggests that the mode of action is not the same. In particular fungi show different sensitivities to the two classes of toxicant for a given fungus the slopes of the dose response curves are different and the

- 40 Brian P W 1944 *Nature (London)* 153 83-84
- 41 Brian P W 1945 *J Soc Chem Ind Trans (London)* 64 315-316
- 42 Brian P W 1949 *Ann Botany* 13 59-77
- 43 Brian P W P J Curtis and H G Hemming 1946 *Trans Brit Mycol Soc* 9 173-187
- 44 Brian P W, J M Wright J Stubbs and A M Way 1951 *Nature (London)* 165 317-349
- 45 Briggs G E 1954 *J Exp Botany* 5 263-268
- 46 Brink F and J M Posternak 1948 *J Cellular Comp Physiol* 32 211-234
- 47 Brook M 1952 *Nature (London)* 10 10 2
- 48 Brucker W 1957 *Arch Mikrobiol* 26 302-306
- 49 Burchfield H P and G L McNew 1950 *Contribs Boyce Thompson Inst* 16 131-161
- 50 Burchfield H P and E E Storrs 1956 *Contribs Boyce Thompson Inst* 18 395-418
- 51 Burchfield H P and E E Storrs 1957 *Contribs Boyce Thompson Inst* 18 429-452
- 52 Burriss R. H and P W Wilson 1942 *J Cellular Comp Physiol* 19 361-371
- 53 Burt, E T 1945 *Ann Appl Biol* 32 247-260
- 54 Byrde R J W J F Harris and D Woodcock 1956 *Biochem J (London)* 64 154-160
- 55 Byrde R J W J T Martin and D J D Nicholas 1956 *Nature (London)* 18 638-639
- 56 Byrde R J W and D Woodcock 1953 *Ann Appl Biol* 40 675-687
- 57 Byrde R J W and D Woodcock 1957 *Biochem J (London)* 65 682-686
- 58 Cade A R and W S Gump 1954 In G F Reddish (ed) *Antisepsis Disinfectants Fungicides and Chemical and Physical Sterilization* p 50-58 Philadelphia Lea and Febiger
- 59 Calvin M 1954 In W D McElroy and B Glass (eds) *A Symposium on the Mechanism of Enzyme Action* p 221-245 Baltimore The Johns Hopkins Press
- 60 Carl E and P Marquardt 1949 *Z Naturforsch* 4b 280-283
- 61 Cartwright K St G and W P K Findlay 1946 *Decay of Timber and Its Prevention* London H M Stationery Office pp 94
- 62 Cavill G W K and J M Vincent 1954 *Nature (London)* 155 301
- 63 Chaberek, S F C Bersworth and A E Martell 1955 *Arch Biochem Biophys* 55 321-337
- 64 Chance B 1952 *Nature (London)* 169 15-22
- 65 Christensen C. H F H Kaufert H Schmitz and J L Allison 1942 *Am J Botany* 29 552-558
- 66 Clark N G and A F Hams 1953 *Biochem J (London)* 55 839-851
- 67 Clifton C. E and W A Logan 1959 *J Bacteriol* 31 523-540
- 68 Cole L J 1952 *J Bacteriol* 64 847-853
- 69 Colhoun J 1954 *Ann Appl Biol* 41 290-304
- 70 Colwell C A and M McCall 1946 *J Bacteriol* 16 9-60
- 71 Committee on Standardization of Fungicidal Tests 1943 *Phytopathol* 33 67-632
- 72 Committee on Standardization of Fungicidal Tests 1947 *Phytopathol* 37 34-356
- 73 Corey R R and H G Shirk 1955 *Arch Biochem Biophys* 56 196-203
- 74 Crowdy S H J F Grove H G Hemming and K C Robinson 1956 *J Exp Botany* 7 42-64
- 75 Cruess W V and P H Richert 1959 *J Bacteriol* 17 363-371
- 76 Cruess W V P H Richert and J H Irish 1951 *Hilgardia* 6 305-314
- 77 Cushing J E M Schwartz and R Bennett 1949 *J Bacteriol* 8 433-44
- 78 Dajsz J and O Karkaryte 1953 *Protoplasm* 38 127-141

- 4 Albert A 1947 *Proc Intern Congr Pure and Appl Chem 11 Congr London* 4 1-5
- 5 Albert A 1950 *Proc Roy Soc (London) B* 136 177-178
- 6 Albert A 1951 *Selective Toxicity* London Methuen and Co Ltd pp 228
- 7 Albert A 1953 *Nature (London)* 172 201
- 8 Albert A 1956 *Nature (London)* 177 525-526
- 9 Albert A M I Gibson and S D Rubbo 1953 *Brit J Exp Pathol* 34 119-130
- 10 Albert A and D Magrath 1941 *Biochem J (London)* 41 534-545
- 11 Albert A S D Rubbo R J Coldacre and B G Balfour 1947 *Brit J Exp Pathol* 28 69-87
- 12 Allen R F R S Shelton and M G Van Campen Jr 1954 *J Am Chem Soc* 76 1158-1159
- 13 Ambrose J F G B Kistakowsky and A G Kridl 1951 *J Am Chem Soc* 73 1232-1236
- 14 Anderson B I and R J Swaby 1951 *Australian J Sci Research Ser B* 4 275-282
- 15 Arca Leão A E de and A da R Furtado 1950 *Mycopathol Mycol Appl* 5 121-124
- 16 Arima K and S Abe 1951 *J Antibiotics (Japan)* 4 342-346
- 17 Arny D C and C Leben 1954 *Phytopathol* 44 380-383
- 18 Badger G M 1946 *Nature (London)* 158 585
- 19 Bailly A J G G Freeman J W Cook and A R Somerville 1950 *Nature (London)* 166 65
- 20 Barron E S G L Nelson and M I Ardao 1948 *J Gen Physiol* 32 179-190
- 21 Barron E S G and T P Singer 1945 *J Biol Chem* 157 218-240
- 22 Bateman F 1943 *U S Dept Agr Tech Bull* 346 1-53
- 23 Bedford C L 1936 *Zentr Bakteriell Parasitenk Abt II* 94 102-112
- 24 Bevers H E F Celdschmidt and H Koller 1952 *Arch Biochem Biophys* 39 236-238
- 25 Beran F A Janke and G Schmidt 1954 *Phytil Phytopharm 1954* (Special Issue) 543-547 (*Chem Abstr* 50 13353 1956)
- 26 Bergman S 1953 *Acta Pathol Microbiol Scand Suppl* No 104 1-130
- 27 Berliner M D 1954 *Am J Botany* 41 93-104
- 28 Biedermann W and E Muller 1951 *Phytopathol Z* 18 301-338
- 29 Blackman G E M H Parke and G Carton 1955 *Arch Biochem Biophys* 54 45-54 65-71
- 30 Bliss C I 1935 *Ann Appl Biol* 29 131-167
- 31 Block S S 1955 *J Agr Food Chem* 3 229-234
- 32 Block S S 1956 *J Agr Food Chem* 4 1042-1046
- 33 Blumer S and J Kunder 1950 *Phytopathol Z* 17 161-199
- 34 Bodnár J and A Terényi 1930 *Z Physiol Chem Hoppe Seylers* 186 151-182
- 35 Bodnár J and T Terényi 1932 *Z Physiol Chem Hoppe Seylers* 207 1892
- 36 Bodnár J I Villányi and A Terényi 1921 *Z Physiol Chem Hoppe Seylers* 163 73-93
- 37 Bradsher C K I C Brown and R J Grantham 1954 *J Am Chem Soc* 76 114-115
- 38 von Brand T E J Tobie B Mehlman and E C Weinbach 1953 *J Cellular Comp Physiol* 41 1-22
- 39 Brewer J H 1954 In G F Reddish (ed) *Antiseptics Disinfectants Fungicides and Chemical and Physical Sterilization* p 214-240 Philadelphia Lea and Febiger

- 10 Gattani M L. 1954 *Plant pathology* 44 113-115
- 11 Geiger W B. 1946 *Arch Biochem* 11 93-32
- 12 Geiger W B. 1948 *Arch Biochem* 16 423-435
- 13 Geiger W B and J E Conn. 1951 *J Am Chem Soc* 67 11 116
- 14 Gibson I A S. 1953 *Trans Brit Mycol Soc* 6 324-334
- 15 Giese, A C. and E L Tatum. 1946 *Arch Biochem* 9 15-23
- 16 Gohre K. 1955 *Arch Forst* 4 93-301 (*Rev Appl Mycol* 6 16 1957)
- 17 Goksoyr J. 1955 *Physiol Plantarum* 8 719-835
- 18 Golden M J and K A Oster. 1947 *J Am Pharm Assoc Sci Ed* 36 359-365
- 19 Goldsworthy M C. and E L Green. 1936 *J Agr Research* 52 517-533
- 20 Goldsworthy M C. and E L Green. 1938 *J Agr Research* 56 489-506
- 21 Goldsworthy M C and E L Green. 1939 *Phytopathology* 29 700-716
- 22 Gonzalez, E. L. and E S G Barron. 1956 *Biochim Biophys Acta* 19 425-437
- 23 Gonzalez F. 1945 *Science* 101 494
- 24 Gottlieb D. 1957 *Phytopathology* 47 59-67
- 25 Gould B S M A Bosniak S Needleman and S Gatt. 1953 *Arch Biochem Biophys* 44 284-297
- 26 Gould B S N A Frigerio and W B Lebowitz. 1955 *Arch Biochem Biophys* 56 466-486
- 27 Graveri R and U Veronesi. 1956 *Exp Cell Resea ch* 11 60-67
- 28 Greig M E. 1946 *J Pharmacol Exp Therap* 87 185-199
- 29 Gross M. 1953 *Science* 118 218-219
- 30 Grove J F. 1948 *Ann Appl Biol* 35 37-44
- 31 Gruenhagen R H P A Wolf and E E Dunn. 1951 *Contribs Boyce Thompson Inst* 16 349-356
- 32 Grunberg E et al. 1950 *Trans N Y Acad Sci II* 13 2-21
- 33 Haerdtl E J and A E Martell. 1956 *J Ag Food Chem* 4 26-37
- 34 Hamilton J M D H Palmater and G L Mack. 1943 *Phytopathology* 3 533-540
- 35 Hailey J L C C McCready J K Brierley and D H Jennings. 1956 *New Phytologist* 55 1-8
- 36 Harry J B et al. 1948 *Contribs Boyce Thompson Inst* 15 195-210
- 37 Hartelius V. 1946 *Compt rend trav lab Carlsberg Sér physiol* 21 18-184
- 38 Hartelius V and K Roholt. 1946 *Compt rend trav lab Carlsberg Sér physiol* 24 163-171
- 39 Hassebrauk K. 1952 *Phytopathol Z* 19 56-8
- 40 Hawkins L A. 1913 *Physiol Resea ch* 1 57-99
- 41 Herrick J A. and J E Kempf. 1944 *J Bacteriol* 48 331-336
- 42 Hessayon G. 1953 *Soil Sci* 75 317-327
- 43 Heuberger J W and J G Horsfall. 1942 *Phytopathology* 32 3 0-3 8
- 44 Hickey R J. 1953 *Arch Biochem Biophys* 46 331-336
- 45 Hill C H. 1952 *J Biol Chem* 199 3 9-332
- 46 Hill C H. 1953 *J Bacteriol* 65 518-80
- 47 Hirschorn E and D E Munnecke. 1950 *Phytopathology* 40 5 4-546
- 48 Hirt R R. 1949 *Phytopathology* 39 31-36
- 49 Hochstein P E and C E Cox. 1952 *Phytopathology* 42 13
- 50 Hochstein P E. and C E Cox. 1956 *Am J Botany* 43 43-441
- 51 Hoffman C T R Schweitzer and G Dalby. 1940 *J Am Chem Soc* 62 988-989
- 52 Hoffman C. T R. Schweitzer and G Dalby. 1941 *Ind Eng Chem* 33 49-51
- 53 Hoffmann Ostenhof O and E Biach. 1947 *Monatsh Chem* 6 319-3 4
- 54 Hoffmann Ostenhof O and E Biach. 1948a *Monatsh Chem* 78 53-57

- 79 Danielli J F 1950 *Cell Physiology and Pharmacology* New York Elsevier Publishing Co pp 156
- 80 Danowski T S and M Tager 1948 *J Infectious Diseases* 80 119-125
- 81 Darby R T and D R Goddard 1950 *Physiol Plantarum* 3 435-446
- 82 Davies W H and W A Sexton 1946 *Biochem J (London)* 40 331-331
- 83 Davis D and A E Dimond 1952 *Phytopathology* 42 563-567
- 84 Davson H and J F Danielli 1943 *The Permeability of Natural Membranes* New York The Macmillan Co pp 361
- 85 Demerec M 1948 *J Bacteriol* 56 63-74
- 86 DiCarlo F J and S Redfern 1947 *Arch Biochem* 15 343-350
- 87 Dickey F H G H Cleland and C Lotz 1949 *Proc Natl Acad Sci U S* 35 581-586
- 88 Dillon Weston W A R and J R Boocor 1935 *J Agr Sci* 25 608-649
- 89 Dimock A W 1936 *Zentr Bakteriell Parasitenk Abt II* 95 341-347
- 90 Dimond A E D Davis R A Chapman and E M Stoddard 1952 *Conn (New Haven) Agr Exp Sta Bull* 557 1-82
- 91 Dimond A E J G Horsfall J W Heuberger and E M Stoddard 1941 *Conn (New Haven) Agr Exp Sta Bull* 451 635-661
- 92 Dimond N S 1941 *Science* 91 420-421
- 93 Dion W M and K A Lord 1944 *Ann Appl Biol* 31 221-231
- 94 Dixon M 1948 *Biochem Soc Symposia (Cambridge Engl)* No 2 39-49
- 95 Dixon M and D M Needham 1946 *Nature (London)* 158 432-438
- 96 Dunshee B R C Leben G W Keitt and F M Strong 1949 *J Am Chem Soc* 71 2436-2437
- 97 Eichhorn G L 1956 In J C Bailar Jr (ed) *The Chemistry of the Coordination Compounds* p 698-742 New York Reinhold Publishing Corp
- 98 Emerson S 1947 *J Bacteriol* 54 195-207
- 99 Engelhard H O Muller and K Bertl 1949 *Zentr Bakteriell Parasitenk Abt I Orig* 155 326-335
- 100 Everitt E L and M A Sullivan 1940 *J Wash Acad Sci* 30 125-131
- 101 Ezekiel W N 1938 *J Agr Research* 56 554-558
- 102 Fawcett C H D M Spencer and R L Wain 1957 *Ann Appl Biol* 45 158-176
- 103 Fechtmeier E T 1949 *Phytopathology* 39 605-615
- 104 Ferguson J 1939 *Proc Roy Soc (London) B* 127 387-404
- 105 Fieser L F M G Etlinger and G Fawaz 1948 *J Am Chem Soc* 70 3 8-3232
- 106 Fieser L F and H Heymann 1948 *J Biol Chem* 176 1363-13 0
- 107 Fildes P 1940 *Brit J Exp Pathol* 21 67-73
- 108 Findlay W P K and J W Vernon 1951 *Ann Appl Biol* 38 816-880
- 109 Finholt R W 1951 *Anal Chem* 23 1038-1039
- 110 Finholt R W M Weeks and C Hathaway 1952 *Ind Eng Chem* 44 101-105
- 111 Fischer E O Hoffmann E Prado and R Boné 1944 *J Bacteriol* 48 439-445
- 112 Fleury C 1948 *Ber schweiz botan Ges* 58 462-477
- 113 Foote M W J E Little and T J Sproston 1949 *J Biol Chem* 181 481-487
- 114 Forsberg J L 1949 *Phytopathology* 39 172-174
- 115 Fries N 1950 *Hereditas* 36 134-150
- 116 Fruton J S and S Simmonds 1953 *General Biochemistry* New York John Wiley and Sons pp 940
- 117 Gale E F 1949 *J Gen Microbiol* 3 369-381
- 118 Gale E F and E S Taylor 1947 *J Gen Microbiol* 1 71-85
- 119 Gale C R 1953 *J Bacteriol* 65 505-508

- 06 Kobs E. and W J Robbins 1935 *Am J Botany* 23 133-139
- 07 Koffler H F H Johnson and P W Wilson 1947 *J Am Chem Soc* 69 1113-1117
- 208 Kølmark G and N H Giles 1955 *Genetics* 40 890-902
- 209 Krebs H A R Whittam and R Hems 1957 *Biochem J (London)* 6 53-60
- 10 Krinsky I and E Racker 1952 *J Biol Chem* 198 91-9
- 211 Kuhn R and H Beinert 1943 *Ber deut chem Ges* 66 B 904-909
- 21 Kuhn R and H Beinert 1944 *Ber deut chem Ges* 77 B 606-608
- 213 Kuhn R and H Beinert 1947 *Chem Ber* 80 101-109
- 214 Kunkel L O 1914 *Bull Torrey Botanical Club* 41 63-93
- 215 Landis L D Kley and V Ercoli 1951 *J Am Pharm Assoc Sci Ed* 40 31-37
- 216 Laskowski W 1955 *Science* 121 999-1000
- 17 Leben C D M Boone and G W Kett 19 *Phytopathology* 4 467-472
- 18 Leben C and G W Kett 1950 *Phytopathology* 40 90-94
- 219 Leben C and G W Kett 1954 *J Agr Food Chem* 2 234-39
- 20 Lehman S G 1943 *Phytopathology* 33 431-448
- 21 Leonard J M and V L Blackford 1949 *J Bacteriol* 57 339-347
- 22 Leopold H and H G German 1940 *Zentr Bakteriell Parasitenk Abt II* 109 65-99
- 23 Leutnitz J Jr 1951 *Trans N Y Acad Sci II* 13 177-181
- 24 Levine S and M Novak 1949 *J Bacteriol* 57 93-94
- 5 Liebermeister K 1950 *Z Naturforsch* 5b 9-86
- 6 Lin C K 1940 *Cornell Univ Agr Exp Sta Mem* 233 1-33
- 27 Lindgren R M and M H Harvey 1959 *J Forest Products Research Soc* 2 1-7
- 28 Little J E T J Sproston and M W Foote 1948 *J Biol Chem* 174 335-347
- 29 Little J E T J Sproston and M W Foote 1949 *J Am Chem Soc* 71 1174
- 30 LoCicero J C D E H Frear and H J Miller 1948 *J Biol Chem* 172 689-693
- 231 Lockwood J L C Leben and G W Kett 1954 *Phytopathology* 44 438-446
- 32 Lohrmann W 1940 *Arch Mikrobiol* 11 59-67
- 33 Loomis W F and F Lipmann 1948 *J Biol Chem* 173 807-808
- 34 Lopatecki L E and W Newton 1952 *Can J Botany* 30 131-138
- 3 Lowry R J A S Sussman and B Heinenhain 1956 *Mycologia* 48 211-222
- 6 Ludwig R A G D Thorn and D M Miller 1954 *Can J Botany* 32 48-54
- 3 Ludwig R A G D Thorn and C H Unwin 1955 *Can J Botany* 33 42-9
- 38 McCalla T M 1940 *J Bacteriol* 40 27-32
- 39 McCalla T M 1941 *J Bacteriol* 41 75-81
- 40 McCallan S E A 1944 *Contribs Boyce Thompson Inst* 13 56-83
- 41 McCallan S E A 1947 *Proc Intern Cong Pure and Appl Chem 11th Congr London* 3 149-156
- 242 McCallan S E A 1948a *Contribs Boyce Thompson Inst* 15 77-90
- 243 McCallan S E A 1948b *Contribs Boyce Thompson Inst* 15 91-117
- 244 McCallan S E A 1949 *Botan Rev* 15 679-643
- 45 McCallan S E A 1951 *Contribs Boyce Thompson Inst* 16 299-307
- 46 McCallan S E A 1956 *Proc Intern Conf Plant Protection 2nd Conf* p 17-9
- 246a McCallan S E A and L P Miller 1957a *Contribs Boyce Thompson Inst* 18 483-493
- 246b McCallan S E A and L P Miller 1957b *Contribs Boyce Thompson Inst* 18 497-506

- 165 Hoffmann Ostenhof O and E Biach 1948b *Monatsh Chem* 79 248-252
- 166 Hoffmann Ostenhof O E Biach S Gierer and O Kraupp 1948 *Monatsh Chem* 79 576-580
- 167 Hoffmann Ostenhof O and E Kriz 1950 *Monatsh Chem* 81 90-100
- 168 Hoffmann Ostenhof O and W H Lee 1946 *Monatsh Chem* 76 180-184
- 169 Hoffmann Ostenhof O and H Moser 1948 *Monatsh Chem* 79 510-512
- 170 Hoffmann Ostenhof O and E Putz 1948 *Monatsh Chem* 79 421-425
- 171 Hoffmann Ostenhof O and E Putz 1950 *Monatsh Chem* 81 703-707
- 172 Horsfall J G 1945a *Fungicides and Their Action* Waltham Mass Chronica Botanica Co pp 239
- 173 Horsfall J C 1945b *Botan Rev* 11 321-398
- 174 Horsfall J C 1956 *Principles of Fungicidal Action* Waltham Mass Chronica Botanica Co pp 219
- 175 Horsfall J G and A E Dimond 1951a *Ann Rev Microbiol* 5 209-222
- 176 Horsfall J C and A E Dimond 1951b *Trans N Y Acad Sci* 12 338-341
- 177 Horsfall J G and S Rich 1951 *Contribs Boyce Thompson Inst* 16 313-347
- 178 Horsfall J G and S Rich 1953 *Indian Phytopathol* 6 1-14
- 179 Horsfall J G and S Rich 1955 *Trans N Y Acad Sci* 18 69-80
- 180 Horsfall J G and G A Zentmyer 1944 *Phytopathology* 34 1004
- 181 Hotchkiss R D 1946 *Ann N Y Acad Sci* 46 479-493
- 182 Hugo W B 1956 *J Gen Microbiol* 15 315-323
- 183 Hwang L and L J Klotz 1938 *Hilgardia* 12 1-38
- 184 James W O and N Garton 1952 *J Exp Botany* 3 310-318
- 185 Janke A F Beran and C Schmidt 1953 *Pflanzenschutz Ber* 10 65-87
- 186 Jarvis F G and M J Johnson 1950 *J Bacteriol* 59 51-60
- 187 Jensen H L 1947 *Proc Linn Soc N S Wales* 71 119-129
- 188 Jensen K A I Kirk G Kølmark and M Westergaard 1951 *Cold Spring Harbor Sympos a Quant Biol* 16 245-261
- 189 Jensen K A C Kølmark and M Westergaard 1949 *Hereditas* 35 521-524
- 190 Johnson F H H Eyring and M J Polissar 1954 *The Kinetic Basis of Molecular Biology* New York John Wiley and Sons pp 814
- 191 Johnston C D 1953 *Arch Biochem Biophys* 44 249-251
- 192 Johnston C D and C S Prickett 1952 *Biochim et Biophys Acta* 9 219-230
- 193 Jowett M 1938 *J Physiol* 92 322-335
- 194 Jowett M and J H Quastel 1937 *Biochem J (London)* 31 1101-1112
- 195 Jurkowska H 1951 *Bull intern acad polon sci Classe sci math nat Ser B I* 1951 161-203
- 196 Kaars Sijpesteijn A M J Janssen and G J M van der Kerk 1957 *Biochim et Biophys Acta* 23 550-557
- 197 Kaars Sijpesteijn A and G J M van der Kerk 1954a *Biochim et Biophys Acta* 13 545-552
- 198 Kaars Sijpesteijn A and G J M van der Kerk 1954b *Biochim et Biophys Acta* 13 69-77
- 199 van der Kerk G J M and J C A Luitjen 1954 *J Appl Chem* 4 314-319
- 200 Kirkham D S 1954 *Nature (London)* 173 690-691
- 201 Kittleson A R 1953 *J Agr Food Chem* 1 677-679
- 202 Klarmann E G and E S Wright 1954 In G F Reddish (ed) *Antiseptics Disinfectants Fungicides and Chemical and Physical Sterilisation* p 429-461 Philadelphia Lea and Febiger
- 203 Kloppeing H L 1951 *Chemical Constitution and Antifungal Action of Sulfur Compounds* Utrecht Schotanus and Jens pp 142
- 204 Klotz I M 1954 In W D McElroy and B Glass (eds) *A Symposium on the Mechanism of Enzyme Action* p 257-284 Baltimore The Johns Hopkins Press
- 205 Knight S G and W C Frazier 1945 *J Bacteriol* 50 502-516

- 279 Martin H R L Wain and E H Wilkinson 1942 *Ann Appl Biol* 29 41-438
- 280 Maselli J A and F F Nord 1952 *Arch Biochem Biophys* 39 406-418
- 281 Mason C L 1948 *Phytopathology* 38 740-751
- 282 Mason C L R W Brown and A E Minga 1951 *Phytopathology* 41 164-171
- 83 Mason C L and D Powell 1947 *Phytopathology* 37 5 7-528
- 84 Merkel J R and W J Nickerson 1953 *Proc Natl Acad Sci U S* 39 1008-1013
- 85 Michaelis L M and J H Quastel 1941 *Biochem J (London)* 35 518-533
- 286 Miles P G and J R Raper 1956 *Mycologia* 48 484-494
- 286a Miller H and W D McElroy 1948 *Science* 107 193-194
- 287 Miller H J 1950 *Phytopathology* 40 326-332
- 88 Miller L P and S E A McCallan 1957 *J Agr Food Chem* 5 116-120
- 89 Miller L P S E A McCallan and R M Weed 1953a *Contribs Boyce Thompson Inst* 17 151-171
- 290 Miller L P S E A McCallan and R M Weed 1953b *Contribs Boyce Thompson Inst* 17 13-19
- 291 Miller L P S E A McCallan and R M Weed 1953c *Contribs Boyce Thompson Inst* 17 282-298
- 292 Miller L P S E A McCallan and R M Weed 1954 *Radioisotope Conf Oxford Engl* 1 381-389
- 93 Minegawa T 1955 *Biochim et Biophys Acta* 16 539-5 2
- 94 Mitchell P 1954 *Symposia Soc Exp Biol* 8 254-61
- 295 Miyahara Y 1953 *Ann Phytopathol Soc Japan* 23 37-40
- 296 Molho D and L Lacroix 1949a *Bull soc chim biol* 31 1348-1356
- 297 Molho D and L Lacroix 1949b *Bull soc chim biol* 31 1357-1361
- 298 Montgomery H B S and H Shaw 1943 *Nature (London)* 151 333
- 99 Mounter L A and A J Chanutin 1953 *J Biol Chem* 204 837-846
- 300 Mucke D 1954 *Flora (NF)* 41 30-50
- 301 Munhead I 1949 *Ann Appl Biol* 36 250-2 6
- 302 Muller E and W Biedermann 1950 *Phytopathol Z* 19 343-3 0
- 303 Muller Stoll W R 1950 *Phytopathol Z* 17 265-86
- 304 Napier E J D I Turner and A Rhodes 1956 *Ann Botany* 0 461-466
- 305 Nason A and A R Lehman 1955 *Science* 120 19-
- 306 Neilands J B and P K Stumpf 1955 *Outlines of Enzyme Chemistry* New York John Wiley and Sons pp 315
- 307 Newton B A 1956 *Bacteriol Revs* 0 14-27
- 08 Neyland M P Dunkel and A L Schade 1955 *J Gen Microbiol* 7 409-416
- 309 Nickerson W J 1946 *Science* 103 484-486
- 310 Nickerson W J and N J W Van Rij 1949 *Biochim et Biophys Acta* 3 461-475
- 311 Nelsen L W 1942 *Cornell Univ Ag Exp Sta Mem* 248 1-44
- 312 Nielsen L W and C E Williamson 1944 *Phytopathology* 34 1026-10 0
- 313 Nordbr ng Herz B 1955 *Physiol Planta um* 8 691-717
- 314 Ohta Y 1944 *Acta Phytochim (Japan)* 14 159-166
- 315 Okazaki R 1956 *Exp Cell Res* 10 489- 04
- 316 Olcott H S and H F Aenkel Conrat 1947 *Chem Revs* 41 151-197
- 317 Oster K A and M J Golden 1947 *J Am Pharm Assoc Sci Ed* 36 283-288
- 318 Oster K A and M J Golden 1954 In G F Reddish (ed) *Antiseptics Disinfectants Fungicides and Chemical and Physical Sterilization* p 132-141 Philadelphia Lea and Febiger
- 319 Owens R G 1953a *Contribs Boyce Thompson Inst* 17 1- 40

- 247 McCallan S L A L I Miller and R M Weed 1934 *Contribs Boyce Thompson Inst* 18 99-98
- 248 McCallan S E A and C Satterstrom 1910 *Contribs Boyce Thompson Inst* 11 325-340
- 249 McCallan S E A and F R Weedon 1940 *Contribs Boyce Thompson Inst* 11 331-342
- 250 McCallan S E A and R H Wellman 1942 *Contribs Boyce Thompson Inst* 12 451-461
- 251 McCallan S E A and R H Wellman 1943 *Contribs Boyce Thompson Inst* 13 93-134
- 252 McCallan S E A R H Wellman and F Wilcoxon 1941 *Contribs Boyce Thompson Inst* 12 49-78
- 253 McCallan S E A and F Wilcoxon 1931 *Contribs Boyce Thompson Inst* 3 13-38
- 254 McCallan S E A and F Wilcoxon 1942 *Contribs Boyce Thompson Inst* 4 233-244
- 255 McCallan S E A and F Wilcoxon 1933 *Contribs Boyce Thompson Inst* 5 173-180
- 256 McCallan S F A and F Wilcoxon 1934 *Contribs Boyce Thompson Inst* 6 419-500
- 257 McCallan S E A and F Wilcoxon 1936 *Contribs Boyce Thompson Inst* 8 151-160
- 258 McCallan S E A and F Wilcoxon 1939 *Contribs Boyce Thompson Inst* 11 5-20
- 259 McCallan S E A and F Wilcoxon 1940 *Contribs Boyce Thompson Inst* 11 309-324
- 260 McClellan W D 1912 *Phytopathology* 32 394-398
- 261 McElroy W D 1947 *Quart Rev Biol* 22 25-58
- 262 McGowan J C P W Brian and H G Hemming 1918 *Ann Appl Biol* 35 25-36
- 263 McKee R K 1931 *Nature (London)* 167 611
- 264 MacLeod R A 1939 *J Biol Chem* 197 751-761
- 265 McNew G L and H I Burchfield 1930 *Contribs Boyce Thompson Inst* 16 163-170
- 266 McNew G L and H P Burchfield 1931 *Contribs Boyce Thompson Inst* 16 357-374
- 267 McNew G L and N K Sundholm 1949 *Phytopathology* 39 791-791
- 268 Mandels G R and R T Darby 1934 *J Bacteriol* 65 16-26
- 269 Mandels C R and R G H Siu 1950 *J Bacteriol* 60 249-262
- 270 Manten A H L Klopping and G J M van der Kerk 1930a *Antonie van Leeuwenhoek J Microbiol Serol* 16 45-55
- 271 Manten A H L Klopping and G J M van der Kerk 1930b *Antonie van Leeuwenhoek J Microbiol Serol* 16 282-294
- 272 Manten A H L Klopping and G J M van der Kerk 1931 *Antonie van Leeuwenhoek J Microbiol Serol* 17 58-68
- 273 Marsh P B 1915 *Phytopathology* 35 54-61
- 274 Marsh P B and M L Butler 1946 *Ind Eng Chem* 38 701-703
- 275 Marsh P B M L Butler and B S Clark 1949 *Ind Eng Chem* 41 216-2184
- 276 Martell A E and M Calvin 1952 *Chemistry of the Metal Chelate Compounds* New York Intenice Hall pp 613
- 277 Martin G J 1931 *Biological Antagonism* New York The Blakiston Co pp 16
- 278 Martin H 1940 *The Scientific Principles of Plant Protection* 3rd ed London Edward Arnold pp 385

- 68 Schutter J and P W Wilson 1950 *J Gen Microbiol* 12 446-454
- 69 Sclarini L J and F F Nord 1943 *Arch Biochem* 3 251-267
- 70 Sempio C and M Castori 1949 *Riv biol (Perugia)* 41 163-172
- 71 Severens J M and F W Tanner 1945 *J Bacteriol* 49 383-93
- 72 Sexton W A 1953 *Chemical Constitution and Biological Activity* 2nd ed New York D Van Nostrand Co Inc pp 424
- 73 Shaw W H R 1954 *Science* 10 61-63
- 74 Shirk H G and H E Byrne 1951 *J Biol Chem* 191 183-186
- 75 Shirk H G and R R Corey Jr 1950 *Arch Biochem Biophys* 38 417-423
- 76 Shirk H G R R Corey Jr and P L Poelma 1951 *Arch Biochem Biophys* 3 92-99
- 77 Shirk H G P L Poelma and R R Corey Jr 1951 *Arch Biochem Biophys* 3 386-391
- 78 Shrift A 1954 *Am J Botany* 41 23-230 345-352
- 79 Simon E W 1953a *J Exp Botany* 4 317-392 393-402
- 80 Simon E W 1953b *Biol Revs Cambridge Phil Soc* 28 453-479
- 81 Simon E W and H Beevers 1951 *Science* 114 124-16
- 82 Simon E W and H Beevers 1952 *New Phytologist* 51 163-190 191-197
- 83 Simon E W and G E Blackman 1949 *Symposia Soc Exp Biol* 3 25-26
- 84 Sisler H D and C E Cox 1954 *Am J Botany* 41 338-45
- 85 Sisler H D and C E Cox 1955 *Am J Botany* 42 3 1-3 6
- 86 Sitt R G H 1951 *Microbial Decomposition of Cellulose* New York Reinhold Publ Corp pp 531
- 87 Sletten O and C E Skinner 1948 *J Bacteriol* 56 6 9-681
- 88 Smalt Sister M A C W Kreke and E S Cook 1957 *J Biol Chem* 224 999-1004
- 89 Smith E L N C Davis E Adams and D H Spackman 1954 In W D McElroy and B Glass (eds) *A Symposium on the Mechanism of Enzyme Action* p 291-312 Baltimore The Johns Hopkins Press
- 90 Smith F G J C Walker and W J Hooker 1946 *Am J Botany* 33 351-356
- 91 Smith J H 1951 *Ann Appl Biol* 8 1-50
- 92 Snell J M and A Weissberger 1959 *J Am Chem Soc* 61 4 0-453
- 93 Snow D and P S Watts 1945 *Ann Appl Biol* 32 102-111
- 94 Soo Hoo G and E Grunberg 1950 *J Invest Dermatol* 14 169-170
- 95 Sorokin E W Roth and H Eilenmeyer 1952 *Helv Chim Acta* 3 1736-1741
- 96 Stallings J H 1954 *Bacteriol Revs* 18 131-146
- 97 Starkey R L and S A Waksman 1943 *J Bacteriol* 45 509-519
- 98 Stedman R L E Kravitz and J D King 1951 *J Bacteriol* 3 650-660
- 99 Steinberg R A 1940 *J Ag Res ch* 60 165-773
- 100 Stevens C M and A Mylroie 1953 *Am J Botany* 40 424-429
- 101 Stewart I and C D Leonard 1952 *Science* 116 564-66
- 102 Stoddard E M and A E D Mond 1949 *Biol Rev* 15 34-3 6
- 103 Stokes A 1954 *Plant and Soil* 5 132-141
- 104 Strecker B 1957 *Z Pflanzenkr* 64 9-35
- 105 Sumner J B and G F Somers 1953 *Chemistry and Methods of Enzymes* 3rd ed New York Academic Press pp 46
- 106 Sussman A S 1954 *J Gen Physiol* 38 59-17
- 107 Sussman A S and R J Lowry 1955 *J Bacteriol* 10 615-685
- 108 Sute C M 1941 *Chem Revs* 28 69-299
- 109 Tatum E L 1950 *J Cellular Comp Physiol* 35 Suppl 1 119-131
- 110 Tatum E L and G W Beadle 1949 *Proc Natl Acad Sci U S A* 34 13
- 111 Tehon L R 1951 *Science* 114 663-664
- 112 Teply L J 1949 *Arch Biochem* 24 383-388
- 113 Ter Horst W P and E L Felix 1943 *Ind Eng Chem* 3 1255-1259

- 320 Owens R C 1933b *Contribs Boyce Thompson Inst* 17 273-282
- 321 Page J E and F A Robinson 1943 *J Chem Soc (London)* 1943 133-135
- 322 Palmer H C R W Greenlee and M M Baldwin 1952 *Phytopathology* 42 472
- 323 Parker Rhodes A F 1942a *Ann Appl Biol* 29 126-145
- 324 Parker Rhodes A F 1942b *Ann Appl Biol* 29 404-411
- 325 Pauson P 1955 *Chem Revs* 55 9-136
- 326 Pennington R J 1957 *Biochem J (London)* 65 534-540
- 327 Petit A 1930 *Ann service botan et agron Tunisie* 6 57-70
- 328 Pfeiffer R E Schraufstatter F Gehring and J Schuk 1949 *Z Naturforsch* 4b 344-350
- 329 Phillips C R 1949 *Am J Hyg* 60 280-288
- 330 Phillips J N and J M Vincent 1948 *Nature (London)* 161 210-211
- 331 Pickett M J and C E Clifton 1943 *J Cellular Comp Physiol* 2 141-165
- 332 Pinckard J A et al 1939 *Phytopathology* 29 177-187
- 333 Plakidas A G 1938 *Phytopathology* 28 307-329
- 334 Pope H and I Christison 1953 *J Bacteriol* 66 639-644
- 335 Potter V R and A E Reif 1952 *J Biol Chem* 194 281-297
- 336 Pramer D 1956 *Arch Biochem Biophys* 62 265-273
- 337 Priddyham T G et al 1956 *Phytopathology* 46 568-575
- 338 van Raalte M H 1952 *Mededel Landbouwhogeschool en Opvoetinstas Staat Gent* 17 163-173 (*Chem Abstr* 46 11543 1952)
- 339 Rahn O and J E Conn 1944 *Ind Eng Chem* 36 185-187
- 340 Ramsey C B M A Smith and B C Herberg 1944 *Botan Gaz* 106 14-23
- 341 Rangaswami G 1956 *Mycologia* 48 800-804
- 342 Reavill M J 1954 *Ann Appl Biol* 41 448-460
- 343 Reif A and V R Potter 1954 *Arch Biochem Biophys* 48 1-6
- 344 Rennerfelt E 1948 *Physiol Plantarum* 1 245-254
- 345 Rennerfelt E and G Nacht 1955 *Scensk Botan Tidskr* 49 419-434
- 346 Rich S and J G Horsfall 1952 *Phytopathology* 42 457-460
- 347 Rich S and J G Horsfall 1954 *Proc Natl Acad Sci U S A* 40 139-145
- 348 Richards C A and R M Addoms 1947 *Proc Am Wood Preservers Assoc* 1947 41-58 (*Chem Abstr* 42 4702-4703 1948)
- 349 Richardson L T 1954 *Can J Botany* 32 335-346
- 350 Rigler N E and G A Creathouse 1940 *Am J Botany* 27 701-704
- 351 Rinderknecht H J L Ward F Bergel and A L Morrison 1947 *Biochem J (London)* 41 463-469
- 352 Ringel S M and F S Bencke 1956 *Mycologia* 48 329-336
- 353 Roblin R O Jr 1946 *Chem Revs* 38 255-377
- 354 Robinson G N 1954 *J Appl Bact* 17 190-195
- 355 Roper J A and E Kafer 1957 *J Gen Microbiol* 16 660-667
- 356 Rothstein A 1954 *Symposia Soc Exp Biol* 8 164-201
- 357 Rothstein A and A D Hayes 1956 *Arch Biochem Biophys* 63 81-99
- 358 Rubbo S D A Albert and M I Gibson 1950 *Brit J Exp Pathol* 31 425-441
- 359 Sampson R E and R A Ludwig 1953 *Can J Botany* 31 531-536
- 360 Sampson R E and R A Ludwig 1956 *Can J Botany* 34 37-43
- 361 Sansome E R 1949 *Trans Brit Mycol Soc* 32 305-314
- 362 Sansome F R 1956 *Trans Brit Mycol Soc* 39 67-78
- 363 Schmidt E W 1924 *Centr Bakteriell Parasitenk Abt II* 60 329-338
- 364 Schopfer W H and E C Grob 1949 *Helv Chim Acta* 32 8 9-838
- 365 Schopfer W H and M Guillaud 1946 *Helv Physiol Pharmacol Acta (Compt rend)* 4 4-95
- 366 Schueler F W 1947 *J Infectious Diseases* 81 139-146
- 367 Schuldt P H and C N Wolf 1956 *Contribs Boyce Thompson Inst* 18 317-393

- 455 Work T S and E Work 1948 *The Basis of Chemotherapy* New York Interscience Publishers pp 435
- 456 Wyss O B J Ludwig and R R Joiner 1947 *Arch Biochem* 1: 415-43
- 457 Yarwood C E 1937 *Phytopathology* 27 931-941
- 458 Yarwood C E 1943 *Hilgardia* 14 595-691
- 459 Yarwood C E 1950 *Phytopathology* 40 178-189
- 460 Yarwood C E 1955 In T H Sternberg and V D Newcomer (eds) *Therapy of Fungus Diseases* p 130-135 Boston Little Brown and Co
- 461 Zentmyer G A 1943 *Phytopathology* 3 1191
- 462 Zentmyer G A 1944 *Science* 100 294-295
- 463 Zentmyer G A 1955 *Phytopathology* 45 398-404

- 414 Thorn G D and R A Ludwig 1954 *Can J Chem* 32 872-879
- 415 Thornberry H H 1950 *Phytopathology* 40 419-429
- 416 Treffers H P 1956 *J Bacteriol* 72 108-114
- 417 Trim A R and A E Alexander 1949 *Symposia Soc Exp Biol* 3 111-149
- 418 Tröger R 1956 *Arch Mikrobiol* 25 166-192
- 419 Valko E I 1946 *Ann N Y Acad Sci* 46 451-478
- 420 Vallee B L and F L Hoch 1955 *Proc Natl Acad Sci U S A* 41 327-338
- 421 Velick S F 1956 *Ann Rev Biochem* 25 251-290
- 422 Verdcourt B 1952 *Mycologia* 44 317-386
- 423 Verrall A F 1919 *J Agr Research* 78 695-703
- 424 Vicklund R E M Manowitz and V J Bagdon 1954 *Mycologia* 46 133-142
- 425 Villemin I F H A Lechevalier and S A Waksman 1953 *Intern Congr Microbiol 6th Congr Rome* 5 146-173
- 426 Vincent J M 1917 *J Soc Chem Ind* 66 149-155
- 427 Volk W A 1954 *J Biol Chem* 208 777-784
- 428 Volkmann C M and E Beerstecher Jr 1955 *J Bacteriol* 70 416-480
- 429 Wain R L and C H Wilkinson 1945 *Ann Appl Biol* 32 240-243
- 430 Waksman S A A H Romano H Lechevalier and F Raubitschek 1952 *Bull World Health Organisation* 6 163-173
- 430a Walker A T 1955 *Iowa State Coll J Sci* 30 229-241
- 431 Walker A T and F G Smith 1952 *Proc Soc Exp Biol Med* 81 556-559
- 432 Weed R M S E A McCallan and L I Miller 1953 *Contribs Boyce Thompson Inst* 17 299-315
- 433 Weinberg E D 1957 *Bacteriol Revs* 21 46-68
- 434 Weiss S J V Fiore and F F Nord 1949 *Arch Biochem* 22 314-322
- 435 Weissman G S and S F Trelease 1955 *Ann J Botany* 42 489-495
- 436 Wellman R H and S E A McCallan 1942 *Contribs Boyce Thompson Inst* 12 431-450
- 437 Wellman R H and S E A McCallan 1946 *Contribs Boyce Thompson Inst* 14 151-160
- 438 West B and F T Wolf 1955 *J Gen Microbiol* 12 396-401
- 439 Whaley F R and J B Harry 1951 *Contribs Boyce Thompson Inst* 16 355-386
- 440 Whiffen A J 1950 *Mycologia* 44 253-258
- 441 Wilcoxon F and S E A McCallan 1930 *Phytopathology* 20 391-417
- 442 Wilcoxon F and S E A McCallan 1931 *Contribs Boyce Thompson Inst* 3 509-528
- 443 Wilcoxon F and S E A McCallan 1932 *Contribs Boyce Thompson Inst* 4 415-424
- 444 Wilcoxon F and S E A McCallan 1935 *Contribs Boyce Thompson Inst* 7 333-339
- 445 Wilcoxon F and S E A McCallan 1939 *Contribs Boyce Thompson Inst* 10 329-338
- 446 Winzler R J 1940 *J Cellular Comp Physiol* 15 343-354
- 447 Wolf C N P H Schuldt and M M Baldwin 1955 *Science* 121 61-62
- 448 Wolf F A et al 1940 *Phytopathology* 30 213-227
- 449 Wolf F T 1947 In W J Nickerson (ed) *Biology of the Pathogenic Fungi* p 91-114 Waltham Mass Chronica Botanica Co
- 450 Wolf F T and F A Wolf 1950 *Bull Torrey Botan Club* 77 77-82
- 451 Wolf P A and W M Westveer 1950 *Arch Biochem* 28 201-206
- 452 Woods D D 1940 *Brit J Exp Pathol* 21 74-89
- 453 Woolley D W 1945 *Proc Soc Exp Biol Med* 60 225-228
- 454 Woolley D W 1952 *A Study of Antimetabolites* New York John Wiley and Sons pp 269

- 455 Work, T S and E Work. 1948 *The Basis of Chemotherapy* New York Interscience Publishers pp 435
- 456 Wyss O B J Ludwig and R R Joiner 1945 *Arch Biochem* 7 41-4
- 457 Yarwood C E 1937 *Phytopathology* 27 931-941
- 458 Yarwood C E 1943 *Hilgardia* 14 595-691
- 459 Yarwood C E 1950 *Phytopathology* 40 173-179
- 460 Yarwood C E 1955 In T H Sternberg and V D Newcomer (eds) *Therapy of Fungus Diseases* p 130-135 Boston Little Brown and Co
- 461 Zentmyer G A 1943 *Phytopathology* 33 1171
- 462 Zentmyer G A 1944 *Science* 100 294-295
- 463 Zentmyer G A 1955 *Phytopathology* 45 398-404

Organism Index

(Binomial names only)

- Abidia* 342 346
blakesleeana 149
glauca 40
spinosa 183
Acetobacter 21
Achlya 373 374
ambisexualis 373
americana 373
biuxualis 373
c. oliviana 40
prolifera 23
Actinomyces 1, 13^o 133
israeli 123 302
Ag. cus (see also *Pisallot*) *campestris*
 2 3 365
melleus 7
Agroclype dura 164
Albugo candida 407
Allomyces 15 159 396 43
arbuscula 40 132 164 306 311 407
javanicus 68 69 81 159 162 248 49
 302
anepus 61 331
macrognus 51
Alnus glutinosa 249
Alternaria 364 366
brassicae 358 40
Alternaria citri 404
oleracea 39 446 473
solani 24 247 327 3 9 3,9 388 390
 441 448
tenuis 180 183 468
Amanita 255
mappa 274
muscaria 73
phalloides 253
Anabaena cylindrica 300
Aph. ionyces eutesches 20
Apodachlya 132
brachynema 248 258
A. miliaria 18
mellea 9 24 27 122 134
Arthrobacter terreus 344 345
Ascobolus 397
stercorarius 315
Ascochyta pisi 316
Asco. dea rubescens 3 8 335
Ashbya gossypii 40 214 216 214 336-
 337 343
Aspe. gilius 25 36 47 46 47 78 86 91
 100 102 107 109 114 115 119 122
 13 133 134 135 137 144 149 153
 1 1 135 1 6 172 1,8 1,9 180 181
 183 184 06 14 216 217 246 2,0

- 253 264 265 275 278 302 306 308
 309 325 328 342 346 347 363 389
 404 406 408 431
aureus 219
batatae 249
echinulatus 390
elegans 122
fischeri 43 45 149 150 390
flavus 20 44 82 179 219 249 281
 282 283 302 316 427
foetidus 118
fumaricus 141 147
fumigatus 16 17 49 130 146 167
 221 227 255 267 280
glaucus 26 280
itaconicus 134 144
janus 364
luchuensis 106 108 394
melleus 430 431
nidulans 17 45 46 48 152 169 213
 305 306 339 390 393 402 449
niger 3 9 20 23 26 36 37 38 42
 43 44 46 49 62 63 64 65 83 88
 100 107 108 109 110 114 117 119
 121 123 131 132 134 135 137 139
 140 141 142 144 145 146 152 170
 174 176 177 180 182 204 208 209
 212 214 216 218 219 220 224 225
 228 241 242 244 245 249 250 261
 263 265 278 279 301 302 303 305
 306 307 308 310 312 313 314 316
 328 334 337 343 370 371 312 316
 392 395 404 413 415 434 435 441
 444 461 468 473 475 481 482
ochraceus 157 309
oryzae 8 9 23 36 37 40 65 81 91
 107 108 110 111 112 113 114 119
 153 178 187 219 224 231 232 252
 253 268 215 219 302 312 419
parasiticus 121 253
quadrilineatus 168
sydowi 43 46 47 48 120
terreus 133 134 144 145 183 306-
 307 316 429-430 433-434 435
versicolor 78 81 83
azotobacter 481

Bacillus globigii 414
subtilis III 256
Basidiobolus ranarum 40

Blakeslea trispora 331 368
Blastocladia 366
pringsheimii 23 24 69 137 133 147
 318 337-338 316
ramosa 340
Blastocladiella 230 366 391 396 411
emersonii 206 224 229 248 266 306
 331 316-377 428
variabilis 40
Blastomyces (see also *Paracoccidioides*)
 11
brasiliensis 206
dermatitidis 10 11 40 46 47 163
 334 454
Boletus 346
bovinus 122
edulis 40 342
granulatus 346
variegatus 20 331
Botryosporium 20
Botrytis 115 133
alli 441 477-478
cinerea 17 110 117 152 133 134 361
 390 400 412 413 415 426
squamosus 428
Bremia lactucae 364
Brevlegnia gracilis 69 91 305
Byssosclamyces fulva 115 116 127 424
 425

Caldariomyces fumago 133 183
Calonectria decora 157
Candida 306 336 346
albicans 10 302 480
pulcherrima (see also *Torulopsis pul-*
cherrima) 283
tropicalis 10
Cantharellus cibarius 40
Capnodium salicinum 15
Cenococcum graniforme 259
Centrospora acerina 428
Cephalosporium acremonium 390 413
constantii 14
salmosynnematum 304
Cephalothecium roseum 156 309
Ceratostomella (see also *Endoconidio-*
phora Ophiostoma) 331 331 316
fimbriata 331 363 371 312
ips 334
microspora 328

- Ceratostamella pinz* 334
ulmi 10 132
Cercospora beticola 18
k kuchi 20
sesami 363
Cercospora herpotrichoides 331
Chaetomium 253 312 342 346 371
aureum 167
funicola 247
globosum 107 114 115 371 372 429 435
thermophile 18
Chalara quercina 20 65 10 75 16 91
Chalaropsis thelaxioides 331 409
Chlorella vulgaris 161 210 410
Choanephora cucurbitarum 247 308 359 364 365
Chondrus crispus 119
Chytridium 20 66 68 13 132 247 248
Circinella 133 342
Citromyces 137
Cladosporium 17 400
Clasterosporium 133
carpophilum 410
Clavaria 83
Claviceps paspali 219
purpurea 46 47 108 264 271 213 274 2 9 363
Clitoria 18
d atrela 164
Clostridium 157
acetobutylicum 336
Coccid o des immitis 43 121 217
Coccomyces hemalis 408 415
Colletotrichum 405
atirrhini 156
destructum 389
gloriosoides 402
g aminicolum 49
hibisci 0
idicum 407
laga a ium 409
li de nutt iu i 21 364
lin 421
pho o des 115
Collybia 18 83
tuberosa 331
tel tips 30 331 36 406
Coniophora cereblla 24 132 133 134
Coniothyrium 156
Coprinus 14 20 21 70 246 217 408
atramentarius 40
ephemerus 311
fimetarius 13 17
hiascens 208
lagopus 331 357 358
radians 37 41 203
similis 167
Cordyceps militaris 122 82 318
Corticium 78 164
centrifugum 134 144
Cortinari 112
glaucopus 331
Cronartium ribicola 407
Cryptococcus (see also *Torulopsis*) 121 300
neoformans 43
Ctenomyces fa inulentis 40
Cunninghamella 133 253 313
blakesleeana 156 107
Curvularia 156
lunata 157
Cylindrocarpon radicola 107
Cytosporina ludibunda 368
Dacrymyces stillatus 159
Dactylum dendroides 156
Daedalea flavida 115
juniperina 160 181
Dahlia 106
Dematium pullulans 10 132 134 130 214 221
Dermocybe sanguinea 168
Diaporthe phaseolorum 339 358 310
Dictyostelium 336
discoideum 41 06
Didymella vadakii 156
Didymium eunigrates 358 362
Diploma pon osae 247 339
Diplodia macrospora 57
natalis 261 277
tubericola 214
Drosophila subtrata 183
Eberthella typhosa 409
Empusa see *Entomophthora*
Eidoconidiophora (see also *Ceatocto mell Ophiostoma*)
adiposa 63
Jagadea 400 400 407 424 4 6

- 2,3 264 265 275 278 302 306 308
 309 325 328 342 346 347 363 389
 404 406 408 431
aureus 249
batatae 249
echinulatus 390
elegans 122
fischeri 43 45 149 150 390
flavus 20 44 82 1,9 219 249 281
 282 283 302 3,6 427
foetidus 118
fumaricus 141 147
fumigatus 16 17 49 13, 146 167
 221 227 2,5 267 280
glaucus 26 280
itaconicus 134 144
janus 364
luchuensis 106 108 394
melleus 430 431
midulans 17 45 46 48 152 169 273
 305 306 339 390 393 402 449
niger 3 9 20 23 26 36 37 38 42
 43 44 46 49 62 63 64 6, 83 88
 100 107 108 109 110 114 117 119
 121 123 131 132 134 135 137 139
 140 141 142 144 145 146 152 170
 174 176 177 180 182 204 208 209
 212 214 216 218 219 220 224 225
 228 241 242 244 245 249 250 261
 263 265 278 2,9 301 302 303 305
 306 307 308 310 312 313 314 316
 328 334 337 343 370 371 3,2 3 6
 392 39, 404 413 415 434 435 441
 444 461 468 473 4,5 481 482
ochraceus 157 309
oryzae 8 9 23 36 37 40 65 81 91
 107 108 110 111 112 113 114 119
 153 178 187 219 224 231 232 252
 2,3 268 2,5 279 30, 312 429
parasiticus 121 253
quadricaneatus 168
sydowi 43 46 47 48 1,0
terreus 135 134 144 145 183 306-
 307 316 429-430 433-434 435
versicolor 78 81 83
Azotobacter 481

Bacillus globigii 414
subtilis 111 2,6
Basidiobolus ranarum 40

Blakeslea trispora 331 368
Blastocladiella 366
pringsheimii 23 24 69 13, 133 142
 328 337-338 3 6
ramosa 340
Blastocladiella 230 366 391 396 411
emersonii 206 224 223 248 266 306
 331 3,6-3,7 428
variabilis 40
Blastomyces (see also *Paracoccidioides*)
 11
brasilensis 206
dermatitidis 10 11 40 46 47 163
 334 434
Boletus 346
bovinus 122
edulis 40 342
granulatus 346
variegatus 20 331
Botryosporium 20
Botrytis 115 133
allii 441 477-4,8
cinerea 17 115 117 132 133 134 364
 390 40, 412 413 415 4,6
squamosus 428
Bremia lactucae 364
Brevilegnia gracilis 69 91 30, 30, 30, 30
Byssochlamys fulva 115 116 1,2 424
 425

Caldanomyces fumago 133 183
Calonectria decora 157
Candida 306 336 346
albicans 10 302 480
pulcherrima (see also *Torulopsis pul-*
cherrima) 283
tropicalis 10
Cantharellus cibarius 40
Capnodium salicinum 15
Cenococcium graniforme 259
Cenitrospora acerina 4,8
Cephalosporium acremonium 395 4,3
costantii 14
salmosynnematum 304
Cephalothecium roseum 1,6 3,9
Cerafostomella (see also *Endoconidio-*
phora Ophiostoma) 331 33, 346
fimbriata 331 363 371 372
ips 334
microspora 328

- Loctarius* 47 164
Lactobacillus 221
 arabinosus 260
 casei 342
 leichmanii 343
 pentosus 221
Lagenidium giganteum 40
Larix 114
Lentinus degener 167
 lepideus 165 461
 omphalodes 259
Lenzites abietina 156
 sepiaria 115 15 410
 trabea 17 346
Leptota naucina 80
Leptographium 208
Leptomitium lacteus 60 81 248 " 8
 60
Leuconostoc mesenteroides 216
Lobaria retigera 167
Lophodermium pinastri 339
Lycogala 108
Lycoperdon giganteum 153

Macrochytrium botrydioides 68 137
Macrophomina phaseoli 115
Macrosporium sarcinaeforme (see also
 Stemphylium sarcinaeforme) 175
 406 481
Magnusia 408 410
 brachytricha 405
 nitida 407
Malbranchea pulchella 184
Maasmus 21 18 83 302 346
 chordalis 134
 fulvobulbillosus 331
 graminum 20
 perforans 331
 rotula 21
Mastigosprium album 17 407
Melampsora lini 391 407 426
Melanconium fuligineum 367
Melanosporea
 destuens 369 372
 zamae 369
Memnoniella echinata 60 70 81 91
 133 327 334 394 402
Merulius 133 134
 confluens 20

Merulius domesticus 83 302
 lacrymans 15 119 132
 niveus 143
Metarrhizium 115
Micrococcus denitrificans 244
Micromonospora 19 254
 vulgaris 18 209
Microsporium 84 265
 audouinii 337
 canis 302
 gypseum 84 306
Mitula paludosa 333
Monascus 347
Monilia (see also *Sclerotinia*) 13 133
 formosa 133 135
 fructigena 115
Monilinia (see also *Sclerotinia*) *fructicola* 9 36 390 39 445 446 451
 473 474
 laxa 476
Monoblepharella 132
Monotostoma lanuginosa see *Thermomyces lanuginosus*
Morchella esculenta 38
Mortierella 133 214
 renispora 251 253
Mucor 40 46 115 137 133 156 214
 250 342 346 413
 dispersus 390 430 431
 guillermondii 10 06
 hiemalis 161 342
 mucedo 82 133 102 374
 plumbeus 132
 pusillus 17
 pyriformis 133
 ceramosus 115 2 1
 ramannianus 331
 ouxii 132
Mycena 27 8 83 403
Mycobacterium 3
 paratuberculosis 344
 tuberculosis 344 484
Mycoderma vini 40
Mycogone perniciosa 0
Mycosphaerella citrullina 66
 pinodis 247 366
Myrothecium verrucaria 106 107 113
 180 20 08 209 217 231 47 209
 327 393 39 400 407 447

- Endoconidiophora fimbriata* 331
moniliformis 134
Endomyces 46 349
decipiens 40
magnusii 331
Endomycopus 149
fibuliger 40
vernalis 265 267
Entomophthora 66 75 86 247
coronata 91
Epidermophyton floccosum 40
Eremaurus fertilis 40
Eremothecium ashbyi 40 259 209 216
 336 337 339-340
Erysiphe 36 424
cichoracearum 408
graminis 363 364 318 393 403 406
 409 410 412 423 428
polygoni 405 407
Escherichia coli 87 209 227 268 271
 310 433 434

Fomes 78 331
annosus 20 102 133 134 403 443
fraxineus 20 115
Fumago vagans 132 133 183
Fusarium 18 46 61 114 115 118 133
 134 143 145 149 170 214 215 216
 217 221 228 255 273 281 283 308
 328 342 346 359 362 366 368 405
 415
aqueductum 20
aurantiacum 20
avenaceum 346
caeruleum 449
conglutinans 363
eumartii 431
graminearum 46 133 273
javanicum 268
lani 123 132 136 151 204 210 214
 216 219 271 246 250 408 421 474
moniliforme 406 410
nivale 17
niveum 246
oxysporum 9 10 11 91 110 117 163
 234 255 256 2 9
roseum 149 401
solanii 6 157 207 208 317 366
tricothecoides 358
vasinfectum 283

Gelasinospora tetrasperma 231
Geotrichum 20 149
candidum 302
Gibberella bacatta 156
fujikuroi 281 283
saubinetii 19
Gliocladium catenulatum 157
roseum 37
Gliomastix concolata 68
Gloeosporium fructigenum 115
Glomerella 254
cingulata 114 115 175 176 177 338
 346 345 401 407 431 473
Gnomonia 17
Gonatortracheia highleyi 312
Gristia confuens 164
Gymnosporangium 395
clavipes 441
globosum 403
Gyrophora esculenta 169

Hansenula anomala 245
Helicostylum piniforme 156
Helminthosporium 172
gramineum 244 359 377
oryzae 400
racenelsii 168
sativum 61 308 389 390
victoriae 234
Hendersoniella 17
Herpobasidium deformans 407
Herpomycetes stylopygae 41
Herpotrichia nigra 17
Hirsutella 347
Hustoplasma 11
capsulatum 10 40 306
farcinosum 10
Hormodendrum 207
Hydnium ferrugineum 167
Hypholoma fasciculare 40 331

Isaria farinosa 347

Karlingia rosea (see also *Rhizoplyctis rosea*) 247 428

Labyrinthula minuta 209
utellii 345

ORGANISM INDEX

- Penicillium terlikowskii* 280
terrestre 180
urticae 63 64 172 309
va ans 43
uniferum 130
uestlingii 49
Peronospora 408
destructor 364
effusa 407
tabacina 398
Pestalotia 389
malorum 30
Peziza 156 164
Phacidium infestans 17 426
Phallus impudicus 273
Phlyctothiza variabilis 20 338
Pholiota aurea 250
Phoma 49
apicola 17 363 391 407
betae 30
terrestris 83
Photobacterium fischeri 27
Phragmidium mucronatum 396 407
411 420
Phycomyces 328 331 360 361 362
blakesleeana 13 20 40 46 48 151
156 19 160 162 160 171 258 263
26 302 34 328 330 332 361 363
30 397 400 414 480
nitens 361 364 46
Phycopsis 17
Phymatotrichum omnivorum 16 37 42
43 63 67 66 246 247 406 461
Physoctenidia obtusa 410
Physarum polycephalum 03 253 366
Phytophthora 40 114 331 399 412
cactorum 70 91
cinnamomi 332
erythroseptica 115
infestans 389 399 407 410 411
pinnulae 17
Pichia membranaefaciens 40
Pilobolus 314 310 360 362
crystallinus 374
kleinii 361
Picula aoryae 247 253 28 260
331 363 360
Pleodomus meliloti 19
Pleurotus lunatellus 27
Polychytrium aggregatum 66 68 84 91
Polyporus 78 115 158 217 331
anceps 133
annosus 83
arcularius 357
benzoinus 165
betulinus 107 302
eucalyptorum 158
hanedai 27
texasus 340
tumulosus 134 160
vaporarius 14 132
versicolor 176 331
Polystictus 78 115
Polytomella caeca 60
Poria cocos 158
vaillantii 134 336
Proactinomyces (see also *Nocardia*) 81
82 83
erythropolis 155
opacus 82 162 163
roseus 156
Propionibacterium pentosaceum 449
Psalliota (see also *Agaricus*) *bisporia* 71
80 81 80 91 247 401
campestris (see also *P. bisporia*) 360
402
Pseudomonas 221 222 227
fluorescens 174 220
putrefaciens 60
Pseudoperonospora humuli 308
Pseudopeziza ribis 363
Puccinia 159 392
chrysa themei 411
cononata 309 400
glumarum 408
graminis 46 163 393 40 410 414
415 48 431
helianthi 414
malvacearum 407
Puccinia pullulans 14 79
Pyronema confluens 307 360
Pythium citrophthora 278
Pythium 20 10 91 132
Pythium 4 247 305
butleri 326
debarianum 14 40 115
Rhipidium 130
Rhizoctonia 115
solanii 18 389

- Nectria cinnabarina*, 40 281
 coccinea 372
 galligena 264
Neurospora 261 326 339 342 394 397
 crassa 5 7 9 10 22 25 38 47 63 67
 87 88 108 115 118 121 133 134
 145 151 152 156 159 160 161 163
 171 172 174 175 176 181 204 210
 214 216 218 219 224 226 228 229
 231 244 245 256 258 261-263 265
 266 277 280 302 304 306 308 310
 312 313 325 326 328 331-342 363
 365 3 6 390 391 393 401-402 413
 424 426 428 429 430 431-432
 433 449
 sitophila 133 254 266 335 365 392
 395 442 451 473 475
 tetrasperma 230 232 391 393-394
 397 401 412 469 481
Nigrospora sphaerica 365
Nitella clavata 452
Nocardia (see also *Proactinomyces*) 431
 343
 acidophilus 164
 corallina 249 265 266
 opaca 155
 sevivorans 424

Oidium 149
Omphalia flavida 27 264
Onygena equina 84 398
Oospora 46 149
 colorans 167
 sulphurea ochracea 166
Ophiobolus graminis 20 23 41
 herpatrichus 156
 myrabeanus 461
Ophiostoma (see also *Ceratostomella*
 Endoconidiophora) 335 346 363
 multiannullatum 10 271 266 2 7
 305 340 100
 pini 333

Paecilomyces varioti 49
Panus stipticus 26
Paracoccidioides brasiliensis (see also
 Blastomyces) 10 11 40
Parmelia 169
Paxillus atromentosus 167

Penicillium 42 43 45 46 48 49 86 91
 100 102 115 132 133 134 135 137
 141 144 149 156 165 166 167 172
 173 174 178 179 180 182 183 184
 214 216 247 252 253 261 263 264
 268 280 302 325 328 342 344 346
 347 370 404 406 424 431 441
 amarum 168
 brevi compactum 43 132 165 173
 221
 capreolinum 43
 charlesii 42 43 180
 chrysogenum 37 44 47 62 81 101
 117 118 135 136 143 157 173 174
 184 186 187 203 208 209 211 212
 216 219 224 225 226 228 230 230
 261 264 265 268 275 302 305 328
 331 432
 cinerascens 180
 crustosum 47
 cyclospium 166 283
 digitatum 46 66 72 75 81 91 111
 134 163 227 341
 duponti 18
 expansum 20
 fumiculosum 119
 gladioli 166
 glaucum 20 42 82 154 156 221 461
 griseofulvum 8 37 133 280
 islandicum 163
 janczewskii 81
 javanicum 46 48 81 86
 lilacinum 48
 luteopurpurogenum 132
 luteum 43 306
 malgouensis 183
 nigricans, 24
 notatum 37 38 153 154 184 185
 216 220 253 262 269 306 308 428
 429 432
 obscurum 280
 puberulum 169
 reticulosum 133 142 220 280
 roquesforti 152
 roseo purpureum 174
 rugulosus 43
 sanguineum 261
 sclerotiorum 43
 spinulosum 107 109
 stipitatum 169

ORGANISM INDEX

- Penicillium terlikowskii* 280
terrestre 180
urticae 63 64 172 309
varians 43
viniferum 133
westlingii 49
Peronospora 408
destructor 364
effusa 407
tabacina 398
Pestalotia 389
malorum 303
Peziza 156 164
Phacidium infestans 17 426
Phallus impudicus 273
Phlycto hiza variabilis 20 338
Pholiota aurea 250
Phoma 249
apicola 17 363 391 407
betulae 310
terrestris 283
Photobacterium fischeri 27
Phragmidium mucronatum 396 407
411 425
Phycomyces 328 331 360 361 362
blakleeanus 13 20 40 46 48 131
136 159 160 162 163 171 258 263
266 302 314 318 320 332 361 363
310 397 402 414 480
nitens 361 364 426
Phycopsis 17
Phymatotrachium omnivorum 16 37 42
43 63 67 16 246 247 426 461
Physalospora obtusa 410
Physarum polyccephalum 203 203 366
Phytophthora 40 114 331 399 412
cactorum 70 91
cinnamomi 332
erythroseptica 115
infestans 389 399 407 410 411
p. mularae 17
Pichia membranaefaciens 40
Pilobolus 344 345 360 362
crystallinus 374
kleinii 361
P. icularia oryzae 247 253 258 266
331 363 366
Plenodomus meliloti 19
Pleurotus lunatus 27
Polychytrium aggregatum 66 68 84 91
Polyporus 78 115 158 247 331
anceps 133
annosus 85
arcularius 357
benzoinus 165
betulinus 107 302
eucalyptorum 158
hanedai 27
texanus 342
tumulosus 134 163
vaporarius 14 132
versicolor 116 331
Polytictus 18 115
Polytomella caeca 60
Poria cocos 158
vallantii 134 336
Proactinomyces (see also *Nocardia*) 81
82 83
erythropolis 135
opacus 82 162 163
roseus 156
Propionibacterium pentosaceum 449
Psalliota (see also *Agaricus*) *bispora* 11
15 80 81 85 91 247 401
campestris (see also *P. bispora*) 363
402
Pseudomonas 221 222 227
fluorescens 174 220
putrefaciens 60
Pseudoperonospora fumuli 338
Pseudopeziza visibis 363
Puccinia 159 392
chrysanthemi 411
coronata 309 40
glumarum 408
graminis 46 163 393 403 412 414
415 418 431
helianthi 414
malvacearum 407
Pullularia pullulans 14 79
Pyronema confluens 337 363
Pythaeocystis citrophthora 218
Pythaeogeton 20 10 91 132
Pythium 4 247 303
butleri 326
debarianum 14 40 11
Rhizidium 132
Rhizoctonia 115
solani 18 389

- Nectria cinnabarina* 40 281
 coccinea 372
 galligena 264
Neurospora 261 326 339 342 394 397
 crassa 5 7 9 20 22 25 38 47 63 67
 87 88 108 115 118 121 133 134
 145 151 152 156 159 160 161 163
 171 172 174 175 176 181 201 210
 214 216 218 219 224 226 228 229
 231 244 245 256 258 261-273 275
 276 277 280 302 304 306 308 310
 312 313 325 326 328 331-342 363
 375 376 390 391 393 401-402 413
 424 426 428 429 430 431-432
 433 449
 sitophila 133 254 276 333 375 392
 393 442 451 473 475
 tetrasperma 230 232 391 393-394
 397 401 412 469 481
Vigrospora sphaerica 363
Vitella clavata 452
Noctardia (see also *Proactinomyces*) 331
 343
 acidophilus 164
 corallina 249 275 276
 opaca 155
 sevivorans 474

Oidium 149
Omphalia flavida 27 264
Onygena equina 84 398
Oospora 46 149
 colorans 167
 sulphurea ochracea 166
Ophiobolus graminis 20 23 413
 terpochrichus 156
 myabeanus 461
Ophiostoma (see also *Ceratostomella*)
 Endoconidiophora 333 346 363
 multiannulatum 10 271 276 277
 303 340 400
 pini 333

Paecilomyces varioti 49
Panus stipticus 26
Paracoccidioides brasiliensis (see also
 Blastomyces) 10 11 40
Parmelia 169
Paxillus atromentosus 167

Penicillium 42 43 45 46 48 49 86 91
 100 102 115 132 133 134 135 137
 141 144 149 156 163 166 167 172
 173 174 178 179 180 182 183 184
 214 216 247 252 253 261 263 265
 278 280 307 305 328 342 344 346
 347 370 404 406 424 431 441
 amarum 168
 brevi compactum 43 137 165 173
 221
 capreolium 43
 charlessi 42 43 180
 chrysogenum 37 44 47 6 81 107
 117 118 135 136 143 157 173 174
 184 186 187 203 208 209 211 212
 216 219 224 225 276 228 240 260
 261 264 265 268 275 302 303 328
 334 432
 cinerascens 180
 crustosum 47
 cyclopium 166 283
 digitatum 36 66 72 75 81 91 111
 134 163 327 341
 dupontii 18
 expansum, 20
 funiculosum 119
 gladioli 166
 glaucum 20 47 82 154 155 21 461
 griseofulvum 8 37 133 280
 islandicum 168
 janczewskii 81
 javanicum 46 48 81 86
 lilacinum 48
 luteopurpureogenum 132
 luteum 43 306
 nalgoviensis 183
 nigricans 24
 notitum 37 38 153 154 184 185
 216 220 253 262 269 306 308 428
 429 432
 obscurum 280
 puberulum 169
 reticulosum 133 147 270 280
 roqueforti 152
 roseo purpureum 174
 rugulosus 43
 sarguineum 261
 serotiorum 43
 spinulosum 107 109
 stiptatum 169

- Streptomyces scabies* 0 113 263 413
thioluteus 280
venezuelae 166 260 264 269
violaceus ruber (see also *S. coelicolor*)
 251
Syncephalastrum racemosum 113 183
Synchytrium endobioticum 412

Tetrahymena pyriformis 228
Thamnidium 363
elegans 115
Thelephora 167
Thermascus aurantiacus 18
Thermomyces lanuginosus 18
Thielaviopsis basicola 88 115
paradoxa 389
Th. australis clacata 313
Thuja plicata 4 5
Tilletia 403 408
carries 216 18 266 466
horrida 331
levis 273
tritici 331 467
Torula 40 46 149
cremoris 302
utilis (see also *Torulopsis utilis*) 263
Torulopsis 43 149
neoformans see *Cryptococcus neoformans*
pulcherrima (see also *Candida pulcherrima*) 308
rotundata 120
sa guinea 402
utilis 210 246 268 2 8 302
T. ameles 113
cinnabarina 283
sua colens 38 16
Tanzschelia punctata 407
Trichoderma 114 115
koningi 251
ignorum 44
viride 40 183 80 309 312 3 8 389
 426
Tricholoma 21 44 8
equestre 302
flavobrunneum 66
fumosum 113
gibbosum 251 258
imbricatum 258
nudum 20
Trichophyton 84 258 259 316
discoideus 3 8 331 333
equinum 338 339
faviforme 339
ferrugineum 40
gypseum 252
mentagrophytes 258 279 431
persicolor 20 256
rosaceum 251
rubrum 123 302
tonsurans 134 259 331
T. chosporon capitatum 10
cutaneum 20
Trichothecium 114
roseum 156
Typhula 17

Uncinula necator 410 414
Urocystis tritici 400 403
Uromyces phaseoli 414 426 420
Ustilago 20 331 346 313 403 408
avenae 412
major 346
scabiosae 258
sphaerogena 230 231 309 323 330
 344
striatiformis 397
violacea 12 40 71 75 86
zeae 3 44 47 108 134 151 206 219
 216 303 407 412 415 429 431
Ustilina vulgata 157 391

Venturia inaequalis 44 63 257 306
 338 342 403 415 449 482
Verticillium 247
albo atrum 12 113 389 40
coccorum 119
malhousii 20
puillioiae 20
tr corpus 14

Vylaria mali 241

Zygo hynchus 346
moelleri 143 224 226
vuilleminii 24
Zygosaccharomyces 312
zoster 40

- Rhizoglyphus rosea* 75 246 247
Rhizopus 18 115 132 133 136 141 156
 214 215 216 251 311 328 342 348
 363 408
bovinus 40
chinensis 17 407
delemar 111
japonicus 37 38 122
nigricans 9 115 134 157 203 221
 227 228 229 302 342-343 390 392
 395 407 473 474
oryzae 9 136 215 216 228
suinus 57 263 328 339 431
Rhodotorula aurantiaca 340
gracilis 150 151
rubra 40 159 160 161
Romania terricola 10
Russula aurata 273

Sabouraudia audouinii (see also *Micro-
 sporium audouinii*) 40
Saccharomyces 40
cerevisiae 345
Saprolegnia delicata 69 91 300
ferax 392
lapponica 40
mixta 6 363 367
Sapromyces elongatus 132 248
reinschii 373
Scenedesmus obliquus 313
Sclerophyllum 78
commune 20 182 181 426
Schizosaccharomyces 40
Schizothecium longicollis 74
Sclerospora 358 364
Sclerotinia (see also *Botrytis Monilia
 Monilinia*) 134 400
camelliae 376 359 340
cinerea 80 110
fruticicola 15 40 175 207 316 359
 391 401 409 410
fructigena 428
laxa 443
Sclerotium (see also *Rhizoctonia*) 331
bataticola 17 115
delphinii 72 132 133 134 144
rolfsii 328 332
Scopulaopsis brevicaulis 5 180 190
 242 243 244 246 255

Sepedonium 17 20
Septoria apii 405
pepi 20
Sordaria fimicola 9 88 326 331 331
 403
Sphaerobolus 365
Sphaerotheca pannosa 363 399
Sphaerulina trifolia 331
Spicaria violacea 119
Sporobolomyces roseus 40
Sporodinia grandis 7 364 367 368
Sporotrichum 17
schenckii 10 40 333
Stachybotrys atra 60 63 73 91 113 115
 122 332
Staphylococcus aureus 480
Stemphylium (see also *Macrosporium*)
radicinum 214
sarcinaeforme 395 446 473 474 476
 482
Stereum 78
frustulosum 25 331
gausapatum 71
hirsutum 132
murrai 331
purpureum 110
Streptococcus faecalis 210
lactis 342
Streptomyces 2 9 17 21 23 41 43 61
 79 80 84 106 111 114 119 132
 133 136 156 157 249 251 252 254
 255 275 277 281 282 311 312 328
 343 346 390 400 432 433 435 462
alba niger 183
albus 120
aureofaciens 155 183
coelicolor (see also *S. violaceus ruber*)
 10 63 73 76 91 133 140 143 201
 208 216 218 219 222 274 220 278
 243 256 283 302
flavescens 431 432 434
flavus 47
fadiae 6 143
griseus 36 110 208 225 276 278 243
 264 266 278 342 343 366 431 433
limosus 283
microflavus 112
nitrificans 219 316
olivaceus 344
proteolyticus 252

- Agrocybin 164 285
 α Alanine metabolism of 257 260 267
 β Alanine 342
 Alcohols effect of on acid formation 141
 toxicity of 457-459 461
 utilization of 83
 see also individual compounds
 Aldolase 204 213 216
 Alkylation and toxicity 463
 N Alkyl α bromoacetamides 461
 N Alkylpyridines 461
 Allantoic acid 4, 278-279
 Allantoinase 279
 Allantoin 218-219
 Allantoinase 279
 Allithiamine 331
 Alternanol 180
 Aluminum 38 307 313
 Amanitins 255
 Amicetin 266 283
 Amidases 265
 Amides 253 258 265
 Amines 273-274 453 461
 α Amino acid oxidases 260-261
 ϵ Amino acid oxidases 261-262
 Amino acid racemases 259
 Amino acids acyl, 265
 Amino acids as autolysis products 8
 as chelating agents 315 49
 biosynthesis of 138 265-272
 deamination of 260-262
 effect of on respiration 211
 of mycelium 44 45
 oxidation of 260-265
 requirements for 248 259
 toxicity of 258 259
 transport of 59-260
 utilization of 83 256-265
 α Aminoadipic acid 270-271
 p Aminobenzoic acid 171 326 327 329
 340-341 345 461
 biosynthesis of 271-272
 γ Aminobutyric acid 265-267
 Aminoethanol and methylated derivatives 181
 Amino hydroxycaproic acid 271
 4 Amino 5 imidazole carboxamide 26
 8 Amino levulinic acid 344
 Amino peptidases 253
 1,2 Amino-1 propanol 219
 Amino sugars 121-122
 Ammonia as autolysis product 8
 in amino acid metabolism 270 265
 266 310
 physiological effects of 249 328 365
 requirement for 326 335
 Ammonium dehydrogenase 249
 Ammonium ion physiological effects of
 19 247-248 391 403
 utilization of 210 213-214 246-249
 333
 Ammonium nitrate utilization of 243-
 244
 Amphomycin 254
 Amygdalin 65-66
 Amylases 110-112 477
 Anaerobic respiration 202
 Aneurin *see* Thiamine
 Aniline 83
 Anisaldehyde 16
 Anthranilic acid 211-212
 Anthraquinones 168 170 172 477
 Antibiotics 440
 Antimalarials 477
 Antimycins 449 462
 Aquatic fungi 23 21 317
 Arabans 114
 D -Arabinose 63 68-73
 L -Arabinose 63 68-73
 D -Arabitol 122
 Arabogalactan 114
 Arabonic acid 130 136
 Arachidic acid 48
 Arginase 28 411
 Arginine metabolism of 260 64 268-
 69 278
 requirement for 259
 utilization of 257
 L Argininosuccinic acid 269
 Aromatic compounds metabolism of
 164-177
 Arsenate 447 448
 Arsenic 182 449 463
 Arsenite 449
 Arsenous acid 182
A. throbacter terreagens factor 344-345
 Arylthioalkanecarboxylic acids 412
 Ascorbic acid 180 345 393
 Ascocin 46

Subject Index

- Accumulation of amino acids* 219-260
 of metals 316 472
 selective 450-452 467
Acetaldehyde 213 214
Acetamide 258 265
Acetanilide derivatives 464
Acetic acid 161 337
 as buffer 22
 formation of 133 143 221
 metabolism *cf* 138 143 145 223 224
 227 462 484
 utilization of 68-73 81
Acetic dehydrogenase 207
Acetoacetic acid 153 161 263
Acetoin 123 214
Acetone formation of 154
Acetonedicarboxylic acid 146
N-Acetyl p-aminobenzoic acid 341
Acetyl choline 213
Acetyl coenzyme A 138 143 202-223
 227
N-Acetyl D-glucosamine 39 121
Acetyl phosphate 221 225 227 228
Acids toxicity of 4
Aconitase 223 227
as Aconitic acid 134 138 223
as Aconitic decarboxylase 145
Acrasin 375
Actidines 4 6-457
Actinavine 449
Actinomycins 254 475
Actiniazic acid 282 283 472
Acylases 265
Adenine 275-277 283 337
Adenosine 275 276
Adenosine diphosphate 215
Adenosine diphosphate ribose 215
Adenosine phosphates 38 203 213 301
Adenosine 3 phosphate 275
Adenosine 5 phosphate 215 277 302
Adenosine triphosphate 302
Adenyl deaminase 275
Adenylc acid *see* Adenosine 3 phosphate
 Adenosine 5 phosphate
Adenylosuccinic acid 277
Adonitol 68-73
Adsorption 452 468-469 472
Aerobic respiration 202
Agar impurities in 325
 in culture media 1 9-10 12-14 324
Agaric acid 47
Age and respiration 206-207
Agmatine 215
Agroclavine 219-280

SUBJECT INDEX

- Carboxypeptidase 253
 Carlic acid 180
 Carlotic acid 180
 Carnosine 240-251
 Carotic acid 180
 Carotinic acid 180
 α Carotene 159 161
 β Carotene 158 159 161
 γ Carotene 159 161 377
 ξ Carotene 159 161
 θ Carotene 159
 Carotenoids 34 3 6
 biosynthesis of 160-162
 formation of 158-160 377
 function of 162 361
 Carvacrol 411
 Casein 241 301
 Catalase 221 231-232 07 008 3 4 471
 Catechol 174 17 063
 Cell disruption 20
 Cell wall 39-41
 Cellobiase 108 113
 Cellobiose 107 108 113
 utilization of 56 66 68- 3
 Cellotriose 113
 Cellulase 113
 Cellulose 40-41 368
 hydrolysis of 110-113
 utilization of 68-73 77-79 83-84 90
 Cephalins 47 153
 Cephalosporin N 184
 Cephalosporin P₁ 158
 Cerebrins 47-48
 Cesium 303 316
 Chelating agents 306 312 315-316 449
 toxicity of 442 454 466-467 471 472
 4 8-184
 Chemical potential 458-459
 Chemostat 3
 Chemotherapy 440
 Chemotropism 391-392
 Chitin 34 37 39-41 377 393
 utilization of 39-80
 Chitinases 119
 Chlamydo pores 3 6 366
 Chloramphenicol 160 166 183
 Chloranil *see* Tetrachloro *p* benzoquinone
 Chlorate toxicity of 443
 Chlorination biological 183-184
 Chlorine 313
p Chloromercuribenzoic acid 463 470
 Chloromycetin *see* Chloramphenicol
 2 Choro 1 4 naphthoquinone 344
 Chlorotetracycline 183
 see also Aureomycin
 Cholesterol 155
 Cholic acid 441
 Choline 38 181-182 273
 Choline sulfate 45
 Chromates 466
 Chromium 313
 Chrysophanic acid 168 170
 Citric acid 22 310 316
 formation of 133 204 225 308-309
 metabolism of 138 219 223- 27
 utilization of 68-73 81
 Citric acid cycle 138 222-226
 Citrimin 308
 Citrovorum factor 342
 L Citrulline 269 277
 Clavacin *see* Patulin
 Clavatin *see* Patulin
 Claviformin *see* Patulin
 Cobalamins *see* Vitamins B₁₂
 Cobalt 253 312 313 316 343
 Cocarboxylase *see* Thiamine pyrophosphate
 Cocoa butter utilization of 82
 Coconut oil utilization of 82
 Coenzyme A 152 161 342
 Colchicine 170
 Compactose 43
 Competitive inhibition 461-469
 Condensing enzyme 243 246 227
 Copper 314 315
 physiological effects of 139 184 309-310 340
 requirement for 309-310
 toxicity of 307 446 447 449 451
 465-472
 uptake of 316 470-452 467-472
 Copper oxine 481
 Copper sulfate 466
 Coprogen 344-345
 Cordycepin 122 280 283
 Corn steep liquor 186
 Cotton as culture medium 3
 Cresosote 448
 Cresols 455

- Asparaginase 26, 41
 Asparagine 265 321 340
 requirement for 258
 utilization of 256-268
 Aspartase 262 266
 Aspartic acid metabolism of 267 265
 266 267 269 270 333
 utilization of 256 257
 Aspergillus acid 28° 283
 Aspergillus 308
 Asperthecin 169
 Asperxanthone 170
 Atramentum 167
 Aureomycin 183 402
 Aureothricin 280
 Autoclaving effects of 51-58
 Autolysis 7 41 241-242 303 304 375-376
 Auxin *see* Indoleacetic acid
 Aversion 374
 Avidin 333
 8 Azaguanine 277
 Azaserine 280 281
 Azelaic acid 334
 Azide 231 455 480

 Barbituric acid 213
 Barium 306
 Barrage 344
 Bavendamta reaction 83-84 115
 Behenic acid 48
 Benzaldehyde 173-174
 Benzamide 265
 Benzoic acid 174 265
 o Benzoquinone 175
 p Benzoquinones formation of 167 170
 toxicity of 415-477
 Benzothiazoles 441 472
 Beryllium 307
 Betaine 183 243
 Betulin 157
 Bicarbonate effect of on reproduction 316-377
 Bioassay of minerals 313-314
 of vitamins 328-329
 Biocyan 334
 Bioluminescence 26-27
 Biotin 283 325 326 327 328 329 332-333 343 371 402
 Biotin sulfone 335
 Biotin sulfoxide 334
 Bisdiisocarbamates 484-485
 Bisphenols 464-465
 Bisulfite 305
 Borax 449
 Bordeaux mixture 466
 Boric acid as buffer 22
 Boron 313
 Bound water 26 36 106
 Bran as culture medium 3
 Bromide ion 183
 Buffers 22
 Bufotenine 274
 1,2 Butanediol 123
 2 Butanol 214
 2 Butanone 214
 2,3 Butylene glycol 123 211
 Butyric acid 153 154

 Cacodylate sodium 189
 Cadaverine 273
 Cadmium 308
 Caffeine 276 277 443
 Calcium 19 38 311-312 311
 Calcium cyanamide 218
 Caldariomycin 183
 Callose 41
 Camphor 390 441
 Canavanine 268 210
 Caprocinose 43
 Capsular polysaccharides 43 191
 Captan 442
 O Carbamyl D-serine 280
 Carbohydrate of mycelium 37 42-43
 of spores 393
 Carbon total 36
 Carbon balance 100-101
 Carbon dioxide fixation of 138 14
 146 210 223 228-229 333 334
 physiological effects of 24 209 363
 366 316-317 413
 Carbon disulfide 483 485
 Carbon monoxide 231
 Carbon nutrition and reproduction 367-369
 and spore germination 401
 Carbonic anhydrase 471
 Carboxylase *see* Pyruvic carboxylase
 Carboxymethylcellulose 77-78
 cis cis β Carboxymuconic acid 174

- Dissociation 454-455
 Disulfide 30 393
 Dithiazolidines 472
 Dithiocarbamates 451 454 472 481-484
 diethyl 480
 dimethyl 471 480 482
 Diurnal cycles 358
 Dithionate 305
 Docosanoic acid 48
 Dodecane 87 163
 Dodecanoic acid 48 154
 Dormancy 395-398
 Dry weight determination of 11-12
 Dulcitol 64 68-73
 Dyes 441 455 464

 Eburicic acid 157 158
 Echinulin 280
 Ecology and vitamin requirements 346-347
 Economic coefficient 89-90
 Edestin 251
 Egg albumen 251
 Eicosanoic acid 48
 Embden Meyerhof pathway 212-217 219
 Emodin 168
 Endogenous respiration 210-211 463
 Endopeptidases 251-253
 Enniatins 255
 Enolase 213
 Enzymes adaptive 15 86-88 207-208
 inactivation of 467-468 470-472 476-477
 induced 75 86-88 207-208
 Epoxides 443
trans-L-Epoxysuccinic acid *see* Ethylene oxide α,β dicarboxylic acid
 Erdin 183
 Ergonovine 2,9
 Ergosterol 49 82 155
 palmitate 49
 Ergot compounds 266 279-280
 Ergothioneine 273-274
 Erythritol 6 68-73 122
 Erythrose-4-phosphate 218
 Erythrulose 123
 Esterases 1 2
 Ethanol 57 3 8 329
 formation of 141 213-216 221
 metabolism of 83 143 214 227
 Ethanol dehydrogenase 213 309
 Ethanolamine 47
 Ethyl acetate 134 143
 Ethyl disulfide 187
 Ethylene 163
 Ethylene chlorhydrin 178-179
 Ethylenediaminetetraacetic acid 265 316 467 480 481
 Ethylene disothiocyanate 485
 Ethylene glycol 123
 Ethylene oxide derivatives, 464
 Ethylene oxide α,β dicarboxylic acid 135 146
 Ethylene thiuram monosulfide 485
 Exopeptidases 251-253
 Expansin *see* Patulin

 Fat coefficient 150
 Fats formation of 149-151
 Fatty acid dehydrogenase 152
 Fatty acids biosynthesis of 151-152 333 462
 oxidation of 153-158
 toxicity of 475 461
 Ferguson's principle 4 8-49
 Fermentation 207 212-217 221-222 394
 Fermentative assimilation 212
 Ferrichromes 309 344-345
 Ferricyanide 205 463
 Ferulic acid 174
 Fibrin 251
 Flavin adenine dinucleotide 220 231 244-245 461 336
 Flavin in enzymes, 231 337
 Flavins as photoreceptors 361 62
 Flavonol 283
 Fluoride 448 449 455
 2-Fluoro-*p*-aminobenzoic acid 340
 Folic acid *see* Pteroylglutamic acid
 Formic acid formation of 138 143 144
 metabolism of 181-182 227 337
 utilization of 68-73 82
 Formylkynurenine 338
N-Formylpteronic acid 342-343
 Freezing injury 425-427
 Fructans 76
 Fructose 60-67 68-73 122
 Fructose 1,6 diphosphate 204 213 418

- Cryptoxanthin 159
 Crystal violet 447 475
 Culture shaken 2
 Cupric ammonium sulfate 466 467
 Cyanide 231 250 455 480
 Cyanocobalamin 337 343-344
 see also Vitamins B₁₂
 Cycloheximide 282 283 442 449
 Cyclopaldic acid 166 172
 Cyclophenin 283
 Cyclopolic acid 166 172
 Cycloserine 280
 Cystathionine 270
 Cysteine 186 205 470
 metabolism of 210 305-306
 utilization of 257 305
 Cysteine desulphydrase 264
 Cysteine sulfinic acid 306
 Cistine 186 305
 Cytase *see* Protopectinase
 Cytidylic acid 277
 Cytochrome oxidase 230-231 411 480
 Cytochrome peroxidase 232
 Cytochrome reductases 230
 Cytochromes 204 229-231 309 330 344
 317 462
 Cytosine 216 283

 Dechlorogriseofulvin 183
 Dehydroacetic acid 172 477
 Dehydrofusanic acid 283
 Δ Dehydropipecolic acid 262
 5 Dehydroquinic acid 171
 5 Dehydroshikimic acid 171
 2 Deoxyglycose 64 220
 4 Deoxypyridine 335
 Deoxyribonucleic acid 41 275
 Depsides 169 170
 Depsidones 169 170
 Dermatophytic fungi 16 21 84 212 218
 Despeptido actinomycin 254
 Desihobiotin 327 333-334
 Detoxification 449 476
 Dextranase 119
 Dextrin 68-73 76
 Dextrinase 119
 Dialkylfluorophosphatase 480
 2,6 Diaminopurine 216-277
 Diastase *see* Amylase
 Diatermynes 164 283

 Diazomethane 443
 Dichlone *see* 2,3 Dichloro-1,4 naphthoquinone
 2,4 Dichloro 6 (o chloroanilino) 1 tria
 zine 457
 1,2 Dichloro 4,5 diamonibenzene 337
 2,3 Dichloro 1,4 naphthoquinone 344
 467 475 416
 2,6 Dichlorophenolindophenol 204 205
 Dichromate potassium 448
 Diethanolamine 209
 Diethylarsonic acid 182
 Digallic acid 116
 Dihydroxyacetone phosphate *see* Triose
 phosphates
 α β Dihydroxy β ethylbutyric acid 210
 α β Dihydroxy β methylbutyric acid 210
 Dihydroorotic acid 277
 2,3 Dihydroxyphenylacetic acid 114
 2,4 Dihydroxyphenylalanine 263
 2,5 Dihydroxyphenylpyruvic acid 263
 3,4 Dihydroxyphthalic acid 165
 2,3 Dimercaptopropanol 410
 Dimethylamine 483
 Dimethylaminoethanol 181
 5,6 Dimethylbenzimidazole 344
 1,2 Dimethyl 4,5 diaminobenzene 337
 343
 Dimethylglycine 181
 Dimethylglyoxime 442
 Dimethylpyruvic acid 135 116
 Dimethyl selenide 182
 Dimethyl sulfide 181
 Dimethyl telluride 182
 Dimorphic fungi 10-11 206 327 333
 m Dinitrobenzene 245
 3,5 Dinitro o cresol 462
 2,4 Dinitrophenol 186 210 432 443
 456 467
 Diosgenin 157
 Dipeptidases 213
 Diphenyl 441
 Diphenylamine 160
 Diphenylthiocarbazone 311
 Dihydropyridine nucleotidase 215
 309 393
 Diposphopyridine nucleotide 1,2 204
 215 217 223 229 338
 Diploican 170
 α α Dipyridyl 479 480

- Growth linear 12-13
 measurement of 11-13 1
 rate of 5-10 102 374
- Guanidine 2/8
- Guanidinobutyramide 264
- Guanine 2/5 2/7 337 3/0
 effect of on spore germination 402
- Guanosine 2/6
- Guanylic acid 2/6
- Gyrophoric acid 169
- Hallochrome 463
- Heavy metals toxicity of 465-472
see also Copper Mercury Silver Uranium
- Helvetic acid 158
- Hemipigments 308
- Hemicelluloses 42 6-17 113-114
- O Heterobiotin 334-335
- Heterothallism 3/4-3/5
- γ Hexachlorocyclohexane 340
- Hexachlorophene 464-46
- Hexacosanoic acid 48
- Hexacosenoic acid 48
- n Hexadecane 163
- Hexadecanoic acid hydroxy 4/
- Hexadecenoic acid 48
- Hexanoic acids 154
- Hexokinase 913
- Hippuric acid 26
- Hippuricase 2 3 96
- Histamine 3 3 0
- Histidine 313
 accumulation of 259
 metabolism of 257 264 261 264
 requirement for 2 0-21 2 9
- L-Histidyl phosphate 2 1
- Histidylhistidine 0-2 1
- Histozyne 26
- Homocysteine 181 2/0 341
- Homocentisic acid 63
- Homologous series and toxicity 4 7-461
- Homoprotocatechuic acid 165
- Homoserine 4/0
- Homonies in reproduction 3/3-3 5
- Hughes press 20
- Humidity effect of on growth 25
 effect of on reproduction 364-365
 effect of on spore germination 5
 399-400 404-406
- Hydrocarbons metabolism of 82 163-164
 toxicity of 457 464
- Hydrogen peroxide 231
- Hydrogen sulfide 4/3-475
- Hydrotropism 364
- Hydroxocobalamin 343
- Hydroxy acids chelation by 4/9
- 3-Hydroxyanthranilic acid 338-339
- p-Hydroxybenzoic acid 444
- a-Hydroxybutyric acid 22
- 5-Hydroxy-N-dimethyl-tryptamine 974
- 2-Hydroxy-5-furan-carboxylic acid 181
- 3-Hydroxykynurenine 263 267 338
- Hydroxylamine 245 249
- Hydroxylamine reductase 245
- Hydroxymethyl ethyl ketone 123
- β -Hydroxy- β -methylglutaryl coenzyme A 161
- p-Hydroxyphenylacetic acid 186
- 2-Hydroxyphenylacetic acid 174
- p-Hydroxyphenylpyruvic acid 263
- Hydroxyproline 257 258 268
- 8-Hydroxyquinoline 315 448
see also Oxine
- 8-Hydroxyquinoline 5-sulfonic acid 481
- 5-Hydroxytryptophan 263 2/4
- Hypominitrite 45
- Hypophosphite 301
- Hyposulfite 05
- Hypoxanthine 275 2 6 3 0 397 402
- Imidazole acetol phosphate 271
- Imidazole glycerol phosphate 2/1
- Imidazoles 484
- Imidazoles 447 448 451 461 462
- Indigo 80 81
- Indole 263 271-272 280 449
- Indoleacetaldehyde 264
- Indoleacetic acid 263-264 361
- Indolepyruvic acid 264
- Indole serine ligase 271 309
- Infrared radiation 432 435
- Inoculum 1 3
- Inoline 276
- Inosine 5-phosphate 276
- Inositol 47 323 326 329 339-340
- Inulase 119
- Inulin 68-73 6
- Invertase *see* Sucrase

- Fructose 6 phosphate 213 218
 Fumarase 223
 Fumaric acid biosynthesis of 142 227
 formation of 133 141-143
 metabolism of 138 223 267 263 266
 269
 utilization of 68-73
 Fumaryl DL alanine 138 142 280
 Fungicidal action 447
 Fungistatic action 443-447
 Fungisterol 49
 Fungitoxic action and structure 451-
 465 477
 Fungitoxin effect of on mitosis 411
 effect of on morphology 441
 Furan derivatives 180
 Furfural 397 453
 Fusatic acid 280 281 283 415
 Fusarubin 170

 Galactocarolose 42 43
 Galactogen 119
 Galactonic acid 132 133
 L Galactono γ lactone 180
 D Galactose utilization of 61-62 68-73
 a Galactosidase 109 110
 β Galactosidase 87-88 108
 D Galacturonic acid 65
 Gallic acid 165 175 176
 Gallium 313
 Gammexane 340
 Gangaleoidin 170
 Gelase 170
 Gelatin 251
 Gentianose 109
 Gentyl alcohol 165 172-173
 Geodin 183
 Geotropism 357
 Germ tube growth 388 391 410
 Gladiolic acid 166 172
 Gladin 251
 Glutoxin 280 283 42
 Glutoxin monoacetate 280
 Gluconic acid formation of 131-135
 137 138 141
 Z keto 65 132 221
 metabolism of 208 220-221
 utilization of 63 68-73
 D Glucono- δ lactone 220
 Glucosamine 121
 Glucosamine 6 phosphate 302
 Glucose effect of concentration 85-86
 metabolism of 208 213 220
 utilization of 60 68-73
 Glucose aerodehydrogenase 135 140-
 221 229
 Glucose dehydrogenases 220
 Glucose oxidase *see* Glucose aerodehy-
 drogenase
 Glucose 6 phosphate 204 213 217-218
 Glucose 6 phosphate dehydrogenase 201
 217 218
 β Glucosidases 108 109 110 471
 Glucosone 121
 Glucuronic acid 65 132 135
 Glutaconic acid 133 146
 Glutamic acid assimilation of 259 480
 metabolism of 225 260 267 264 266-
 268
 requirement for 248
 utilization of 256 257
 Glutamic decarboxylase 265
 Glutamic dehydrogenase 262 266 309
 Glutamic γ semialdehyde 263
 Glutamine 258 265 268
 ester of 280
 γ Glutamyl transferase 256
 Glutaric acid 134 146
 Glutathione 186 253-254 470 481
 Glyceraldehyde 3 phosphate *see* Triose
 phosphates
 Glycerol 123 214 469
 utilization of 65 68-73
 Glycerol phosphates 123 302
 Glycerophosphatase 153 302
 Glycine copper complex of 466-467
 metabolism of 143 257 260 265 269
 toxicity of 258
 Glycogen 42 43 121 365
 Glycogens utilization of 68-73 6
 Glycolaldehyde 173
 Glycolic acid 138 143 269
 Glycols 123
 Glycylglycine dipeptidase 253
 Glyoxylic acid formation of 134 145
 145
 metabolism of 138 145 226 227 269
 278
 Grisofolm 164
 Grisofolium 183 441

- Malachite green 4/5 4/6
 Malate synthetase 227
 Malic acid 316 466-467
 formation of 133 140-143
 metabolism of 138 222 223 225 227
 utilization of 68-73 81
 Malic dehydrogenase 2 3 227
 Malic enzyme 149
 Malonic acid 81 146 224 225 2 6
 Maltose 66 68-73 107 108
 Manganese 314 315
 physiological effects of 139 310 310
 4/9 480
 requirement for 310
 Manganese dioxide 203
 Mannans 43 114
 Mannitol 64 68-73 122 212
 Mannose 63
 Mannonic acid 132 133
 Mannose 60-62 68-73
 Mannose 1 phosphate 62
 Mannosidases 110
 Mannosidostreptomycin 110
 Mappin 274
 Marine fungi 16 96
 Media culture 4
 Melanins 263 377 428
 Melanin 109
 Melibiose 109
 Melibiose 67 68-73 109
 Menadione 344
 Menadione reductase 177
 Mercaptide formation 463 4 0
 Mercaptobenzothiazole 472
 Mercury toxicity of 307 443 446 448
 463 465-472
 Metaphosphates 38 302
 Methanol 83
 Methionine metabolism of 181 186
 261 270 303-306 327 341
 toxicity of 2 8
 utilization of 13 7
 p-Methoxy tetrachlorophenol 183
 4-Methoxy 2,5-p-toluquinone 167
 Methylamyl ketone 154
 Methylarsenate sodium 182
 Methylation 181-182 341
 2-Methyl 5-bromomethyl-6-aminopyr-
 dine 330
 2-Methylbutene 2 163 415
 Methyl cinnamate 163 1/2
 β -Methylcrotonaldehyde 161
 β -Methylcrotonyl coenzyme A 161
 S-Methylcysteine 2 0
 γ -Methylcytosine 182
 Methyl-diethylarsine 182
 Methylene blue 204 03 469 4/3
 2-Methylenebis (3,4,6-trichlorophenol)
 464-465
 Methylene sulfide 189
 α -Methylglucoside 63 110
 β -Methylglucoside 65
 Methylheptenone 161-162
 4-Methyl 5- β -hydroxyethylthiazole 330
 2-Methyl 3-hydroxy 1,4-naphthoquinone
 344
 N-Methylisoleucine 966
 Methyl ketones 154
 α -Methylmannoside 110
 Methyl mercaptan 182
 Methyl p-methoxybenzoic acid 402
 Methyl p-methoxycinnamic acid 1/2
 2-Methyl 1,4-naphthoquinone 344 4/3
 Methylnonyl ketone 154
 Methylpropyl ketone 154
 D- α -Methylserine 266
 γ -Methylketonic acid 180
 N-Methylvaline 266
 Mineolitic acid 47
 Mineral oil 4
 Minerals 37-38 57 300-322
 and reproduction 310-371
 and spore germination 401
 Molybdenum 244-245 310-311 314 31
 Monoallyl phosphoric acid 1,3 302
 Monomethylamine 466
 Monomethylaminoethanol 181
 cis-cis-Muconic acid 174
 Mucopolysaccharases 119-120
 Muscarine 273
 Mutarotase 220
 Mutation 429 432 434-435 443
 Mycelianamide 280
 Mycelium decomposition of in soil 43
 Mycodextran 43
 Mycogalactan 43
 Mycomycin 164
 Mycophenolic acid 166
 Mycorrhizal fungi 91 347
 Myokinase 203

- Iodoacetamide 391
 Iodoacetic acid 463
 Iodosobenzoic acid 463
 Ionization 454-457
 Ionizing radiation 433-435
 β Ionone 161-162
 Iron 38 307-308 313
 physiological effects of 139 172-173
 304-308 336-337 340
 Isoamyl alcohol 262
 Isoamylamine 273
 Isocitric acid 138 143 145 223 225 227
 Isocitric dehydrogenase 223 226
 Isocitritase 226 227
 D Isoleucine 261
 DL Isoleucine 257
 L Isoleucine 270
 Isomaltose 102 107
 Isonicotinic hydrazide 160
 Isopropyl alcohol 474
 Itaconic acid 134 138 144-145
 Itatartaric acid 134 145
 Itatartaric acid lactone 145

 Javanicin 168 170 475
 Junipal 181

 Keratin 84 241-252
 β keto acid decarboxylases 154
 α ketoadipic acid 271
 β ketoadipic acid 174
 α ketobutyric acid 262
 2 ketogluconic acid 65 132 221
 α ketoglutaric acid metabolism of 138
 223-225 262 266 267 377 423
 α ketoglutaric acid oxidase 2 6
 α ketosocaproic acid 240
 α ketosovaleric acid 342
 ketones α β unsaturated 477-478
 5 keto octadecanoic acid dodecyl ester
 47
 6 keto octadecanoic acid 47
 Kopic acid 178-179 219
 kynurenic acid 263
 kynurenine 267 338-339
 kynurenine transaminase 264

 Laccase 142-146
 Lachrymators 448
 Lactase 87-88 103
 Lactic acid biosynthesis of 136 215-
 216 228
 formation of 136 138 185
 utilization of 68-73
 Lactic dehydrogenase 136 230
Lactobacillus casei factor 342-343
 Lactobionic acid 139
 Lactose metabolism of 107 108
 utilization of 56 67 68-73 74 185
 Lag phase 6
 Lecanoric acid 169
 Lecithins 47 153
 D-Leucine 261
 DL Leucine 256 257
 L Leucine 259 260 262 240
 L Leucyl L proline anhydride 255
 Levansucrase 120
 Levomycin 254
 Lichenase 114 119
 Light 160 357-360 404-428
 Lignin 41 83 90
 Lignoceric acid 48
 Lignosulfonic acid 84
 Lime sulfur 472-473
 Linoleic acid 47 48
 Linolenic acid 48
 Lipases 152-153
 Lipids 82 374
 metabolism of 142-164
 of cells 35 37 41 45-49 392-393
 see also Fats Fatty acids Phospholipids
 Lipositol 340
 Lipoxidase 155
 Luminescence 26-27
 Luteic acid 43
 Lycopersin 24-25 266
 Lycopene 158 159 161
 Lycopanthin 149
 Lyophilization 4
 Lysergic acid 249 280
 α D Lysine 261
 α L Lysine 257 259 261-262 271
 β Lysine 266
 Lysozyme 453

 Magnesium of cells 37 38
 physiological effects of 19 139 306-
 307 313 316 371
 requirement for 304 306-307

- Penicillin 70
 Penicillic acid 180 47
 Penicillin 15 184-187 308 441
 Penicillin amidase 187
 Penicillinase 187
 Penicillin B 270
 Penicillic acids 187
 Pentaacetylglucose 64
 Pentonic acids 2
 Pentosans 42 43 77 113-114
 Pentose phosphates 217-218
 Pentoses metabolism of 143 208 221-222
 Peptidases 65 49
 see also Endopeptidases Exopeptidases
 Peptides 250-251 255-256 479
 Periodicity 367
 Permeability 706-707 397 442-443 449
 453-454 458-460 470
 to acids 80
 to cations 391
 to metals 468
 to vitamins 326 340
 Peroxidases 232
 Peroxides 433 443
 Persulfate 303
 pH effect of ammonium utilization on 247-248
 effect of organic acid utilization on 80 248
 effect of on growth 14-15 19-23 6
 effect of on metabolism 19 141 150
 178 208 337
 effect of on mineral nutrition 97
 312 316
 effect of on permeability 80 3 6
 effect of on reproduction 363 66
 effect of on spore germination 403-404 409
 effect of on substrate utilization 9
 81 44 246
 effect of on toxicity 454-457 468
 effect of on vitamin requirements 333 337 40
 effect of on zonation 3 9
 internal 2 23
 Phallin 7
 Phalloidine 23
 Phenanthraquinones 477
 o Phenanthroline 49 480
 Phenazine derivatives 205
 Phenol 83 441
 phenols oxidation of 174-176
 toxicity of 449 453 455 4 7 463 464
 phenolsulfatase 306
 phenothiazone 472
 Phenylacetic acid 173-174 186 208
 Phenylalanine 171 207-209 271-272
 Phenylethylamine 186 273
 a Phenylglycine 271
 Phenylmercuric acetate 461
 Phenylpyruvic acid 211-212
 Phomazurin 28
 Phosphatases 103 123 275 302-303
 Phosphate 22 249 301-304
 Phosphatides 46 47 153
 Phosphite 301
 Phosphoenolpyruvic transphosphorylase 213
 Phosphofructokinase 213
 Phosphogluconate oxidation pathway 19 217-220 221-222
 6 Phosphogluconic acid 217
 Phosphoglyceric acids 213 372
 Phosphoglyceryl kinase 213
 Phosphoglyceryl mutase 213
 Phosphohexoisomerase 213
 Phospholipases 153
 Phospholipids 152 339
 see also Phosphatides
 Phosphomonoesterases 123
 Phosphoproteins 38 44
 Phosphorus 8 139 143 301-304 469
 bioassay for 313 314
 of cells 37 38 316
 Phosphorylases 121
 Photoreduction oxidative 213 462
 Photoreactivation 433
 Phototaxis 360
 Phototropism 358 360-362 392
 Phthalic acid 72
 Phycomycetes physiological characteristics of 18 90 1 304-305 347 366
 Ithyic acid 301 302
 Phytin 339-340
 Ihytoene 159 161
 Phytofluene 1 9 161
 Icterococcalin 280
 Iimelic acid 334

- Myristic acid 48
 Nalgiolaxin 183
 Nalgiotensin 183
 Naphthoquinone derivatives 168 170
 344 415-477
 ω (2 Naphthylloxy) n alkylcarboxylic
 acids 461
 Neopyrithiazine 332
 Neurosporene 159 161
 Nicotinamide 337
 Nicotinic acid 326 327 328 337-339
 345
 Nicotinic acid esters 461
 Nidulin 169 170 183
 Nigera 43
 Nitrate 242-245
 Nitrate reductase 244-245
 Nitrite metabolism of 249 335
 physiological effects of 215 243 246
 utilization of 246
 Nitrite reductase 245
 m Nitroaniline 245
 Nitroaryl reductase 245
 Nitrobenzene 83
 Nitrogen total 12 36 43-44
 Nitrogen fixation 249
 Nitrogen mustard 432 443
 Nitrogen supply physiological effects of
 19 41 79 139 149 178-179 210
 247-248 275 369-370
 Nitrophenols 462-464
 p Nitrophenyl β glucoside 107
 p Nitrophenyl phosphate 302
 β Nitropropionic acid 280 81
 ω Nitrostyrene derivatives 418
 Notatin *see* Glucose acrodehydrogenase
 Nuclei of spores 389-390
 Nucleic acids 45 274-277
 Nucleosides 275-277
 Nucleotides 203 275-277

 Octadecadienoic acid 48
 Octadecanoic acid 48
 keto derivatives of 47
 Octadecatrienoic acid 48
 Octadecenoic acid 48
 Octanoic acid 154 155
 Oleic acid 48 82 335

 Oligosaccharides 66-75 105-109 363-
 369
 Oosporein 167
 Order of death at high temperature
 424-425
 Ornithine 261 267 268
 Ornithine cycle 268-269
 Orotic acid 277
 Orotidine 277
 Orotidine 5 phosphate 277
 Osmotic pressure 27-28 406
 Oxalacetic acid 138 141-143 223- 9
 330 333
 Oxalacetic decarboxylase 266
 Oxalic acid 248 379
 formation of 134 137 138 141 144
 145 180 303
 utilization of 68-73 81
 Oxalic acid decarboxylase 144
 Oxalosuccinic acid 223
 Oxalosuccinic decarboxylase 223 266
 β Oxidation 153-155
 Oxidation reduction balance 100-101
 Oxidative assimilation 212 462-463
 Oxidative phosphorylation 203 467
 Oxime 315 478 480-481
 see also 8 Hydroxyquinoline
 dl Oxibiotin 331-335
 Oxygen determination of 203
 diffusion of 205
 physiological effects of 23-24 208-
 209 365-366 391 412-413 435
 Oxygen demand 2-3
 Oxysavonin 170
 Oxytetracycline 183
 Ozone 366

 Palmitic acid 48 82
 Pantethine 161
 Pantoic acid 342
 Pantothenic acid 161 329 341-347 345
 Paraffins 82
 Partition coefficient 458-459
 Patulin 172-175 179-180 477
 Lectase *see* Pectin methyltransferase
 Lectic materials 41 80 114-119
 Lectinase *see* Pectin polygalacturonase
 Pectin depolymerase 116 118
 Pectin methyltransferase 115-119
 Pectin polygalacturonase 115-119

- Penatin 220
 Penicillic acid 180 477
 Penicillin 15 184-187 308 441
 Penicillin amidase 187
 Penicillinase 187
 Penicillin B 270
 Penicillic acids 187
 Pentaacetylglucose 64
 Pentonic acids 722
 Pentosans 42 45 47 115-114
 Pentose phosphates 217-218
 Pentoses metabolism of 143 208 221-222
 Peptidases 76, 4 9
 see also Endopeptidases Exopeptidases
 Peptides 250-251 253-256 4,9
 Periodicity 367
 Permeability 706-707 397 442-443 449
 453-454 458-460 4 0
 to acids 80
 to cations 331
 to metals 468
 to vitamins 3,6 340
 Peroxidases, 237
 Peroxides, 433 443
 Persulfate 300
 pH effect of ammonium utilization on 74, 248
 effect of organic acid utilization on 80 748
 effect of on growth 14-1 19-23 26
 effect of on metabolism 19 141 1,0 1 8 208 337
 effect of on mineral nutrition 207 312, 316
 effect of on permeability 80 3,6
 effect of on reproduction 263 266
 effect of on spore germination 403-404, 403
 effect of on substrate utilization 9 81 214, 216
 effect of on toxicity 4 4-4,7 463
 effect of on vitamin requirements 330 337 340
 effect of on zonation 2
 internal 22 23
 Phallin, 2,
 Phalloidine, 2,
 Phenanthraquinones 4 7
 o phenanthroline 4,9 480
 Phenazine derivatives 200
 Phenol 83 441
 Phenols oxidation of 1,4-1 6
 toxicity of 449 453 450 457 463 464
 Phenolsulfatase 306
 Phenothiazone 4,2
 Phenylacetic acid 173-1,4 186 208
 Phenylalanine 1,1 257-259 271-272
 Phenylethylamine 186 2,3
 α Phenylglycine 771
 Phenylmercuric acetate 46
 Phenylpyruvic acid 2,1-2 7
 Phomazarin 283
 Phosphatases 100 123 2,3 30, 303
 Phosphate 22 749 201- 04
 Phosphatides 46 47 103
 Phosphate 301
 Phosphoenolpyruvic transphosphorylase 213
 Phosphofructokinase 213
 Phosphoglucuronate oxidation pathway 1 9 217-220 221 222
 6 Phosphogluconic acid 21,
 Phosphoglyceric acids 213 3 2
 Phosphoglyceryl kinase 713
 Phosphoglyceryl mutase 213
 Phosphohexosomerase 713
 Phospholipases 153
 Phospholipids 152 3 9
 see also Phosphatides
 Phosphomonoesterases 123
 Phosphoproteins 78 44
 Phosphorus 8 139 149 201-204 469
 bioassay for 313 314
 of cells 37 3 3 6
 Phosphorylases 171
 Phosphorylation oxidative 213 462
 Photoreaction 433
 Phototaxis 260
 Phototropism 3 8 60-62 7
 Fthalic acid 22
 Phycomycetes physiological characteristics of 8 20 21 204-30, 317 266
 Phytic acid 201 262
 Phytin 339-240
 Phytoene 1 9 161
 Phytofluene 1,2 161
 Picroxcelin 200
 Limelic acid 3,4

- Ribose 64 68-73
 Ribose 1 phosphate 16
 Ribose 5 phosphate 218
 Ribulose 5 phosphate 218
 Rubber 84-85
 Rubidium 303 316
 Rubrofusarin 1 0
 Rugulose 43
 Rust fungi 26 3/8 426
 spore germination in 390 397 398
 399-401 403 405 408 412 414

 D-Saccharic acid 60 132 130
 Saponin 57
 Sarcosine 266
 Sawdust as culture medium 3
 Scandium 313
 Sclerotia 306-307
 Sclerotiose 43 1-1
 Sedoheptulose phosphates 18
 Selenate sodium 182
 Selenide sodium 180
 Selenium 182 4/5
 Self inhibition 413-415
 Serine 47 143 157 26 269 271
 Serine dehydrase 262
 Shaker 2
 Shikimic acid 171 1/2 271-2 2 341
 Silicon 38 313
 Silver toxicity of 446 447 450 453
 460-472
 uptake of 402 468-472
 Sodium 38 300 303
 Soil as culture medium 3
 in preservation of cultures 4
 Soil fungi 23 247
 Solanin 441
 Solanone 1,0
 Solubility 458-4 9
 Sonication 200
 Sorbitol 68-73 122
 Sorbose 122 90
 toxicity of 10 62-63
 utilization of 62 68-73
 Spectrophotometry 04
 Spiculisporic acid 47
 Spinosin 167
 Spirilloxanthin 159
 Spore discharge 364- 65
 Spore germination effect of ionizing ra-
 diation on 433-430
 effect of light on 428
 effect of pH on 22 403-404
 effect of temperature on 18 407-410
 latent period of 388 391 410-411
 rate of 388 390-391 395-396 400
 410-411
 Spore starch 42
 Spores maturation of 90-398
 nuclear number of 389-390
 respiration of 393-390
 selling of 391 406 413
 Sporulation effect of calcium on 312
 effect of zinc on 309
 inhibition of 481
 Squalene 157
 Stachyose 109
 Staling 8 249
 Stannic chloride 467
 Stannous chloride 467
 Starch 68-73 75-76
 Stearic acid 48 82
 Sterilization chemical 3,7-378
 Steroids requirement for 340
 Sterols 46 49 82 155-107
 Stipitatic acid 169
 Streptolisin 04
 Streptomycin 110 160 108 308 441 452
 Streptothricin 254
 Strontium 311
 Structure and fungitoxicity 464-465
 Styrene 4 8
 Succinate ketoglutarate isocitrate cycle
 224
 Succinic acid metabolism of 142 145
 146 2 3 220-223 428
 formation of 133 142-143
 utilization of 68-73 81
 Succinic dehydrogenase 223
 Succinoxidase 230
 Sucrose 105-107 374 393 471
 Sucrose 3 4
 breakdown of 105-107 393
 utilization of 66-74 369
 Sucrose phosphorylase 107
 Sugar alcohols 56 122
 Sugar fungi 91
 Sugar phosphates hydrolysis of 100 302
 Sugars decomposition of 57-58

- Pinoselic acid 168
 Pipelic acid 262
 Plant pathogenic fungi 19 21 115 347
 366 377-378
 Plasmoptysis 403
 Polyenes 164
 Polyenyne 164
 Polyfructoses of mycelium 43
 Polygalactoses of mycelium 43
 Polyglucoses of mycelium 43
cis Polyisoprene 164
 Polymyxin 453
 Polyphenol oxidases 84 173 174-176
 178 263 480
 Polyphosphates 302 479
 Polyporenic acids 158
 Polysaccharides 42-43 120-121 368-369
 Polyuronide of mycelium 43
 Porphyrins 479
 Potassium 303-304 316 371 468
 bioassay for 314
 of cells 37 78
 permeability to 468
 Powdery mildew fungi spore germina-
 tion in 396 398 399 401 403-406
 408 412
 Prephenic acid 271-272
 Preservation of cultures 4-5
 Pressure 423 433
 Progesterone 157
 Proline 257 260 265 268
 β Propiolactone 454
 Propionic acid 68-73 81 469
 Propylene oxide 377-378
 Proteinases 251-253
 Protein of cells 57 41 44-45 393
 requirement for 345
 utilization of 250-252
 Proteinases *see* Endopeptidases
 Protocatechuic acid 171-172 174
 Protopectinase 115-116
 Pteroylglutamic acid 341-343
 Thiocol 344
 Puberulonic acid 169
 Pulcherrimin 283 309
 Purines 245-248 462
 Putrescine 273
 Pyrazine derivatives 283
 Pyrazoles 464
 3 Pyridinethiol 472
 Pyridoxal 335
 Pyridoxal phosphate 267 271 335
 Pyridoxamine 335
 Pyridoxine 326 328 329 335 345
 Pyrimidine moiety of thiamine 330-339
 Pyrimidines 275-277
 Pyriothiamine 332
 Pyrone derivatives 172 178-180
 Pyrophosphate 38 302
 Δ Pyroline 5 carboxylic acid 268
 Pyruvic acid 379 397
 carboxylation of 142 228 333
 from serine 143 262
 role of in respiration 138 213-216
 218 222-228
 in valine synthesis 270
 Pyruvic carboxylase 213 216 309
 Pyruvic oxime 249
- Q* of growth 15
 of respiration 208
 of thermal death 475
 Quinic acid 171 172
 Quinoline derivatives 283
 Quinolinic acid 338-339
 Quinone reductase 177
 Quinones in respiration 177-178
 toxicity of 449 463 475-478
- Raffinose 67-73 107 109
 Ravenelin 168
 Resistance 378
 Respiration aerobic 202
 anaerobic 202
 endogenous 210-211 463
 inhibition of 442-443 484
 of spores 343-39
 terminal 272 232
 Respirometry 203
 Resting cells 101
 Resting spores 396-398
 Rhamnose 63 68-73
 Rhizopterin 342-343
 Rhodoviolascins 178 179
 Riboflavin 307 326 328 396-397 343
 372 402
 Riboflavin monophosphate 272
 Ribonucleic acid 47 275

- s* Triazines 441-452 453 457 463 464
 Tri *n* butyltin acetate 467
 N (Trichloromethylthio) 4 cyclohexene
 1, *m* dicarboxamide 44
 Trimethylamine 273
 Trimethylarsine 182
 Trimethylethylene 163 41, *s*
 Triose phosphate dehydrogenase 204
 213 484
 Triose phosphate isomerase 913
 Triose phosphates 213 218 221
 Tripalmitin 82
 Triphenylmethanes 464 475
 Triphosphate 302
 Triphosphopyridine nucleotide 04
 17-218 223 229 245 267 338
 Tristearin 87
 Triterpenes 157-158
 Tropisms 391-392
 Tropolone derivatives 169 170 475
 Trypsin 2 7
 Trypsinogen 5
 Tryptophan 171 56 457 259
 metabolism of 263 271-2 2 338-339
 Tryptophan synthetase 271 309
 Tungstate sodium 441
 Turbidity as measure of growth 12
 Tyramine 263 273
 Tyrosinase 175-176 263 316
 Tyrosine 171 257 59 263 271-272

 Ultraviolet radiation 429-433
 effect of on reproduction 357-358
 59 316
 Uracil 475 216
 Uranium 448 454 4 0
 Urea 45 249 277-279
 metabolism of 68- 69 6-273
 Urease 278 477
 Ureidosuccinic acid 277
 Urethane 83 49
 γ (3,4 Ureylene-cyclohexyl)butyric acid
 335
 Uric acid 275 78- 79
 Uricase 278
 Uridine 5 phosphate 277
 Urocanic acid 264
 Ustilagic acid 47 104 219
 Ustilic acids 47

 Utilization definitions of 55- 6

 Valine 186 257 261 270
 Vanadium 313
 Vanillin 174
 Vapor pressure 458-459
 Varianose 43
 Variation 3 102
 Versene *see* Ethylenediaminetetraacetic
 acid
 Viomycin 254
 Viridicatin 281 283
 Vitamin A 345
 Vitamin B *see* Thiamine
 Vitamin B₁₂ *see* Riboflavin
 Vitamin B *see* Pyridoxine
 Vitamin D 345
 Vitamin E 345
 Vitamins B₁₂ 343-344
 Vitamins K 344 46
 Vitamins as impurities 57
 effect of on reproduction 371-372
 effect of on spore germination 407
 D-Volemitol 122
 Volutin 38

 Water and growth 25-46
 see also Bound water Humidity
 Water content of cells 36
 Wilt fungi 18-19 23
 Wood destroying fungi 8 83-84 175
 347

 Xanthates 47, *s*
 Xanthine 275 276 78
 Xanthine oxidase 275 278
 Xanthenes 168 170
 Xanthophylls 162
 Xanthurenic acid 263
 X rays *see* Ionizing radiation
 Xylanases 114
 Xylans 114
 Xylobiose 114
 Xylonic acid 132 136
 Xylose 63 68-73 421
 Xylulose 5 phosphate 218

 Yeast extract 375

 Zeaxanthin 159

- Sulfanilamide 2/6 442 461
 Sulfate 182 186 304-306
 Sulfhydryl compounds 11 393
 Sulfhydryl enzymes inhibition of 462-463 470-472 476-478 485
 Sulfide 303
 Sulfite 303
 Sulfonamides 341 456 461
 Sulfur 8 139 182
 requirement for 304-306
 toxicity of 450 472-475
 Sulfur mustard 443
 Sulochrin 166
 Surface action 453-454
 Surface tension 458-459

 Tannase 1 6
 Tannic acid 83-84 175
 Tannins 83
 Tartaric acid 68-73 81 131 143
 Taurocholate sodium 441
 Taxonomy 39-41 9 90-91 103 346-347
 Telemorphosis 3/4
 Tellurite potassium 182
 Tellurium 182 473
 Temperature and vitamin requirements 326 336
 and zonation 359
 effect of on growth 13-19
 effect of on metabolism 79 208
 effect of on pH response 19
 effect of on reproduction 359 362-364
 effect of on spore germination 389 397 399 407-412
 lethal effects of 423-427 433
 trophic effects of 362
 Terminal respiration 229-232
 Terresin 183
 Terresinic acid 180
 Tetrachloro *p* benzoquinone 4/6
 Tetrachloronitrobenzenes 419
 Tetracosanoic acid 48
 Tetracosenoic acid 48
 Tetracycline 183
 Tetradecane 82 163
 Tetradecanoic acid 48
 Tetrahydrophytoene 161
 Tetramethylthiuram disulfide 482-483
 Tetramethylthiuram monosulfide 483
 Tetric acid 180
 Thelephoric acid 167
 Theophylline 2/6
 Thermal death point 423-424
 Thermal death time 424
 Thermophily 18
 Thiaminase 329
 Thiamine 13 215 326 328 329-337
 333 339 345 347 371-372
 Thiamine pyrophosphate 329-330
 Thiazole moiety of thiamine 330-332
 Thigmotropism 392
 Thiobis phenols 472
 Thiocyanate 305
 Thiocyanate derivatives 472
 Thioglycolic acid 477
 Thiol compounds protection by 463 470 476-477
 Thiolutin 280
 Thionophosphate 302
 Thiosulfate 305 434
 Thiourea 184 245 304 305 4/2
n-Threitol 122
 Threonine 257 262 264 2/0
n-Threose 64
Thuyaphycin 170 475
 Thunberg method 202-203
 Thunberg Wieland reaction 147 143 227
 Thymine 275
 Thymol 441
 Tin 467
 Tolerance to toxicants 448-449
 Toluquinone 173
 Torularhodin 159
 Torulene 159
 Toxicity apparent 451
 specific 451
 Transacetylase 143
 Transaldolase 218
 Transamination 258 267
 Transfructosidase 106-107
 Transgalactosidase 107 108
 Transglucosidase 107 108
 Transglycosidases 107-108
 Transketolase 218 279
 Transmethylation 181-187
 Trehalase 108
 Trehalose 66 68- 5 108-109

Zein 251

Zinc 38 308-309

induction of variants by 443

permeability to 468

physiological effects of 139 142 157

1,3 203 2,5 308-309 320 314 371

toxicity of 446

Zonation 18 308-359

Zygotropism 374

Zymonic acid 181

